

PCT

From the INTERNATIONAL BUREAU

**NOTIFICATION OF THE RECORDING
OF A CHANGE**

(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

To:

CHARI, Santosh, K.
Orange & Chari, Suite 4900, P.O.
Box 190
66 Wellington Street West
Toronto Dominion Bank Tower
Toronto-Dominion Centre
Toronto, Ontario M5K 1H6
CANADA

Date of mailing (day/month/year) 02 November 2000 (02.11.00)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference 00264-0003	
International application No. PCT/CA00/00430	International filing date (day/month/year) 20 April 2000 (20.04.00)

1. The following indications appeared on record concerning: <input type="checkbox"/> the applicant <input type="checkbox"/> the inventor <input checked="" type="checkbox"/> the agent <input type="checkbox"/> the common representative		
Name and Address CHARI, Santosh, K. Orange & Chari 4900-55 King Street West, P.O. Box 190 Toronto Dominion Bank Tower Toronto-Dominion Centre Toronto, Ontario M5K 1H6 Canada	State of Nationality	State of Residence
	Telephone No. 416 601-8440	
	Facsimile No. 416 601 8454	
	Teleprinter No.	
2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning: <input type="checkbox"/> the person <input type="checkbox"/> the name <input checked="" type="checkbox"/> the address <input type="checkbox"/> the nationality <input type="checkbox"/> the residence		
Name and Address CHARI, Santosh, K. Orange & Chari, Suite 4900, P.O. Box 190 66 Wellington Street West Toronto Dominion Bank Tower Toronto-Dominion Centre Toronto, Ontario M5K 1H6 Canada	State of Nationality	State of Residence
	Telephone No. 416 601-8440	
	Facsimile No. 416 601 8454	
	Teleprinter No.	
3. Further observations, if necessary: 		
4. A copy of this notification has been sent to: <input checked="" type="checkbox"/> the receiving Office <input checked="" type="checkbox"/> the designated Offices concerned <input type="checkbox"/> the International Searching Authority <input type="checkbox"/> the elected Offices concerned <input type="checkbox"/> the International Preliminary Examining Authority <input type="checkbox"/> other:		

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer S. De Michiel Telephone No.: (41-22) 338.83.38
--	--

THIS PAGE BLANK (USPTO)

PCT

**NOTIFICATION OF THE RECORDING
OF A CHANGE**

(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

GRAVELLE, Micheline
Bereskin & Parr
40th floor
40 King Street West
Toronto, Ontario M5H 3Y2
CANADA

Date of mailing (day/month/year) 14 June 2001 (14.06.01)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference 6850-226	
International application No. PCT/CA00/00430	International filing date (day/month/year) 20 April 2000 (20.04.00)

1. The following indications appeared on record concerning:	
<input type="checkbox"/> the applicant	<input type="checkbox"/> the inventor
<input checked="" type="checkbox"/> the agent	<input type="checkbox"/> the common representative
Name and Address CHARI, Santosh, K. Orange & Chari, Suite 4900, P.O. Box 190 66 Wellington Street West Toronto Dominion Bank Tower Toronto-Dominion Centre Toronto, Ontario M5K 1H6 Canada	State of Nationality State of Residence Telephone No. 416 601-8440 Facsimile No. 416 601 8454 Teleprinter No.
2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:	
<input checked="" type="checkbox"/> the person	<input checked="" type="checkbox"/> the name
<input checked="" type="checkbox"/> the address	<input type="checkbox"/> the nationality
<input type="checkbox"/> the residence	
Name and Address GRAVELLE, Micheline Bereskin & Parr 40th floor 40 King Street West Toronto, Ontario M5H 3Y2 Canada	State of Nationality State of Residence Telephone No. 416 957-1682 Facsimile No. 416 361-1398 Teleprinter No.
3. Further observations, if necessary: A new agent has been appointed. Please also note the new agent's file reference.	
4. A copy of this notification has been sent to:	
<input checked="" type="checkbox"/> the receiving Office	<input type="checkbox"/> the designated Offices concerned
<input type="checkbox"/> the International Searching Authority	<input checked="" type="checkbox"/> the elected Offices concerned
<input checked="" type="checkbox"/> the International Preliminary Examining Authority	<input checked="" type="checkbox"/> other: CHARI, Santosh, K.

The International Bureau of WIPO 34, chemin des Colombettes 1211 Genève 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer F. Baechler Telephone No.: (41-22) 338.83.38
--	--

THIS PAGE BLANK (USPTO)

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner
US Department of Commerce
United States Patent and Trademark
Office, PCT
2011 South Clark Place Room
CP2/5C24
Arlington, VA 22202
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year)

04 December 2000 (04.12.00)

International application No.

PCT/CA00/00430

Applicant's or agent's file reference

00264-0003

International filing date (day/month/year)

20 April 2000 (20.04.00)

Priority date (day/month/year)

23 April 1999 (23.04.99)

Applicant

FORSBERG, Cecil, W. et al

1. The designated Office is hereby notified of its election made:



in the demand filed with the International Preliminary Examining Authority on:

09 November 2000 (09.11.00)



in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was



was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Facsimile No.: (41-22) 740.14.35

Authorized officer

Nestor Santesso

Telephone No.: (41-22) 338.83.38

THIS PAGE BLANK (USPTO)

REC'D 31 JUL 2001

WIPO

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference 00264-0003	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/CA00/00430	International filing date (day/month/year) 20/04/2000	Priority date (day/month/year) 23/04/1999
International Patent Classification (IPC) or national classification and IPC A01K67/027		
Applicant UNIVERSITY OF GUELPH et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 7 sheets, including this cover sheet.

- ☒ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 7 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☒ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 09/11/2000	Date of completion of this report 27.07.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer SCHEFFZYK, I Telephone No. +49 89 2399 8602 

THIS PAGE BLANK (USPTO)

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/CA00/00430

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

Description, pages:

1-56 as originally filed

Claims, No.:

1-57 with telefax of 04/07/2001

Drawings, sheets:

1/58-58/58 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:

THIS PAGE BLANK (USPTO)

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/CA00/00430

☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

IV. Lack of unity of invention

1. In response to the invitation to restrict or pay additional fees the applicant has:

- ☐ restricted the claims.
☐ paid additional fees.
☐ paid additional fees under protest.
☐ neither restricted nor paid additional fees.

2. ☐ This Authority found that the requirement of unity of invention is not complied and chose, according to Rule 68.1, not to invite the applicant to restrict or pay additional fees.

3. This Authority considers that the requirement of unity of invention in accordance with Rules 13.1, 13.2 and 13.3 is

- ☐ complied with.
☒ not complied with for the following reasons:
see separate sheet

4. Consequently, the following parts of the international application were the subject of international preliminary examination in establishing this report:

- ☒ all parts.
☐ the parts relating to claims Nos. .

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	4,6,9,11,12,14-16,18-28,30-48
	No:	Claims	1-3,5,7,8,10,13,17,29,49-57
Inventive step (IS)	Yes:	Claims	4,6,9,11,12,14-16,18-28,36-43,48
	No:	Claims	1-3,5,7,8,10,13,17,29-35,44-47,49-57

THIS PAGE BLANK (USPTO)

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/CA00/00430

Industrial applicability (IA) Yes: Claims 1-18,29-57
 No: Claims 19-28

2. Citations and explanations
 see separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:
see separate sheet

THIS PAGE BLANK (USPTO)

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/CA00/00430

SECTION IV-----

The IPEA is of the opinion that present claims are directed to different problems which are not linked together by a single inventive concept as required by Rule 13.1-13.3 PCT:

The first problem which can be identified is the provision of animals containing/expressing a phytase gene and constructs suitable to prepare such animals which are selected from the group consisting of pigs, goats, sheeps, cows, horses, fish and poultry (claims 13-18,26-28,33,34,36-43, 44-57).

The second problem can be seen in the provision of a method suitable for expressing a protein in the gastrointestinal tract of pigs, goats, sheeps, cows, horses, fish and poultry (claims 19-25.29) and, finally

the third problem which can be identified is the provision of a method suitable for producing a protein via salivary gland secretion in any non-human animal (claims 30-32 and 35)

The only common concept underlying the subject-matter of all claims can be seen in the use of non-human animals to express polypeptides. This concept, however, is not considered suitable to establish unity of presently claimed subject-matters since said concept is already well-known in the prior art. Therefore, the IPEA holds the view that present application does not meet the requirements of Rule 13.1-13.3 PCT.

SECTION V-----

Novelty:

Present claims 1-3, 5,7,8, 10, 13, 17, 53 and 54 do not contain a **technical feature** which would be suitable to render the claimed animal/animal cells clearly distinguishable from any natural occurring ones. Concerning claims 5 and 7 it is noted that the expression "derives from" does not convey an exact meaning but encompasses any PSP or PRP since eventually any sequence may be considered to be derived from another sequence. With respect to claims 13, 17, 29, 53 and 54 which require that the gene encodes phytase it is noted that according to the

THIS PAGE BLANK (USPTO)

information given in the application on page 1, lines 23 and 24 cows express phytase at a low level. Thus, claims 1-3,5,7,8,10,13, 17, 29, 53, 54 do not meet the requirements of Art. 33(2)(3) PCT.

In addition, claims 49-52, 55-57 also lack novelty and inventive step since phytase isolated from E.coli as well as antibodies directed against said enzyme are already taught in WO 99/08539 (1). Crossreactions thereof with the phytase expressed by a nucleic acid sequence according to SEQ.ID.NOS. 3, 5 and 7 cannot be excluded. For the sake of completeness it is noted that even if presently claimed antibodies were novel over those available in the prior art the presence of an inventive step still could not be acknowledged since APPA of E.coli already was available at the filing date of present application. The provision of antibodies directed to a known protein, however, lacks inventive activity since it only involves standard methods.

Inventive step:

Claims 44-47 concern nucleic acid constructs containing a gene encoding phytase and one regulatory sequence for gastrointestinal tract specific expression of said gene. However, taking into account that the preparation of such constructs and the expression thereof in salivary glands is already suggested in WO 97/48812 (2) (see page 18, lines 27-31) the subject-matter of claims 44-47 and claims 30-35 which relate to a process for producing a polypeptide which involves the expression of said polypeptide in salivary glands cannot be considered to be inventive. Concerning claims 30-35 it is noted that it appears that the inventive step establishing feature resides in the unexpected successful expression of a polypeptide in a specific animal (see below). However, claims 30-35 do not specify the non-human animals used for expression of the protein. Thus, they do not contain a feature suitable to establish the presence of an inventive step.

According to the Applicant it is not possible to make reliable predictions concerning the successful expression of polypeptides in the saliva or gastrointestinal tract of non-human animals without demonstrating the gene function in the animal, in particular when the non-human animal used to express the polypeptide is larger. The IPEA basically concurs with the Applicant in this position. However, with respect to this the IPEA would like to point out that present

THIS PAGE BLANK (USPTO)

application only exemplifies the expression of APPA in pig! Thus, for obvious reasons present claims cover subject-matter which is evidently not enabling disclosed in present application and hence unworkable (Art. 5 and 6 PCT).

SECTION VII-----

- 1). Newly-filed set of claims extends beyond the content of the application as originally filed since according to the application as filed the transgene constructs had to be heterologous. However, according to new claims said constructs also can be homologous (Art. 34(2)(b) PCT).
- 2). In addition, no basis can be found in the application as originally filed for the subject-matter of claim 42 (Art. 34(2)(b) PCT).
- 3). With respect to the expression "incorporated by reference" Applicant's attention is drawn to Guidelines C-II 4.4 and 4.17 PCT.

SECTION VIII-----

- 1). It is highly questionable whether all animals specified in claims 1, 13 and 36 actually are suitable to successfully express functional polypeptides (see also above, section V).
- 2). The language used in claim 48 ("includes") creates ambiguity since it is unclear whether the molecule according to claim 44 contains one of the sequences recited in claim 48 additional or not.
- 3). Claims 19-28 relate to subject-matter considered by this Authority to be covered by the provisions of Rule 67.1(iv) PCT. Consequently, no opinion will be formulated with respect to the industrial applicability of the subject-matter of these claims (Article 34(4)(a)(i) PCT).

THIS PAGE BLANK (USPTO)

OCT 24 2000

**THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE
PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:**

1. A transgenic non-human animal that carries in the genome of its somatic and/or germ
5 cells a nucleic acid sequence including a heterologous transgene construct, said construct
including a transgene encoding a protein, said transgene being operably linked to a first
regulatory sequence for salivary gland specific expression of said protein.
2. The animal of claim 1 wherein said first regulatory sequence comprises a saliva
10 protein promoter/enhancer sequence, whereby said animal expresses said protein in its saliva.
3. The animal of claim 1 wherein said animal is a mammal.
4. The animal of claim 3 wherein said animal is chosen from the group comprising pigs,
15 goats, sheep, cows, horses, rabbits, rodents, cats and dogs, and in addition, fish and poultry..
5. The animal of claim 1 wherein said saliva protein promoter/enhancer sequence
comprises a parotid secretory protein (PSP) promoter/enhancer, a proline-rich protein (PRP)
promoter/enhancer or a salivary amylase promoter/enhancer.
20
6. The animal of claim 5 wherein said promoter/enhancer is a parotid secretory protein
(PSP) promoter/enhancer.
7. The animal of claim 6 wherein said parotid secretory protein (PSP)
25 promoter/enhancer is derived from a mouse.
8. The animal of claim 5 wherein said promoter/enhancer is a proline-rich protein (PRP)
promoter/enhancer.
- 30 9. The animal of claim 8 wherein said proline-rich protein (PRP) promoter/enhancer is
derived from a rat.

THIS PAGE BLANK (USPTO)

10. The animal of claim 1 wherein said transgene is further operably linked to one or more second regulatory sequences including enhancers, transcription regulatory sequences, termination sequences, and polyadenylation sites.

5 11. The animal of claim 1 wherein said transgene comprises a gene encoding a protein having phytase activity.

12. The animal of claim 1 wherein said transgene encodes a phytase or a homologue thereof.

10 13. The animal of claim 1 wherein said animal is a pig, said transgene comprising a gene encoding a protein having phytase activity and wherein said first regulatory sequence comprises a parotid secretory protein (PSP) promoter/enhancer or a proline-rich protein (PRP) promoter/enhancer.

15 14. The animal of claim 1 wherein said transgene construct comprises a nucleic acid sequence according to SEQ ID NO:3, SEQ ID NO:5; or SEQ ID NO:7.

20 15. A transgenic non-human animal that carries in the genome of its somatic and/or germ cells a nucleic acid sequence including a heterologous transgene construct, said construct including a transgene encoding phytase or a homologue thereof.

16. The animal of claim 15 wherein said transgene is operably linked to a first regulatory sequence for salivary gland specific expression of said phytase.

25 17. The animal of claim 16 wherein said first regulatory sequence comprises a parotid secretory protein (PSP) promoter/enhancer, a proline-rich protein (PRP) promoter/enhancer or a salivary amylase promoter/enhancer.

30 18. The animal of claim 17 wherein said animal is a mammal.

THIS PAGE BLANK (USPTO)

19. The animal of claim 18 wherein said phytase or a homologue thereof is expressed in saliva or in the gastrointestinal tract of said animal.

20. The animal of claim 15 wherein said transgene construct comprises a nucleic acid sequence according to SEQ ID NO:3, SEQ ID NO:5; or SEQ ID NO:7.

21. A method of expressing a protein, the method comprising the steps of:

a) introducing a transgene construct into a non-human animal embryo such that a non-human transgenic animal that develops from said embryo has a genome that comprises said

transgene construct, wherein said transgene construct comprises:

i) a transgene encoding said protein, and

ii) at least one regulatory sequence for gastrointestinal tract specific expression of said protein,

b) transferring said embryo to a foster female; and,

c) developing said embryo into said transgenic animal

wherein said transgene is produced in the gastrointestinal tract of said animal.

22. The method of claim 21 wherein said regulatory sequence provides for salivary gland or pancreatic gland specific expression of said protein.

23. The method of claim 21 wherein said regulatory sequence provides for salivary gland specific expression of said protein.

24. The method of claim 23 wherein said salivary gland is a parotid gland, submaxillary gland, or a submandibular gland.

25. The method of claim 23 wherein said transgene is expressed in the saliva of said animal.

26. The method of claim 21 wherein said transgene is heterologous.

THIS PAGE BLANK (USPTO)

27. The method of claim 21 wherein said at least one regulatory sequence comprises a salivary protein promoter/enhancer sequence.

28. The method of claim 21 wherein said protein is a glycoprotein.

29. A transgenic animal adapted for expressing a protein according to the method of claim 21, or a progeny thereof.

30. The method of claim 21 wherein said protein is a phytase or a homologue thereof.

31. The method of claim 21 wherein said transgene construct comprises a nucleic acid sequence according to SEQ ID NO:3, SEQ ID NO:5, or SEQ ID NO:7.

32. A process for producing a protein comprising the steps of:

a) obtaining saliva containing said protein from a non-human transgenic animal, said animal containing within its genome a transgene construct, wherein said transgene construct comprises:

i) a transgene encoding said protein, and

ii) at least one regulatory sequence for salivary gland specific expression of said protein, and

extracting said protein from said saliva.

33. The process of claim 32 wherein said transgene is heterologous.

34. The process of claim 32 wherein said at least one regulatory sequence comprises a salivary protein promoter/enhancer sequence.

35. The process of claim 32 wherein said protein is a glycoprotein.

36. The process of claim 32 wherein said transgene construct comprises a nucleic acid sequence according to SEQ ID NO:3, SEQ ID NO:5; or SEQ ID NO:7.

THIS PAGE BLANK (USPTO)

37. The process of claim 32 wherein said protein is a phytase or a homologue thereof.

38. The process of claim 32 wherein said salivary gland is a parotid gland, submaxillary, or a submandibular gland.

39. A method for expressing a phytase or a homologue thereof in a non-human animal, said method comprising:

a) constructing a nucleic acid sequence including a transgene construct comprising:

i) a transgene encoding said phytase or a homologue thereof, and

ii) at least one regulatory sequence for gastrointestinal tract specific expression of said protein, and

b) transfecting the animal with said nucleic acid sequence;

whereby said animal carries within the genome of its somatic and/or germ cells said transgene construct and wherein said animal expresses said phytase or a homologue thereof in its gastrointestinal tract.

40. The method of claim 39 wherein said transgene construct results in salivary gland or pancreatic gland specific expression of said phytase or a homologue thereof.

41. The method of claim 40 wherein said regulatory sequence provides for salivary gland specific expression of said phytase or a homologue thereof.

42. The method of claim 41 wherein said salivary gland is a parotid gland, submaxillary, or a submandibular gland.

43. The method of claim 41 wherein said phytase or a homologue thereof is expressed in the saliva of said mammal.

44. The method of claim 41 wherein said transgene construct comprises a nucleic acid sequence according to SEQ ID NO:3, SEQ ID NO:5; or SEQ ID NO:7.

THIS PAGE BLANK (USPTO)

45. The method of claim 39 wherein said nucleic acid sequence is introduced into said animal in the form of a transgene construct.

46. The method of claim 45 wherein said transgene construct is a nucleic acid molecule.

47. The method of claim 46 wherein said plasmid comprises a nucleic acid sequence according to SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:6.

48. The method of claim 39 wherein said animal is chosen from the group comprising pigs, goats, sheep, cows, horses, rabbits, rodents, cats, dogs, fish and poultry.

49. The method of claim 48 wherein said animal comprises a mouse or a pig.

50. A nucleic acid molecule comprising a nucleic acid sequence including a gene encoding a protein, said gene being operably linked to at least one regulatory sequence for gastrointestinal tract specific expression of said protein.

51. The molecule of claim 50 wherein said at least one regulatory sequence comprises a salivary protein promoter/enhancer sequence, whereby expression of said protein is salivary gland specific.

52. The molecule of claim 51 wherein said salivary protein promoter/enhancer sequence comprises a parotid secretory protein (PSP) promoter/enhancer, a proline-rich protein (PRP) promoter/enhancer, a salivary amylase promoter/enhancer, or a SV40 promoter/enhancer.

53. The molecule of claim 51 wherein said protein comprises a phytase or a homologue thereof.

54. The molecule of claim 53 wherein said molecule is a transgene construct.

55. The molecule of claim 54 wherein said molecule is a nucleic acid molecule.

THIS PAGE BLANK (USPTO)

56. The molecule of claim 55 wherein said molecule comprises a nucleic acid sequence according to SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, or SEQ ID NO:6.

57. The molecule of claim 53 wherein said molecule includes a nucleic acid sequence according to SEQ ID NO:3, SEQ ID NO:5; or SEQ ID NO:7.

58. An antibody specific to a protein expressed by a nucleic acid sequence according to SEQ ID NO:3, SEQ ID NO:5; or SEQ ID NO:7.

59. The antibody of claim 58 wherein said antibody is monoclonal.

60. The antibody of claim 58 wherein said antibody is polyclonal.

61. A hybridoma secreting the antibody of claim 59.

62. A host cell transfected with molecule of claim 50.

63. A host cell transfected with the molecule of claim 56.

64. A host cell transfected with the molecule of claim 57.

65. The host cell of claim 63 wherein said cell is an bacterial cell.

66. The host cell of claim 64 wherein said cell is an animal cell.

67. A diagnostic kit for immunologically detecting a protein expressed by a nucleic acid sequence according to SEQ ID NO:3, SEQ ID NO:5; or SEQ ID NO:7, the kit including an antibody specific to said protein.

68. The kit of claim 67 wherein said antibody is monoclonal.

69. The kit of claim 68 wherein said antibody is polyclonal.

THIS PAGE BLANK (USPTO)

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference 00264-0003	FOR FURTHER ACTION		see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.
International application No. PCT/CA 00/ 00430	International filing date (day/month/year) 20/04/2000	(Earliest) Priority Date (day/month/year) 23/04/1999	
Applicant UNIVERSITY OF GUELPH			

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 4 sheets.



It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the language, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.



the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of the sequence listing :



contained in the international application in written form.



filed together with the international application in computer readable form.



furnished subsequently to this Authority in written form.



furnished subsequently to this Authority in computer readable form.



the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.



the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ Certain claims were found unsearchable (See Box I).

3. ☐ Unity of invention is lacking (see Box II).

4. With regard to the title,



the text is approved as submitted by the applicant.



the text has been established by this Authority to read as follows:

5. With regard to the abstract,



the text is approved as submitted by the applicant.



the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the drawings to be published with the abstract is Figure No.



as suggested by the applicant.



because the applicant failed to suggest a figure.



because this figure better characterizes the invention.



None of the figures.

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 00/00430

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A01K67/027 C12N9/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

STRAND, EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 48812 A (UNITED KINGDOM GOVERNMENT ;YANKE LINDSEY JAY (CA); CHENG KUO JOAN) 24 December 1997 (1997-12-24) page 18, line 27 - line 31 ---	1-69
X	WO 99 17610 A (UNIV CALIFORNIA) 15 April 1999 (1999-04-15) page 7, line 12 - line 26 page 10, line 21 -page 11, line 22 ---	1-5,10, 50-52,62 6-9
Y	---	
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

29 September 2000

Date of mailing of the international search report

13/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sprinks, M

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No

T/CA 00/00430

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SELINGER, L. B. (1) ET AL: "The rumen: A unique source of enzymes for enhancing livestock production." ANAEROBE, (1996) VOL. 2, NO. 5, PP. 263-284., XP000940728	21,22, 26,28, 29,50
Y	abstract page 267, column 2 page 276, column 1 -page 277, column 1 ---	15,30, 39,40, 45,46, 48,49
X	ZHANG J X ET AL: "Expression of a bacterial endo (1-4)-beta-glucanase gene in mammalian cells and post translational modification of the gene product" BIOCHIMICA ET BIOPHYSICA ACTA-MOLECULAR CELL RESEARCH, (27 JUN 1997) VOL. 1357, NO. 2, PP. 215-224. PUBLISHER: ELSEVIER SCIENCE BV, PO BOX 211, 1000 AE AMSTERDAM, NETHERLANDS. ISSN: 0167-4889., XP000940605 abstract ---	50,62
Y	WO 99 08539 A (DIVERSA CORP) 25 February 1999 (1999-02-25) Discloses a phytase - relevant for present SEQ ID NO:7 figure 1 ---	11-20, 30,31, 36,37, 39-49, 53-61, 63-69
Y	DATABASE EMBL 'Online! ID: AF062078, 18 May 1998 (1998-05-18) DENNIS ET AL.: "Cloning vector p34S-Cm" XP002148835 Relevant for SEQ ID NO:4 abstract ---	56,63,65
Y	DATABASE EMBL 'Online! ID: MMPSPG, 3 November 1992 (1992-11-03) MIKKELSEN: "M. musculus Psp gene for parotid secretory protein" XP002148836 Relevant for SEQ ID NO:1,7 abstract --- -/--	1-7, 10-14, 16-69

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 00/00430

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>DATABASE EMBL 'Online! ID: BLCAT3DNA, 26 February 1992 (1992-02-26) LUCKOW: "Plasmid pBLCAT3 gene for beta-lactamase and CAT gene for chloramphenicol acetyltransferase" XP002148837 Relevant for SEQ ID NO:2,6 abstract</p> <p>---</p>	47, 56, 63, 65
Y	<p>DATABASE EMBL 'Online! ID: MMU73190, 4 November 1996 (1996-11-04) LAURSEN ET AL.: "Mus musculus positive acting regulatory region of the parotid secretory protein (PSP) gene" XP002148838 Relevant for SEQ ID NO:7 abstract</p> <p>---</p>	1-7, 10-14, 16-69
Y	<p>DATABASE EMBL 'Online! ID: CVCATBLA, 24 January 1992 (1992-01-24) BOSHART ET AL.: "Cloning vector pBLCAT6 encoding chloramphenicol acetyltransferase (CAT) and beta-lactamase (bla) genes" XP002148839 Relevant for SEQ ID NO:3,5 abstract</p> <p>---</p>	14, 20, 31, 36, 44, 47, 57-61, 64-69
Y	<p>DATABASE EMBL 'Online! ID: RNRP15, 12 May 1991 (1991-05-12) LIN ET AL.: "Rat salivary proline-rich protein (RP15) gene" XP002148840 Relevant for SEQ ID NO:3,5 abstract</p> <p>---</p>	1-5, 8-14, 16-69
Y	<p>DATABASE EMBL 'Online! ID: ECAPPAA, 5 February 1991 (1991-02-05) DASSA ET AL.: "E. coli periplasmic phosphoanhydride phosphohydrolase (appA) gene" XP002148841 Encodes a phytase - relevant for present SEQ ID NO:3,5,7 abstract</p> <p>-----</p>	11-20, 30, 31, 36, 37, 39-49, 53-61, 63-69

THIS PAGE BLANK (USPTO)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 00/00430

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9748812	A	24-12-1997	US 5939303 A	17-08-1999
			US 5985605 A	16-11-1999
			AU 716845 B	09-03-2000
			AU 3021697 A	07-01-1998
			BR 9709793 A	11-01-2000
			CA 2257101 A	24-12-1997
			EP 0904385 A	31-03-1999
			NO 985804 A	10-02-1999
			PL 330532 A	24-05-1999
WO 9917610	A	15-04-1999	US 6004944 A	21-12-1999
			AU 9668998 A	27-04-1999
			EP 1030557 A	30-08-2000
WO 9908539	A	25-02-1999	US 5876997 A	02-03-1999
			AU 8904498 A	08-03-1999
			EP 1003379 A	31-05-2000
			US 6110719 A	29-08-2000

THIS PAGE BLANK (USPTO)

SEQUENCE LISTING

<110> University of Guelph
 Forsberg, Cecil W.
 Golovan, Sergeui
 Phillips, John P.

<120> Transgenic Animals Expressing Salivary Proteins

<130> U Guelph - 03

<140>

<141>

<150> 60/130,508

<151> 1999-04-23

<160> 7

<170> PatentIn Ver. 2.1

<210> 1

<211> 20623

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Lama2/APPA
 plasmid


<400> 1

```

tcgagagtat ctttgtcagc tgtgcctcca acaaaggggt actgttgcc acatagaaag 60
atctaaacta attaattaat ccctcaccgc caaatctttc agtcactaag ttagcacgat 120
tggtgaacaa gttctccaaa ggagagatac agatgagtgc gtataggggtg gacctggctg 180
ctgaggagac acctgcatct gactaagaag agccacgggtg ttagttgaat ggtgtggagt 240
aggggtgggtc tgtgggacag tagaaaatcg agaggcatgt gccgtttagt gaactgatgg 300
aagctacccc aaacgacaga gattgtcagt caggccaatc cgtttcgagt ttgatgggca 360
gccggacagt gagacagaca cacctactca gttggaggaa ggatgagaac aatggccagc 420
agggattgag agaccctgac aggcgcaagg ccctaacaca cacacctacc acctcacttg 480
acaaagctgc caaagaccaa agacttggtc tccattagaa atgacagctg gcttgacccg 540
acagcataat aagcagagtg tactctgatt ggagaacttt aatgtgtttc attcagtatt 600
ataaaaggac agtattacag attttgttgt acactgctgt tacatgtggg gcagtgtgtc 660
tttaagtagg gtaaagtact ctttaaaaat gggtcctaga tattttttcc tttaactcaa 720
gtctcttact gtttaaatga tttttatatt gtttaatatg gaggaaaaag aagcgtaa 780
ggacaatata tatttagaga aagatgggta gctgtcagaa aaatatgcaa atcaaaatca 840
caccaagact gcagcacacc cctgtcagat ggctgtgatc aagaaaataa atgacaatga 900
gtgggtgggtg agatgtacta aagggaacaa cacacacaca cacacacaca cacacacaca 960
cacactggag caaccactgt ggaaatcagt atgaatgggtc ctcaaaaacc tgaagataga 1020

```


ggggggctg gtggcataca cttttattcc cagcactggg gaggcagagg cagggtggatc 1080
 tctgagttcc aggccagcct ggtctatagc acagggttcta ggacagccag ggctacacag 1140
 aaaaaccctg ccttgattaa accaaaccaa accaaaccaa accaaaccaa accaaaccaa 1200
 accaaaccaa accaaaccag accaaaccaa aacactgaag atagaacttc agtattccat 1260
 tcctagatat atacccaatg gagactaagt cagcaagaca cctgcacagc catgttccact 1320
 actacactgt tcaccacagc caggctgtgg aaccagcctg agtgtccatg ataaatgaat 1380
 ggataggtaa ctttcaagggt aaatggactc tgctgtgtac atgcctcaca ttctgtttat 1440
 tcatttttct ttatgagggtg tccattcagg agtcacatgg tagttctatt ttcagtcttc 1500
 tgaagatact acactgggtc ccacagttta cactttttatc agcagtgaat aaggggttcct 1560
 ctatccttac catcatttgt tgtaattttt cttgatgacc ctctttctga cagggatagg 1620
 atgtaatatc agtgtgagga agtacaactt gttttctaag tatttattgg ccccttgcat 1680
 ttcttctttt gaaaactgtc ggttcctgac atctgctcag gtattcattg gatgttggtt 1740
 ctttggtggt tgagttctta tgaattctag atgttaaate cctgcctgtg gttctctccc 1800
 attctgtagg ctgcctcctc accctggcaa ttgttgctct tgttttgtag aaacttttga 1860
 cttcatggaa tctcatttgt cagttttccc tctctgcta tagcctgagc taatgcaactg 1920
 gtttttacag agccctgggtc tatgccttta tctcctctg gcagcttcgg agtttcattt 1980
 cttacattta gatctttgat ccactttgaa caagttttgg agcaggggtga gagatacgaa 2040
 tctagttcca ttcttccata tgtgatccta gtttacatag catcgttggg tgaagaggtt 2100
 ttattttatt tttaaataat gtgtcataaa aaacgaggtg gttgtagcag tgtggatttg 2160
 tttctttgtc ctttgatcta caggctctgt tttgtgtcag tctcatgatg ttttattgct 2220
 atggctctgt catacagtct gaggtcagggt attgtgatat accttcagta ttgctccctc 2280
 agactcagggt ttgctttggc caggagtcac cttactcagt gctcttagag ctccccccagc 2340
 atgtagctgc tactattctt agttgataaa tcaggaaact ggggctcaga gagattaact 2400
 gtcttgaact acttctgggg aggtgaaacg tggagacact aaactgtgtt taccctgtac 2460
 tgctccagta gctgtcgggt gctgggctac agcaaagcac ctatactata tattactcag 2520
 gaggtggaaa aactcagcct cccttgggggt tcccaagctc ccagggtgtcc agtcactgct 2580
 ggaaacctca tggagtctga aaggaagggt tgagggtaca tggggcagcg atgaggagcc 2640
 tggggctggg atctcccaa cacctggata tccagatgcc actgggtcag ggggagttgg 2700
 gaacagagtt gggatgtcca tggacctgtg acaaggccag ggccaggggg aggataactc 2760
 tggctttact aatttgcgaa agtccttagc ttagcagcag ttgtctggga gcacagaggg 2820
 gccttctgta agaggctcag gcagtgccgc tctgtaggcg aagggtcttct ccatgttccc 2880
 catggtggtt cttgatgaaa gagacagtcc ttggctccaa actgggtttat tgattgttca 2940
 ttgtggaaaa tgggtgcaca ccaccttctc aggggtggacc agagatcaaa taccttttgc 3000
 agggaggaat atctgggaag ggacgcttac tggctaaacc ctccaggcct ctagatacat 3060
 cattagcatg gagaactctg ttctgggcta catgaccaca ggccacattt ccacaagcca 3120
 catgtgggaa gtgtggcaca tgttctaggc caggaatctg gtagggagcg tggagccacc 3180
 taccatccca ggtgggtgcc tgggtgccag ggacctgaa cccgctcaac cttaccaagt 3240
 ttctggcag ggtccactgt cctacacaga agctggagga ggtgtgaggg ttgtgtcttt 3300
 gtggaatgtc ccatgctgct tggggctcag tttctccacc tgtacctcat tggtttgggt 3360
 ataaaaagtg gggatacttt attattctct gactcggtcc tgaggaaaaa gcatcgtggc 3420
 agtccaggaa ccacacctg aggttcctgc actgaaggga ctccctaagt ctctggagtc 3480
 tctccccttc acagagctgc caaagtctag gttcttttga ggataacaga gccatgcttg 3540
 gtaagcagac aacagcattt gtttactcaa cttctttttg tcagctccct cttcataaac 3600
 aagttgagac accatgctgg cttgaggaag acttctaaag ccagacaact gtgcaaggaa 3660
 gaagaagaag gggcaagtgg agttagcctg gatgtagccc tcaaagtctc cagagaccag 3720
 ccatgaaggc tcaagtggag ggcaagacct gcagcagcca agcatctggc aggagaggat 3780
 cctgggaacc cctctaccat gacacacatt cttcctgcag gtcacactta ataggccatt 3840
 tcttatttgg atctatcatg gtgttctgtg cgagattaat gaggtgttat gctgcgaaca 3900



```

gaaagttata taaaaacaag tccccccccc ttgtcactgc tgctaagaat gtagcagaaa 3960
ttgtctcaag tgtctctcta atcagaaaca ataaagggtct ccttggattc aagccctcca 4020
gtttcctcct tccttgctga gccttggaca cccatacaaa cctcctggat gctacagctc 4080
tgggcagaga ctccaagggtg gggagagact gatggtacaa aagcaaaata cttgtttggg 4140
ggtacaccca ctctctgcc tgtgtggttc ctgcagtcag tcctgcagac aggccctcag 4200
tgggtcttcc atgggcaaca cgcagagga ggcaatggat gggaataccc acaccctggg 4260
tagtttacct cggccatgct ctctgctctt catccctcct ctgccctctg ccacggcttt 4320
ctctgcagga atcatatctt catattggcc cacagggtgtt ctctcacc tagctatgat 4380
gtttacttta gagtgacctt agcagggtct gtgggaatga gttctagaag gctcacggag 4440
atgctaggga agaaacgtct tctaactact gaggttacta agttcctggg gggtgtctct 4500
gcctttccct tgttaaagtc acctgaagt tagtgcagaa gaaatcagag ccagtcaca 4560
gagtaaatat ggtcctgaag atttcctttg agtgcccaga atccatgaca tttcaagagc 4620
cctctttgta ccttaagtca tttggggttg tatcttctgc ttgatgtatg tgtgtgtgtt 4680
tatcaaagag tgagatggtt acataagagg tgctctaaag gacagagagg atttgcaatt 4740
gtggcatgtg acatcctcag gccttgctct ggtgccagga ggaactgatg cagaaaagag 4800
taagagggtca tttcctggag gctgtcacta tagaggagat cttacagtgc attccctcct 4860
ccaggccctg cctgaggata gacatgtgct gactgcaact gaaacagagg cttgggatgg 4920
agagttaggt tcacagaagg gaggggtggga gatggatgct tgctgggttc tgggtctcat 4980
caccagctcc tgaccacccg gtcagcccat gtgcttattc catagctttc ttttgctatg 5040
tttactcagt gtggtgtttg ttgggaccca gcagaagcca gtcccaggct gacagctgtg 5100
gatacacagg gcagcatgag ggtcctcagc ctgaagcagt caggctggca gaagagaaaag 5160
accagcacac attccttcaa ccaactatgt cttgaaaaac aaacatatta tatcacatat 5220
attgcattta tgagacagct aaaatgtact cgggtagcat gactccaggg ggggatattc 5280
gcaagtgcc tgaagtggcag agggacagcc aatgtgaggc aagaaggaaat tctggctcaa 5340
cacagcttag ctccctgggtg ttggttcaaa ctttgagagt ttgaccacaa gcactttatt 5400
tttgacatat ttaaacagag cacaactttg ggaaaaagtt ttcttatgaa aattatcaca 5460
ataaagctta aggcattgact acattaaaat gcctttgcaa agtatatgtg ccctcttcca 5520
caagaatggt tctattgact gagaaataat gttcaggata aagatccagg aagaaaagat 5580
cagggtataag taaaatacta aactcttttg caaagtacat agaccctctt tcataacaat 5640
gggttctatt gactgacaag cactgctcag gagttgggaa agagtctagc ataagcacga 5700
tagcctggag actctagtga ggtctagtct tacagacagc aaaaatcacc aggttacaaa 5760
ctacattcat ttccagtttt ctgatcaggc acaggatatga atccctctctg ttgaagagaa 5820
aagtccatgt gtttaaaata tctggtttct ccagtgttat tagcgagaag acttgagccc 5880
tatacaactc ccacctggag tgacatcctg tcttcattggt atattacata cctagacacg 5940
ctcatctcac agacttagga ctttgtcttc tgatctccat ttctgatccc acttccacct 6000
ttgccttgat agtgtcattt tcttactgct cttggtgaca accatgttat cctctgtgta 6060
tttgagtgtt accattttca gattttacct gtatgcaaga tcacacagtc tttgtctttc 6120
tgtctggatg catgctaata tctacacaac aacccttccc cgtcactcag atcttctctc 6180
attaacacat acatggtgct gaagaggcta gggagcttcc cttcagtggg gagctagctg 6240
gctattgggc ctttttgact gtccaggaag gcccgaatt gctgagacaa gaacttagat 6300
tcttcattat tgactctaac tcatgtatca agcagaagct aatgaatagt tatcaacagg 6360
atcagagggt ccagtgtaa gacatttgac atgaaagaac ggaggaagga cagatggatg 6420
cataaaagca ggaccactgc cccaggaagg tcctggaaac tgatgcaggg caaaggacag 6480
gttataaacc aaatcttagg gagtcaggaa gagcacagag gagctcaacc aactgaccac 6540
tgcttagggg ctaccaaccc aatcctcct gtgggaacag ctaagctatc agccaagggt 6600
aataaacagg caggacctgt ggatgacatg gagagcatag ggaccctggg tccagccttt 6660
agcacctgca ctctcaggat actccacct tgtgtcttag agagcctagg gatactgggt 6720
ccagcctttg gtaccttcac tctcagggt ccccatcact gtgtcttgga gagcctaggc 6780


```


accctgggtc cagccttcag tacctgcgct ctcaggacac cccaccattg tctcttgccc 6840
 cgtctcttct tectcttctt ccttttcatt gtctcttctc tgtttctttc ttgactctcc 6900
 tttccctca caccctcact ctagttctcc ccttccctct ctgcatcacc ctattctctc 6960
 tgtggtccct ccactttctt ttatctctca tgcctctctc ctcctcaca tacttgctac 7020
 ccactatact tcaggggcca gctctagtga caaagctgtt aatagcaaga ctctcagatc 7080
 tccaacggct cagaggagcc agaccacca agaactctct ccagggtcaa tttcagggtc 7140
 cttcgaaagc tttcagcaaa tgctcaggga acatgccact aacaagaaga tgcaaattcc 7200
 agttgagagt gggaaaggcc cttgcgtagg tcccatcttc caggccaagg tcagaggggc 7260
 tctgtgtaat cgggattgac agggctcaga acaatgtttt gtttttaagg tttatttatt 7320
 ttaggtgtta gtgtctttgc ttgcatgacc ttatgtgcat catgtgtgtg cagggtcctg 7380
 atgacagtag aggagggtt tgaatccctg gggataggaa gttacaggaa attataagct 7440
 gctttgtggg tcttctagct ttcccaacag aagtgaatgc tcttcaccac tgagccatct 7500
 ctctaggccc aagagacatt gctttatgga tataattgtg tgtgtgtgtc aacattgagg 7560
 aaagggaat aaaaaaaaaa cttcagccgc taagggtgtg cagtttctact aattgctact 7620
 ttagttgtg ataaaatggc aggtgcttca acatttatat atacaaaaac ttccctgctg 7680
 gtggttcaac tgtgagaact ggggtaagt ggtgagttct ctttttctgt ctctgtctct 7740
 gtctctctcc ttccattctt tcttaaagga aataaacatt gcagctgggt tatagctcat 7800
 caatatggaa gttacagaag tgaaaaaagg cattgccttg gtgggtgggt ttaccagctg 7860
 atttttggtt gtcctgcaag gaggtctggg gactggctgc tctgtctctg tctgtatgag 7920
 tgagggaagt ctggggagca gattccctaa ccttcagcct ggctggttc ctgagtgaac 7980
 ccagcctctc tggctctagt agctttttcc aaacaggaat ctgagtgggt acaggggaaca 8040
 agtaccagcc cattgcttaa gtgccagggt tagtgagggc aggaagctgc catagctggg 8100
 attagtagtt gtattggatg taggaagtc tctctggga cagctaattc ttaatgcttc 8160
 actggagatt ttcaatgaga aatttatccc acggcccata tggcccatc cttttgtctc 8220
 caacagccaa gtattttcca ttagaggaga cttcctgtac acttgatgga tgctcattcc 8280
 aagggtgactt ggggcagtca gtacagactt gggatgacct ctgacagcct aacctctccc 8340
 caacaagggc cctctatgtt tgctatgtaa tgtaatgtca gacattgtca ggagtgtccg 8400
 cagcacagcc tgcccagtggt gagggctctc atagggttcc cactgtctta tctacacagg 8460
 gataacgagg aggtaagctg cagttcccag tctcacttca cagaggaaga gataaccca 8520
 tcccagggtca tgtagccagc agtggaaga atgaggattt gaactcaggt cttccaagtc 8580
 ccattgatag catctctca caagtccctt gccacctca cgatgcctta gacacttgc 8640
 tgccctttat actaaggaga tgcaggtaaca aggggtttac ccatgtagca gctgaggcag 8700
 ctggggatag ataccagcag caggcctgat gtcaccactc taactccagc atcccagtc 8760
 tgtgttctct gagtgtgaaa atccctactt aacaagattg tgcaacagtc cttggctctg 8820
 tgacccatag ctggaaacag gattctcatt gatttgtgga acatgggtggc agccagccaa 8880
 aaagagggtc tgcatacaga agacacgtgt ggcaaggcca cagcagactc tgactacctt 8940
 agcttacaga attacaaggt cataatgtcc tctgctttgg tcacctcatg ttaaggacag 9000
 gccctaataga agatggggca gaagactgaa ggaatggcca accaataact ggcccaactt 9060
 gagacccatc ctacaggcaa gcatcaattc ctgacactac taatgatact ctgttatgct 9120
 tgcagacaga agcctagcat aactatctc cgagaggtcc acccagcaac tgactgaaac 9180
 agaaaaagat atccacaggc aaacagtggg tggaggtcag ggactattat gggagagctg 9240
 tgggaaggat taaaaaccct gaaggggata ggaacccac aggaagacca acagagtcaa 9300
 ctaagagacc tgtgggagct ctcagagact gagccaccaa ccaaagagca tacacaggcc 9360
 ggtccgaggc acctggcacg tgtgaagcag acatgcagct cagtctccat gtaggtcctc 9420
 caataagcgg tagcctgact gcagtatcca atccccaaca gggctgcata gtctggcctc 9480
 agtgggggag gatgccccta atcctgcaga gacttgatga gtggagagct atccaggggg 9540
 aaccacccct ctctgagaag ggaatgggga tgggggaggg actctgtgaa gaggggacaa 9600
 ggacaaacaa gaacctcaaa taggtcaggc cctaaaggct tgctaagtag cagtggccca 9660


```

gctctgtcct gttcctcagc ccaaggetca gctcccacct gtttctgtgt ttttctggct 9720
tttcatgggc ctaggacttg gtgaccagtt caaacaatgg ggccctgtgga agacacaata 9780
tacaagacta gggacattcc tgttctgctg actatccata gcctgatgta ggtggaagga 9840
cccaatcact ggatttctac ccttgacaaa ccttgacagc tgagggcctc tcagaaacct 9900
atcttctcca ctgaaaaatg agactctcaa atgaacgtcg tgacaatcat caggcttatt 9960
aaagaggtgt atctaacctg aatggcaagc agacagcagg caaatgtctg tatcaacctc 10020
taggaaggac aagaactgct cactgctgcc ccccaggagg ccatttgctg aaacagctgc 10080
tctcctgctg gtgcacaggc cctgccttct cattgcagcc acagcccctt cctgtctgaa 10140
cctcctgtca ggtcactggg aaacagatca agatggaaca ggacagctcc tgatggtaaa 10200
taaaaaacag tggcatggc tattcatagg ggtttatgct tcttcagtc acactgtgaa 10260
gagctgtggg catgaaccac agtgctcgag gtagagttgg ggttctgaaa ttcacagtgg 10320
ggtgagctca gtaaagtga gctggaggtc actcgtgaga cacacagtcc tgctgcttct 10380
gttcccaata tcctgaggag acgacacatc tactttgttc agaggccaca gtctagtga 10440
cctgagagtt accagtttct tatttgtgtg tgtgtgtgtg tgtgtgtgtg tgtgtgtgtg 10500
tgttgttcgt gtgtgagtgc aggtgcacat atgatagcgt acacgttgag gtcagaggat 10560
aactatcagg cgttgtcccc tcctactttt cctcggactc tggagaacaa acatgggtcc 10620
ttattccagg ggagcaagtc gctgttggtc gacacatctt gctcacatac attttaccta 10680
gacaatggag cctccatcag agtattactt tagctcctca ccgatggcaa tgcaccacct 10740
ctctaccac ataggagttg ggtctccaca cccccacac ccccttcac caaaacgttt 10800
tcagttactt tatctggtaa agttcatcag agaatgaagc cagtattaag aacatggaat 10860
catttgaggaa cctggatcta gcaatacccc accctagatg gagttgctga gttttcacct 10920
cagattataa tcccccccta gcttctatgg tttattctga aaccagggga actcgattcc 10980
tccctttgga ccacagacat cctggcttgt gaattcacat gtcactctact gctaaccat 11040
tggtagtatg tggctcacag agacacacta cagtcatggc caatgtcaag gtaggacaga 11100
tgtgaatcat tccccagtc ctgctgtttt catgactaac cctcctcagc acagtgacca 11160
tgaacctact tttccctcc ttttattttt agaattgctg gaattttcta ttttgagaaa 11220
taatagcctt gggcagcatt aaacaaaatc atctagaaag ctggtttaaa atacagatgg 11280
ttgagtcagt gaaagagtga ggaatgtcat tattggcccc tcacagaggc tggctcactc 11340
cagcagaggt ggttgaagct cttggacacg ggtcaggtgc ataggaaagg tngtctggga 11400
cactgagaac cacaattgaa caaacagAAC tggttgcttt ttttttttta aatgagttct 11460
caaaaaatga ctggctagct taggcaaata cttcgagcca acccaacaga acattcttcc 11520
attgattcat tctggatctt ctttctagac aatactgaac tgacccttg ttggcagctc 11580
caagtttgac aacatagggc tttgaacttg gcacaaggtc catcactgtc acccaagcat 11640
cctgggtgac ctttgggttg gaatatcttg gctaacctta gatattttct ttggagtatc 11700
tttagaacat ccaggaaata gggcttgatt ctcatcctgg gaccacaata taagtcaccc 11760
tagaatccca ggagatcgtg cagagaaaca aggatctctc tcgtgtgcat ccttcttcaa 11820
agcagtgagt agtgactcca ctaaaactgag ttcccatctg agagtccaca ggaggctttg 11880
gggcaagaag cagaggggaag gcaactgttg tgttggtaaa gttttgactc taacaaattt 11940
gaagacatag atgacattgt gtcagactaa caacaaccta gactcatgtg ggttctgttt 12000
agggatcaga ttttattcat caatgacttg tcttagtgta tagagaaagg cttcctactg 12060
gagtgtaggc tcaataatga cagaagagat agctatttcc cctagggact gtgctgctcc 12120
aagtttggtg gagaaaggca gtggggaacc tagatgtgct ctctggggag ggggtctgaa 12180
gctggcttca tagaaggtgt gaagttttgc tgaaacatct aaacagaatt atagcttagg 12240
aaagtgagca ggcaaggcag ggaatgtgtt gcatatgtat atgtacatga atatatatg 12300
ttatagatac acacacattt gaacctcatt tgcagatgac agaaaatagg ttattttgcc 12360
tctcttaact gctaagcaca atgacttcca gttccatcca tttcctgaaa tgccacaatt 12420
tcatttttca ttgtggctga ataaaattcc attgcagact gggccctact tcatccactc 12480
ctgagggcag gcatatcccc tggctccatt tcttacctat tgtgaagaga agtgcaactg 12540

```



```

tcttgttgaa aggcaagcgt gagagaggca ggcactaatt gtggggtttt gtttcttctt 12600
cctgctatga ctctccattt gtcagaacca aagatcgata aaagccgcca ccatgaaagc 12660
catcttaatc ccatttttat ctcttctgat tccgttaacc ccgcaatctg cattcgctca 12720
gagtgaagcg gagctgaagc tggaaagtgt ggtgattgtc agtcgtcatg gtgtgctgctc 12780
tccaaccaag gccacgcaac tgatgcagga tgtcacccca gacgcatggc caacctggcc 12840
ggtaaaactg ggttggtgta caccgcgcgg tggtagacta atcgccatc tcggacatta 12900
ccaacgccag cgtctggtag ccgacggatt gctggcgaaa aagggtgctc cgcagtctgg 12960
tcaggctcgc attattgctg atgtcgacga gcgtaccctg aaaacaggcg aagccttcgc 13020
cgccgggctg gcacctgact gtgcaataac cgtacatacc caggcagata cgtccagtcc 13080
cgatccgtta ttaaatctc taaaaactgg cgtttgcaa ctggataacg cgaacgtgac 13140
tgacgcgatc ctacgacggg caggagggtc aattgctgac ttaccgggc atcggcaaac 13200
ggcgtttcgc gaactggaac ggggtgcttaa ttttcgcaa tcaaacttgt gccttaaagc 13260
tgagaaacag gacgaaagct gttcattaac gcaggcatta ccatcggaac tcaaggtag 13320
cgccgacaat gtctcattaa ccggtgcggt aagcctcgca tcaatgctga cggagatatt 13380
tctcctgcaa caagcacagg gaatgccgga gccgggggtg ggaaggatca ccgattcaca 13440
ccagtggaac accttgctaa gtttgcataa cgcgcaattt tatttgctac aacgcacgcc 13500
agagggttgc cgcagccgcg ccaccccggt attagatttg atcaagacag cgttgacgcc 13560
ccatccaccg caaaaacagg cgtatggtgt gacattacc acttcagtgc tgtttatcgc 13620
cggacacgat actaatctgg caaatctcgg cggcgactg gagctcaact ggacgcttcc 13680
cggtcagccg gataacacgc cgccagggtg tgaactggtg tttgaacgct ggcgtcggct 13740
aagcgataac agccagtgga ttcaggtttc gctggtcttc cagactttac agcagatgcg 13800
tgataaaacg ccgctgtcat taaatacgcc gcccgagag gtgaaactga ccctggcagg 13860
atgtgaagag cgaaatgcgc agggcatgtg ttcgttggca ggttttacgc aaatcgtgaa 13920
tgaagcacgc ataccgcgtt gcagtttgta aggtaccggg ggatcacaaac ttgccctctg 13980
aagaggaaga acagaaggat gccacaactc tctgctggc tactctccag tggtttcac 14040
ttacttctga tggcatttcc ctctagaaag tgctactatc atccacacat ttctacctga 14100
gaccaccaa aggacctcc caaattctct tctctctga gtagtctcca cacctgttac 14160
caccatccca gaattaaaat cctaactgca ctctggcggt tgacttgccct cagtccctgc 14220
aataagagtt gttggcagtg ccaggcgtgg tggcgcacgc ctttaattcc agcacttggg 14280
aggcagaggc aggcggattt ctgagttcga ggccagcctg gtctacagag tgagttccag 14340
gacagccagg gctatacaga gaaacctgtg gtcgaaaaac caaaaaaaa aaaaaagtt 14400
gttggcagag tgtgggttat ataccagggt gagatttcaa atgagtggtc gaagctgtag 14460
ccagaaggaa cttagaggat agctcataac ttaaaaagaa atgtagagag tagcagaaac 14520
attgagagag tgggcacaca gccactgtgt gaatgtggca gaacacaatc cagccagcta 14580
tacatgcata agtgtatatt ggcgccatcc tgactgatga gacacaggaa aacagataga 14640
cggggttagg tggccatggc ctttctctgc tgcctcttcc taagggtcat ctcaagacct 14700
tatgctctct taactcttcc attgctactt agcttctaga tatcacctcc agattagtct 14760
ccttgggtac atcagtgatc ctggtgatat ccagggtctc ctgattccat ctttgtcata 14820
gaggctgcaa ctaaagaggt cttcttaata cttcacacc tgatgcaaaa aggaagacac 14880
agaagtacac agaggtgaag tgattcatgt aggacataca gtgagcaagc atcagggtcc 14940
ggattatctg actctactct aacttttatg taaatgtgct ttatgccatt aacactgtca 15000
ttcctgtgct tcagctctgg gagactccca agcactctta ggcacaagcc acaattaagg 15060
gactctgaca ctctgcattg attaattagc atgggtggtc ctatgtttcc agattcatga 15120
ttgtttcact ttccatatag gctatgaagg gtgtgaggaa attttttggg gacagaattg 15180
gaggcaatcc acctctctca ggaagcctct atctggaaaa gcttacaact cagggacagt 15240
aactgtaggc ccagtccttg gtgtccaaaa tgggttttat ggtttgaatc tgcaaagcct 15300
tccatgtgct caaaggtttg aacatggagc ctctctctgg taacactgta ttggaggctt 15360
ttgagactgg atgctctttg gtcccatggt ttgctacatc atctgtcaag atatgacca 15420

```



```

ggcatgctac cagctaccac agactatgcc tctccagctt tcatgtttctc cccaccatga 15480
tagacttgta tctcctaaaa atggaatcaa agcaaacttt tcttgcatga agttttttttt 15540
tttctgttaa gtgttttggtc acagggacaa gaaaacactc aatacagata attagtacca 15600
gagttgaggt tcattgctct agcaagttgg atcaaatttt tagggctttg gaactgattt 15660
ataagagaca tgtagaagag tctgaagctg tgggctacag aagtgtcacc agttttttaag 15720
aatagtttaa tacaccatgg gaattgtgaa aatcagaatg ctcacacaaa ggcagacagg 15780
aaaacgtgag catgtggcgt gtgagagggc ataagaagga acctaggggg aaatgagcta 15840
gaagccattc ggctacgtta gggaacgtgt gtggctgtgc ttggcccatg ccctggcaat 15900
ctgaatgagg ccaaatttta aaggagtggg ctaactcgat tgtcagagaa aatatcaaga 15960
cagaccacca ctcaggctat gccgtgtttg tgaccgacca gctactctta gccagctcta 16020
ttgtgaaatt ccagagcaat tatcagagca tgaagataca tacagtttag tgaagtaagg 16080
ggtgtgggtc cctaagtggg tgggtgcataa atctatgtag gtgatgccta agtgacactt 16140
gataatccaa aatatcagca atgtggaatg tcttccaagg agacctgtag acacacattt 16200
tagaactttg ctcattggctg taataaatag ctagctagaa atcatttcct gaagaggtta 16260
gtctgagtta cggttccagg gcaaacattc agtgatggca aggaaggcat tgcagtcagg 16320
agccaaaggt cagctgggtc cattgcatca agagtagaga gtcagagtgt gagtagaaag 16380
aggatacagg ttataaaacc tcaactgtcca ctctcagcaa tccattttct cctaaaaggc 16440
tttaccttct aaagatttta gtcttcaaaa ccagtaccag tagcctggga acaaaagttg 16500
aaacaaatga gcctttgtgg ggcatttcac acttaaaaca gggcatcacc taggaggagc 16560
cctgtgtgca gtaggaagtg tggcctctgt gtcaggaatg ctcaggctaa taaggggtcc 16620
tctatctgag ggaccctatg aagattcaac aagtagttgt gagaattccc tgtaaattgga 16680
tgctaccaat ttgacatttg tagacctgct attgtgtgct tctttatttg gctctcccat 16740
ctcccaactt tccaacccat attccacatt aatcccttcc accaccatgc aacactaggt 16800
aggagagaag gaaggttaga agagaaagtg ggtatagatc tatttagact acttctgct 16860
gattaggggc aagtccaatc gtcattgtca ggatacctcc aaccagcaac cagcaaacca 16920
gcaaatcaga aacagcaaaa gcagccaaca aggagcact aaccagcagg attggggctg 16980
gtagcgtggg agcagtcact actggtcttc tcatggcttt ggcattaata ctctctcaag 17040
aaattccgta attttttccc caccacctga aattccgtaa ttttaaattg aaactatcta 17100
cagctggcaa aaatcacatc tctcctagag cacaagacaa atcatagtta ctggctattt 17160
gcaatctgaa gcatctcaat atcccacacc tgggattaaa acaaaaacat attcacatca 17220
cataactgtt ttttttttcc aattttttat taggtatttt ctttatttac atttcaaattg 17280
ctatcccgaa agtcccctat accctcccac ctccctgctc ccctacacac ccactcccac 17340
tttttgacct tggagtccc cggtactggg gcatataaag tttgcaagac caaggggcct 17400
ctcttcccag tgatggcga ctaagccatc ttctgtaca tatgcagata gagacacgag 17460
ctctgggggt actagttagt tcatattgtt gttccaccta tagggctcga gacccttca 17520
gctccttggg tactttgtct agctcctcca ctgggggctc tgtgttttat ctaatagatg 17580
actgtgagca tccacttctg tatttgacag gcactggcct agcgtcacat gagccagcta 17640
tatcaggggc ctttcagcaa aaccttgctg gcatgtgcaa tagtgtctgc gtttgggtgg 17700
tgattatggg atggatccac tagttctaga gcggccgcca ccgcggtgga gctccagctt 17760
ttgttccctt tagtgagggt taattgcgcg cttggcgtaa tcatgggtcat agctgtttcc 17820
tgtgtgaaat tgttatccgc tcacaattcc acacaacata cgagccggaa gcataaagtg 17880
taaagcctgg ggtgcctaata gagtgagcta actcacatta attgctgtgc gctcactgcc 17940
cgctttccag tcgggaaacc tgtcgtgcca gctgcattaa tgaatcggcc aacgcgcggg 18000
gagaggcggt ttgctgattg ggcgtctctc cgcttctctg ctcaactgact cgctgcgctc 18060
ggtcgttcgg ctgcggcgag cggtatcagc tcaactcaaa gcggtaatac gggtatccac 18120
agaatcaggg gataacgcag gaaagaacat gtgagcaaaa ggccagcaaa aggccaggaa 18180
ccgtaaaaag gccgcgttgc tggcgttttt ccataggctc cgccccctg acgagcatca 18240
caaaaatcga cgctcaagtc agaggtggcg aaaccgcaca ggactataaa gataccaggc 18300

```



```

gtttccccct ggaagctccc tcgtgcgctc tcctgttccg accctgccgc ttaccggata 18360
cctgtccgcc tttctccctt cgggaagcgt ggcgctttct catagctcac gctgtaggta 18420
tctcagttcg gtgtaggtcg ttcgctccaa gctgggctgt gtgcacgaac cccccgttca 18480
gccccaccgc tgcgccttat ccggtacta tcgtcttgag tccaacccgg taagacacga 18540
cttatcgcca ctggcagcag ccactggtaa caggattagc agagcgaggt atgtaggcgg 18600
tgctacagag ttcttgaagt ggtggcctaa ctacggctac actagaagga cagtatttgg 18660
tatctgcgct ctgctgaagc cagttacctt cggaaaaaga gttggtagct cttgatccgg 18720
caaacaaacc accgctggta gcggtggttt ttttgtttgc aagcagcaga ttacgcgcag 18780
aaaaaaaagga tctcaagaag atcctttgat cttttctacg gggctctgacg ctcagtggaa 18840
cgaaaactca cgtaagggga ttttggctat gagattatca aaaaggatct tcacctagat 18900
ccttttaaat taaaaatgaa gttttaaatc aatctaaagt atatatgagt aaacttgggtc 18960
tgacagttac caatgcttaa tcagtgaggg acctatctca gcgatctgtc tatttcgttc 19020
atccatagtt gcctgactcc ccgtcgtgta gataactacg atacgggagg gcttaccatc 19080
tgcccccagt gctgcaatga taccgcgaga cccacgctca ccggctccag atttatcagc 19140
aataaaccag ccagccggaa gggccgagcg cagaagtggc cctgcaactt tatccgcctc 19200
catccagtct attaattggt gccgggaagc tagagtaagt agttcgccag ttaatagttt 19260
gcgcaacgtt gttgccattg ctacaggcat cgtgggtgtca cgctcgtcgt ttggtatggc 19320
ttcattcagc tccggttccc aacgatcaag gcgagttaca tgatcccca tgttggtgcaa 19380
aaaagcgggt agctccttcg gtcctccgat cgttgctcaga agtaagttgg ccgcagtgtt 19440
atcactcatg gttatggcag cactgcataa ttctcttact gtcatgccat ccgtaagatg 19500
cttttctgtg actggtgagt actcaaccaa gtcattctga gaatagtgtg tgccggcgacc 19560
gagttgctct tgccccgcgt caatacggga taataccgcg ccacatagca gaactttaaa 19620
agtgtctatc attggaaaac gttcttcggg gcgaaaactc tcaaggatct taccgctgtt 19680
gagatccagt tcgatgtaac ccactcgtgc acccaactga tcttcagcat cttttacttt 19740
caccagcgtt tctgggtgag caaaaacagg aaggcaaaat gccgcaaaaa aggggaataag 19800
ggcgacacgg aaatgttgaa tactcatact ctctcttttt caatattatt gaagcattta 19860
tcagggttat tgtctcatga gcggatacat atttgaaatg atttagaaaa ataaacaaat 19920
aggggttccg cgcacatttc cccgaaaagt gccacctaaa ttgtaagcgt taatatatttg 19980
ttaaatttcg cgtaaaattt ttgttaaata agctcatttt ttaaccaata ggccgaaatc 20040
ggcaaaatcc cttataaatc aaaagaatag accgagatag ggttgagtgt tgttccagtt 20100
tggaacaaga gtccactatt aaagaacgtg gactccaacg tcaaagggcg aaaaaccgtc 20160
tatcagggcg atggccact acgtgaacca tcaccctaata caagtttttt ggggtcgagg 20220
tgccgtaaag cactaaatcg gaaccctaaa gggagcccc gatttagagc ttgacgggga 20280
aagccggcga acgtggcgag aaaggaaggg aagaaagcga aaggagcggg cgctagggcg 20340
ctggcaagtg tagcgggtcac gctgcgcgta accaccacac ccgccgcgct taatgcgccg 20400
ctacagggcg cgtcccatc gccattcagg ctgcgcaact gttgggaagg gcgatcgggtg 20460
cgggcctctt cgctattacg ccagctggcg aaagggggat gtgctgcaag gcgattaagt 20520
tggttaacgc cagggttttc ccagtcacga cgttgtaaaa cgacggccag tgagcgcgcg 20580
taatacgact cactataggg cgaattgggt accgggcccc ccc 20623

```

<210> 2

<211> 6708

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: R15/APPA +

intron plasmid with pBLCAT3 vector

<400> 2

```

ggatcccctt tgctatgtag tttttaatgg aaattacaac ccatagtgtg ttgataaata 60
gagagtccctg tttgggttta gcaacctctg tttctcataa actccataaa aacaggaata 120
ctctttgttt ctagcataac caaaagattt agtgaattga aaacaatggt cccttagagt 180
ataggtctaa taaccccgaa aatattacca tgatactgag catttgtaag tatctcatag 240
catgtagtat ccatagtcca tcaatgagag agacatttaa catgattttc attaatcagg 300
tggaagagac atgacaacat tcacaggcac tgcacagaac atagtgggtcc accttgacac 360
tatttcacta aactagggtt atctattttg ttgctttctc taacatctct gcaatgaagc 420
aggtcaacag tgccacatat cctttactta acctaaggaa cacaaaaaat tttctacata 480
tatcctgggt agagagtgtc taaaataagt tttccaagaa tggaaaagaa atgttctgac 540
ttaacaatta agacagtatt tatttaaagc aagaaatatg aggcacacaa gaaaatattt 600
tggaagagaa ccatttggtg aacaatattt caaataaaaa tagacaaaca tagttaattg 660
taaacatat gtttgaccag cccttctttt caatagggtt aatgtgaata aaatgttaaa 720
gattctcttt ggggtggtgc aaattgtcca cgaataagac aaaatataaa aataaggact 780
gagtctcaca aaatgaaaag gaaatatatt cagaaagaga atcttgagag aatgtgttgt 840
cacaaattaa agaaaacctg tgggtgaatga catcctgagg cctgagctat tactgacatt 900
taagataaag gtaactgtat acatttgtcc cattgagggg acaagaaagc tgctctcatg 960
ttcagctcta taattcttgc cttaaacaac ttaaatagaa tgatttaaaa tatggagctg 1020
tccatggacc tttgaaatat aaaatagtca agcaacttat caaggaatta cagattcctt 1080
gatactaaca caggtaaata ccacacgtgt tttgagacta catttgctgg gattttattg 1140
atgtaatagg tcacatgttt ttcgggccaa tgttgctgtt attcgggtac ttcaagagaa 1200
tagtggaac tgatgctatg tattctaggg gtttgaagtg atgtttcatg attgaaattt 1260
gtaaaagaat aacatcatca ttcttaacaa tagaacatat aaagtcacac agaagtgaca 1320
gtgtttaagc tgtactattg atcaaagaaa tttattacct tcagtttcaa tggaaataat 1380
tactgataat acaaacatgt gtgaacacac actaatccta tccaaatgca cagtataca 1440
cagaaaatat tagcaagtag aatgcaatat ttatataacg attgtattta tcaatcaatt 1500
gtatgtatca atatatgggc tattttctta cacatgattt tattcaaatt tactctaata 1560
attgttgaac catttagaaa aggcatctg gcaacttttc cttacctcat ccagctgggc 1620
aaaagtccca gtgtggagta aaggatgcaa gatttctctg tctgttaagt ataaaataat 1680
agtatgaatt caaagggtgc attcttctgc ttctagttat aaaggcagtg cttgcttctt 1740
ccagcacaga tctggatctc gaggagcttg gcgagatttt caggagctaa ggaagctaaa 1800
agccgccacc atgaaagcca tcttaatccc atttttatct cttctgattc cgtaacccc 1860
gcaatctgca ttcgctcaga gtgagccgga gctgaagctg gaaagtgtgg tgattgtcag 1920
tcgtcatggt gtgcgtgctc caaccaaggc cagcgaactg atgcaggatg tcacccaga 1980
cgcatggcca acctggccgg taaaactggg ttggctgaca ccgcgcggtg gtgagctaat 2040
cgctatctc ggacattacc aacgccagcg tctggtagcc gacggattgc tggcgaaaaa 2100
gggctgccg cagtctggtc aggtcgcgat tattgctgat gtcgacgagc gtaccgtaa 2160
aacaggcgaa gccttcgccc ccgggctggc acctgactgt gcaataaccg tacataacca 2220
ggcagatacg tccagtcccg atccgttatt taatcctcta aaaactggcg tttgccaaact 2280
ggataacgag aacgtgactg acgcgatcct cagcagggca ggaggggtcaa ttgctgactt 2340
taccgggcat cggcaaaccg cgtttcgcga actggaaccg gtgcttaatt ttccgcaatc 2400
aaacttgtgc cttaaaccgt agaaacagga cgaaagctgt tcattaacgc aggcattacc 2460
atcggaactc aaggtgagcg ccgacaatgt ctcatatacc ggtgcggtaa gcctcgcatc 2520
aatgctgacg gagatatttc tctgcaaca agcacaggga atgccggagc cggggtgggg 2580
aaggatcacc gattcacacc agtggaacac cttgctaagt ttgcataacg cgcaatttta 2640
tttgctacaa cgcacgccag aggttgcccc cagccgcgcc accccgttat tagatttgat 2700

```


caagacagcg	ttgacgcccc	atccaccgca	aaaacaggcg	tatggtgtga	cattacccac	2760
ttcagtgtcg	tttatcgccg	gacacgatac	taatctggca	aatctcggcg	gcgcactgga	2820
gctcaactgg	acgcttcccc	gtcagccgga	taacacgccg	ccaggtgggtg	aactgggtgtt	2880
tgaacgctgg	cgtcggctaa	gcgataaacg	ccagtggtatt	caggtttcgc	tggtcttcca	2940
gacttttacag	cagatgcgtg	ataaaaacgcc	gctgtcatta	aatacgccgc	ccggagaggt	3000
gaaactgacc	ctggcaggat	gtgaagagcg	aaatgcgcag	ggcatgtgtt	cgttggcagg	3060
ttttacgcaa	atcgtgaatg	aagcacgcat	acccgcttgc	agtttgtaag	gtataaggca	3120
gttattggtg	cccttaaacg	cctggtgcta	cgctgaata	agtataata	agcggatgaa	3180
tggcagaaat	tcgccggatc	tttgtgaagg	aaccttactt	ctgtgggtgtg	acataattgg	3240
acaaactacc	tacagagatt	taaagctcta	aggtaaatat	aaaattttta	agtgtataat	3300
gtgttaaact	actgattcta	attgtttgtg	tatttttagat	tccaacctat	ggaactgatg	3360
aatgggagca	gtgggtggaat	gcctttaatg	aggaaaacct	gttttgctca	gaagaaatgc	3420
catctagtga	tgatgaggct	actgctgact	ctcaacattc	tactcctcca	aaaaagaaga	3480
gaaaggtaga	agacccaag	gactttcctt	cagaattgct	aagttttttg	agtcagtctg	3540
tgtttagtaa	tagaactctt	gcttgctttg	ctatttacac	cacaaaggaa	aaagctgcac	3600
tgctatacaa	gaaaattatg	gaaaaatatt	ctgtaacctt	tataagtagg	cataacagtt	3660
ataatcataa	catactgttt	tttcttactc	cacacaggca	tagagtgtct	gctattaata	3720
actatgctca	aaaattgtgt	accttttagct	ttttaatttg	taaaggggtt	aataaggaat	3780
atgtgatgta	tagtgccttg	actagagatc	ataatcagcc	ataccacatt	tgtagagggt	3840
ttacttgctt	taaaaaacct	cccacacctc	cccctgaacc	tgaaacataa	aatgaatgca	3900
attgttggtg	ttaacttggt	tattgcagct	tataatgggt	acaaataaag	caatagcatc	3960
acaaatttca	caaataaagc	atttttttca	ctgcattcta	gttggtggtt	gtccaaactc	4020
atcaatgtat	cttatcatgt	ctggatcgat	ccccgggtac	cgagctcgaa	ttcgtaatca	4080
tggtcatagc	tgtttcctgt	gtgaaattgt	tatccgctca	caattccaca	caacatacga	4140
gccggaagca	taaagtgtaa	agcctggggg	gcctaattgag	tgagctaact	cacattaatt	4200
gcgttgcgct	cactgcccgc	ttccagtcg	ggaaacctgt	cgtgccagct	gcattaatga	4260
atcgccaac	gcgcggggag	aggcgggttg	cgtattgggc	gctcttcgc	ttcctcgctc	4320
actgactcgc	tgcgctcggt	cgctcggtg	cggcgagcgg	tatcagctca	ctcaaaggcg	4380
gtaatacggg	tatccacaga	atcaggggat	aacgcaggaa	agaacatgtg	agcaaaaaggc	4440
cagcaaaagg	ccaggaaccg	taaaaaggcc	gcgttgctgg	cgtttttcca	taggctccgc	4500
ccccctgacg	agcatcacia	aaatcgacgc	tcaagtcaga	ggtggcgaaa	cccgacagga	4560
ctataaagat	accaggcggt	ttcccctgga	agctccctcg	tgcgctctcc	tgttccgacc	4620
ctgccgctta	ccggatacct	gtccgccttt	ctcccttcgg	gaagcgtggc	gctttctcaa	4680
tgctcacgct	gtaggtatct	cagttcggtg	taggtcgttc	gctccaagct	gggctgtgtg	4740
cacgaacccc	ccgttcagcc	cgaccgctgc	gccttatccg	gtaactatcg	tcttgagtc	4800
aaccgggtaa	gacacgactt	atcgccactg	gcagcagcca	ctggtaacag	gattagcaga	4860
gcgaggtagt	taggcgggtg	tacagagttc	ttgaagtggg	ggcctaacta	cggctacact	4920
agaaggacag	tatttggtat	ctgcgctctg	ctgaagccag	ttaccttcgg	aaaaagagtt	4980
ggtagctctt	gatecggcaa	acaaaccacc	gctggtagcg	gtgggttttt	tgtttgcaag	5040
cagcagatta	cgcgcagaaa	aaaaggatct	caagaagatc	ctttgatctt	ttctacgggg	5100
tctgacgctc	agtggaaacga	aaactcacgt	taagggattt	tggtcatgag	attatcaaaa	5160
aggatcttca	cctagatcct	tttaaattaa	aatgaagtt	ttaaatcaat	ctaaagtata	5220
tatgagtaaa	cttggtctga	cagttaccaa	tgcttaatca	gtgaggcacc	tatctcagcg	5280
atctgtctat	ttcgttcatc	catagttgcc	tgactccccg	tcgtgtagat	aactacgata	5340
cgggagggct	taccatctgg	ccccagtgct	gcaatgatac	cgcgagaccc	acgctcaccg	5400
gctccagatt	tatcagcaat	aaaccagcca	gccggaaggg	ccgagcgag	aagtggtcct	5460
gcaactttat	ccgcctccat	ccagtctatt	aattgttgcc	gggaagctag	agtaagtagt	5520
tcgccagtta	atagtttgcg	caacgttggt	gccattgcta	caggcatcgt	ggtgtcacgc	5580



11

tccatggacc tttgaaatat aaaatagtca agcaacttat caaggaatta cagattcctt 1080
gatactaaca caggtaaata ccacacgtgt tttgagacta catttgctgg gattttattg 1140
atgtaatagg tcacatgttt ttcgggccaa tgttgctgtt attcggttac ttcaagagaa 1200
tagtggcaac tgatgctatg tattctaggg gtttgaagt atgtttcatg attgaaattt 1260
gtaaaagaat aacatcatca ttcttaacaa tagaacatat aaagtcacac agaagtgaca 1320
gtgtttaagc tgtactattg atcaaagaaa tttattacct tcagtttcaa tggaaataat 1380
tactgataat acaaacatgt gtgaacacac actaatccta tccaaatgca cagtgatata 1440
cagaaaatat tagcaagtag aatgcaatat ttatataacg attgtattta tcaatcaatt 1500
gtatgtatca atatatgggc tattttctta cacatgattt tattcaaatt tactctaata 1560
attgttgaac catttagaaa aggcatactg gcaacttttc cttacctcat ccagctgggc 1620
aaaagtccca gtgtggagta aaggatgcaa gatttcctgc tctgttaagt ataaaaataat 1680
agtatgaatt caaagggtgc attcttctgc ttctagttat aaaggcagtg cttgcttctt 1740
ccagcacaga tctggatctc gaggagcttg gcgagatttt caggagctaa ggaagctaaa 1800
agccgccacc atgaaagcca tcttaatccc atttttatct cttctgattc cgttaacccc 1860
gcaatctgca ttcgctcaga gtgagccgga gctgaagctg gaaagtgtgg tgattgtcag 1920
tcgtcatggg gtgcgtgctc caaccaaggc cagcgaactg atgcaggatg tcaccccaga 1980
cgcatggcca acctggccgg taaaactggg ttggctgaca ccgcgcggtg gtgagctaat 2040
cgcttatctc ggacattacc aacgccagcg tctggtagcc gacggattgc tggcgaaaaa 2100
gggctgcccg cagtctggtc aggtcgcgat tattgctgat gtcgacgagc gtacccgtaa 2160
aacaggcgaa gccttcgccg ccgggctggc acctgactgt gcaataaccg tacataacca 2220
ggcagatacg tccagtcacc atccgttatt taatcctcta aaaactggcg tttgccaaact 2280
ggataacgcy aacgtgactg acgcgatcct cagcagggca ggagggtcaa ttgctgactt 2340
taccgggcat cggcaaacgg cgtttcgcga actggaacgg gtgcttaatt ttccgcaatc 2400
aaacttgctg cttaaacgtg agaaacagga cgaaagctgt tcattaacgc aggcatcacc 2460
atcggaactc aagggtgagc ccgacaatgt ctcatcattc ggtgcggtaa gcctcgcac 2520
aatgctgacg gagatatttc tcctgcaaca agcacaggga atgccggagc cgggggtggg 2580
aaggatcacc gattcacacc agtggaacac cttgctaagt ttgcataacg cgcaatttta 2640
tttgctacaa cgcacgccag aggttgcccc cagccgcgcc accccgttat tagatttgat 2700
caagacagcy ttgacgcccc atccaccgca aaacaggcg tatggtgtga cattaccac 2760
ttcagtgtg tttatcgccg gacacgatac taatctggca aatctcggcg gcgactgga 2820
gctcaactgg acgcttcccc gtcagccgga taacacgccg ccagggtggtg aactggtgtt 2880
tgaacgctgg cgtcggctaa gcgataacag ccagtggatt caggtttcgc tggctctcca 2940
gactttacag cagatgcgtg ataaaacgcc gctgtcatta aatacgcgc ccggagaggt 3000
gaaactgacc ctggcaggat gtgaagagcg aaatgcgcag ggcattgtgt cgttggcagg 3060
ttttacgcaa atcgtgaatg aagcacgcac acccgcttgc agtttgtaag gtataaggca 3120
gttattggtg cccttaaacy cctggtgcta cgcctgaata agtgataata agcggatgaa 3180
tggcagaaat tcgccggatc tttgtgaagg aaccttactt ctgtggtgtg acataattgg 3240
acaaactacc tacagagatt taaagctcta aggtaaatat aaaattttta agtgtataat 3300
gtgttaaact actgattcta attgtttgtg tattttagat tccaacctat ggaactgatg 3360
aatgggagca gtggtggaat gcctttaatg aggaaaacct gttttgctca gaagaaatgc 3420
catctagtga tgatgaggct actgctgact ctcaacattc tactcctcca aaaaagaaga 3480
gaaaggtaga agacccaag gactttcctt cagaattgct aagttttttg agtcatgctg 3540
tgtttagtaa tagaactctt gcttgctttg ctatttacac caciaaggaa aaagctgcac 3600
tgctatacaa gaaaattatg gaaaaatatt ctgtaacctt tataagtagg cataacagtt 3660
ataatcataa catactgttt tttcttactc cacacaggca tagagtgtct gctattaata 3720
actatgctca aaaattgtgt acctttagct ttttaatttg taaaggggtt aataaggaat 3780
atgtgatgta tagtgccctg actagagatc ataatcagcc ataccacatt ttagagaggt 3840
ttacttgctt taaaaaacct cccacacctc ccctgaacc tgaaacataa aatgaatgca 3900




```

attgttggtg ttaacttggt tattgcagct tataatgggt acaaataaag caatagcatc 3960
acaaatttca caaataaagc atttttttca ctgcattcta gttgtggttt gtccaaactc 4020
atcaatgtat cttatcatgt ctggatcgat ccccggttac 4060

```

<210> 4

<211> 6116

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: R15/APPA
plasmid with pBLCAT3 vector

<400> 4


```

ggatccccctt tgctatgtag tttttaatgg aaattacaac ccatagtgtg ttgataaata 60
gagagtccctg tttgggttaa gcaacctctg tttctcataa actccataaa aacaggaata 120
ctctttgttt ctagcataac caaaagattt agtgaattga aaacaatgtt cccttagagt 180
ataggtctaa taaccccgaa aatattacca tgatactgag catttgtaag tatctcatag 240
catgtagtat ccatagtcca tcaatgagag agacatttaa catgattttc attaatcagg 300
tggaagagac atgacaacat tcacaggcac tgcacagaac atagtgggtc accttgaca 360
tatttcacta aactaggttt atctattttg ttgctttctc taacatctct gcaatgaagc 420
aggtcaacag tgccacatat cctttactta acctaaggaa cacaaaaaat tttctacata 480
tatcctgggt agagagtgtc taaaataagt tttccaagaa tggaagaa atgttctgac 540
ttaacaatta agacagtatt tatttaaacg aagaaatatg aggcacacaa gaaaatattt 600
tggaagaaa ccatttggtg aacaatattt caaataaaaa tagacaaaca tagttaattg 660
taaaacatat gtttgaccag cccttctttt caataggctt aatgtgaata aaatgttaa 720
gatttctctt ggggtggctg aaattgtcca cgaataagac aaaatataaa aataaggact 780
gagtctcaca aaatgaaaag gaaatatatt cagaaagaga atcttgagag aatgtgttgt 840
cacaaattaa agaaaacctg tggatgaatg catcctgagg cctgagctat tactgacatt 900
taagataaag gtaactgtat acatttgtcc cattgagggg acaagaaagc tgctctcatg 960
ttcagctcta taattcttgc cttaacaac ttaaatagaa tgatttaaaa tatggagctg 1020
tccatggacc tttgaaatat aaaatagtca agcaacttat caaggaatta cagattcctt 1080
gatactaaca caggtaaact ccacacgtgt tttgagacta catttgctgg gattttattg 1140
atgtaatagg tcacatgttt ttcgggccaa tgttgctgtt attcggttac ttcaagagaa 1200
tagtggcaac tgatgctatg tattctaggg gtttgaagtg atgtttcatg attgaaattt 1260
gtaaaagaat aacatcatca ttcttaacaa tagaacatat aaagtcacac agaagtgaca 1320
gtgtttaagc tgtactattg atcaaagaaa tttattacct tcagtttcaa tggaaataat 1380
tactgataat acaaacatgt gtgaacacac actaatccta tccaaatgca cagtgatata 1440
cagaaaatat tagcaagtag aatgcaatat ttatataacg attgtattta tcaatcaatt 1500
gtatgtatca atatatgggc tattttctta cacatgattt tattcaaatt tactctaata 1560
attgttgaac catttagaaa aggcatactg gcaacttttc cttacctcat ccagctgggc 1620
aaaagtccca gtgtggagta aaggatgcaa gatttcctgc tctgttaagt ataaaataat 1680
agtatgaatt caaaggtgcc attcttctgc ttctagttat aaaggcagtg cttgcttctt 1740
ccagcacaga tctggatctc gaggagcttg gcgagatttt caggagctaa ggaagctaaa 1800
agccgccacc atgaaagcca tcttaatccc atttttatct cttctgattc cgttaacccc 1860
gcaatctgca ttcgctcaga gtgagccgga gctgaagctg gaaagtgtgg tgattgtcag 1920
tcgtcatggt gtgcgtgctc caaccaaggc cagcgaactg atgcaggatg tcacccaga 1980

```



cgcctagcca	acctggcccg	taaaactggg	tgggtgaca	cgcgcggtg	gtgagcta	2040
cgcctatctc	ggacattacc	aacgccagcg	tctggtagcc	gacggattgc	tggcgaaaaa	2100
gggctgccc	cagtctggtc	aggtcgcgat	tattgctgat	gtcgacgagc	gtacccgtaa	2160
aacaggcgaa	gccttcgccg	ccgggctggc	acctgactgt	gcaataaccg	tacataacca	2220
ggcagatacg	tccagtcce	atccgttatt	taatectcta	aaaactggcg	tttgccaact	2280
ggataacgcg	aacgtgactg	acgcgatcct	cagcagggca	ggaggggtcaa	ttgctgactt	2340
taccgggcat	cggcaaacgg	cgtttcgcga	actggaacgg	gtgcttaatt	ttccgcaatc	2400
aaacttgtgc	cttaaacgtg	agaaacagga	cgaaagctgt	tcattaacgc	aggcattacc	2460
atcggaaactc	aagggtgagcg	ccgacaatgt	ctcattaacc	ggtgcggtaa	gcctcgcatc	2520
aatgctgacg	gagatatttc	tcctgcaaca	agcacaggga	atgccggagc	cgggggtgggg	2580
aaggatcacc	gattcacacc	agtggaacac	cttgctaagt	ttgcataacg	cgcaatttta	2640
tttgctacaa	cgcacgccag	aggttgccc	cagccgcgcc	accccgttat	tagatttgat	2700
caagacagcg	ttgacgcccc	atccaccgca	aaaacaggcg	tatggtgtga	cattaccac	2760
ttcagtgtctg	tttatcgccg	gacacgatac	taatctggca	aatctcggcg	gcgactgga	2820
gctcaactgg	acgcttccc	gtcagccgga	taacacgccg	ccaggtgggtg	aactgggtgtt	2880
tgaacgctgg	cgtcggctaa	gcgataacag	ccagtggtatt	caggtttcgc	tggctctcca	2940
gactttacag	cagatgcgtg	ataaaacgcc	gctgtcatta	aatacgccgc	ccggagaggt	3000
gaaactgacc	ctggcaggat	gtgaagagcg	aaatgcgcag	ggcatgtgtt	cgttggcagg	3060
ttttacgcaa	atcgtgaatg	aagcacgcac	acccgcttgc	agtttgtaag	gtataaggca	3120
gttattgggtg	cccttaaacg	cctgggtgcta	cgcctgaata	agtgataata	agcggatgaa	3180
tggcgaaaat	tcgccggatc	tttgtgaagg	aaccttactt	ctgtgggtgtg	acataattgg	3240
acaaactacc	tacagagatt	taaaaaacct	cccacacctc	cccctgaacc	tgaaacataa	3300
aatgaatgca	attgttggtg	ttaacttggt	tattgcagct	tataatgggt	acaaataaag	3360
caatagcatc	acaaatttca	caaataaagc	atTTTTTTca	ctgcattcta	gttgtgggtt	3420
gtccaaactc	atcaatgtat	cttatcatgt	ctggatcgat	ccccgggtac	cgagctcgaa	3480
ttcgtaatca	tgggtcatagc	tgtttctgt	gtgaaattgt	tatccgctca	caattccaca	3540
caacatacga	gccggaagca	taaagtgtaa	agcctgggg	gcctaattgag	tgagctaact	3600
cacattaatt	gcgttgcgct	cactgccgc	tttccagtcg	ggaaacctgt	cgtgccagct	3660
gcattaatga	atcggccaac	gcgcggggag	aggcggtttg	cgtattgggc	gctcttccgc	3720
ttcctcgctc	actgactcgc	tgcgctcggt	cgttcggctg	cggcgagcgg	tatcagctca	3780
ctcaaaggcg	gtaatacgg	tatccacaga	atcaggggat	aacgcaggaa	agaacatgtg	3840
agcaaaaggc	cagcaaaagg	ccaggaaccg	taaaaaggcc	gcgttgctgg	cgtttttcca	3900
taggctccgc	ccccctgacg	agcatcaca	aaatcgacgc	tcaagtca	ggtggcgaaa	3960
ccgcacagga	ctataaagat	accaggcggt	tccccctgga	agctccctcg	tgcgctctcc	4020
tgttccgacc	ctgccgctta	ccggatacct	gtccgccttt	ctcccttcgg	gaagcggtgc	4080
gctttctcaa	tgctcacgct	gtaggtatct	cagttcgggtg	taggtcgttc	gctccaagct	4140
gggctgtgtg	cacgaacccc	ccgttcagcc	cgaccgctgc	gccttatccg	gtaactatcg	4200
tcttgagtc	aacccggtaa	gacacgactt	atcgccactg	gcagcagcca	ctggtaacag	4260
gattagcaga	gcgaggtatg	taggcgggtg	tacagagttc	ttgaagtgg	ggcctaacta	4320
cggctacact	agaaggacag	tatttggtat	ctgcgctctg	ctgaagccag	ttaccttcgg	4380
aaaaagagtt	ggtagctctt	gatccggcaa	acaaaccacc	gctggtagcg	gtgggttttt	4440
tgtttgcaag	cagcagatta	cgcgcagaaa	aaaaggatct	caagaagatc	ctttgatctt	4500
ttctacgggg	tctgacgctc	agtggaacga	aaactcacgt	taagggattt	tgggtcatgag	4560
attatcaaaa	aggatcttca	cctagatcct	tttaaattaa	aatgaagtt	ttaaatcaat	4620
ctaaagtata	tatgagtaaa	cttgggtctga	cagttaccaa	tgcttaatca	gtgaggcacc	4680
tatctcagcg	atctgtctat	ttcgttcatc	catagttgcc	tgactccccg	tcgtgtagat	4740
aactacgata	cgggagggct	taccatctgg	ccccagtgct	gcaatgatac	cgcgagaccc	4800
acgtccaccg	gctccagatt	tatcagcaat	aaaccagcca	gccggaagg	ccgagcgcgag	4860



aagtgtcct	gcaactttat	ccgcctccat	ccagtctatt	aattgttgcc	gggaagctag	4920
agtaagtagt	tcgccagtta	atagtttgcg	caacgttggt	gccattgcta	caggcatcgt	4980
ggtgtcacgc	tcgtcgtttg	gtatggcttc	attcagctcc	ggttcccaac	gatcaaggcg	5040
agttacatga	tccccatgt	tgtgcaaaaa	agcggtttagc	tccttcgggtc	ctccgatcgt	5100
tgtcagaagt	aagttggccg	cagtgttatc	actcatgggt	atggcagcac	tgcataattc	5160
tcttactgtc	atgccatccg	taagatgctt	ttctgtgact	ggtgagtact	caaccaagtc	5220
attctgagaa	tagtgtatgc	ggcgaccgag	ttgctcttgc	ccggcgtaaa	tacgggataa	5280
taccgcgcca	catagcagaa	ctttaaaagt	gctcatcatt	ggaaaacggt	cttcggggcg	5340
aaaactctca	aggatcttac	cgctgttgag	atccagttcg	atgtaacca	ctcgtgcacc	5400
caactgatct	tcagcatctt	ttactttcac	cagcgtttct	gggtgagcaa	aaacaggaag	5460
gcaaaatgcc	gcaaaaaagg	gaataagggc	gacacggaaa	tgttgaatac	tcataactctt	5520
cttttttcaa	tattattgaa	gcatttatca	gggttattgt	ctcatgagcg	gatacatatt	5580
tgaatgtatt	tagaaaaata	aacaaatagg	ggttccgcgc	acatttcccc	gaaaagtgcc	5640
acctgacgtc	taagaaacca	ttattatcat	gacattaacc	tataaaaaata	ggcgatatcac	5700
gaggcccttt	cgtctcgcgc	gtttcgggtga	tgacggtgaa	aacctctgac	acatgcagct	5760
cccggagacg	gtcacagctt	gtctgtaagc	ggatgccggg	agcagacaag	cccgtcaggg	5820
cgcgtcacgc	ggtgttggcg	ggtgtcgggg	ctggcttaac	tatgcggcat	cagagcagat	5880
tgtactgaga	gtgcaccata	tgcggtgtga	aataccgcac	agatgcgtaa	ggagaaaaata	5940
ccgcatcagg	cgccattcgc	cattcaggct	gcgcaactgt	tgggaagggc	gatcggtgcg	6000
ggcctcttcg	ctattacgcc	agctggcgaa	agggggatgt	gctgcaaggc	gattaagttg	6060
ggtaacgcc	gggttttccc	agtcacgacg	ttgtaaaacg	acggccaagt	ccaagc	6116

<210> 5

<211> 3470

<212> DNA


<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: R15/APPA
transgene

<400> 5

ggatcccctt	tgctatgtag	tttttaatgg	aaattacaac	ccatagtgtg	ttgataaata	60
gagagtccctg	tttggtttaa	gcaacctctg	tttctcataa	actccataaa	aacaggaata	120
ctctttgttt	ctagcataac	caaaagattt	agtgaattga	aaacaatgtt	cccttagagt	180
ataggtctaa	taaccccgaa	aatattacca	tgatactgag	catttgtaag	tatctcatag	240
catgtagtat	ccatagtcca	tcaatgagag	agacatttaa	catgattttc	attaatcagg	300
tggaaaagac	atgacaacat	tcacaggcac	tgcacagaac	atagtgggtcc	accttgca	360
tatttcacta	aactaggttt	atctattttg	ttgctttctc	taacatctct	gcaatgaagc	420
aggtcaacag	tgccacatat	cctttactta	acctaaggaa	cacaaaaaat	tttctacata	480
tatcctgggt	agagagtgt	taaaataagt	tttccaagaa	tggaaaagaa	atgttctgac	540
ttaacaatta	agacagtatt	tattttaagc	aagaaatatg	aggcacacaa	gaaaatattt	600
tgggaagaaa	ccatttggtg	aacaatattt	caaataaaaa	tagacaaaca	tagttaattg	660
taaaacatat	gtttgaccag	cccttctttt	caataggctt	aatgtgaata	aatgtttaa	720
gatttctctt	gggtggctgc	aaattgtcca	cgaataagac	aaaatataaa	aataaggact	780
gagtctcaca	aaatgaaaag	gaaatatatt	cagaaagaga	atcttgagag	aatgtgttgt	840
cacaaattaa	agaaaacctg	tggtgaatga	catcctgagg	cctgagctat	tactgacatt	900



```


taagataaag gtaactgtat acatttgtec cattgagggg acaagaaagc tgctctcatg 960
ttcagctcta taattcttgc cttaaacaac ttaaatagaa tgatttaaaa tatggagctg 1020
tccatggacc tttgaaatat aaaatagtca agcaacttat caaggaatta cagattcctt 1080
gatactaaca caggtaaata ccacacgtgt tttgagacta catttgctgg gattttattg 1140
atgtaataag tcacatgttt ttcgggccaa tgttgctgtt attcggttac ttcaagagaa 1200
tagtggcaac tgatgctatg tattctaggg gtttgaagt atgtttcatg attgaaattt 1260
gtaaaagaat aacatcatca ttcttaacaa tagaacatat aaagtcacac agaagtgaca 1320
gtgtttaagc tgtactattg atcaaagaaa tttattacct tcagtttcaa tggaaataat 1380
tactgataat acaaacatgt gtgaacacac actaatccta tccaaatgca cagtataca 1440
cagaaaatat tagcaagtag aatgcaatat ttatataacg attgtattta tcaatcaatt 1500
gtatgtatca atatatgggc tattttctta cacatgattt tattcaaatt tactctaata 1560
attgttgaac catttagaaa aggcatactg gcaacttttc cttacctcat ccagctgggc 1620
aaaagtccca gtgtggagta aaggatgcaa gatttcctgc tctgttaagt ataaaataat 1680
agtatgaatt caaaggtgcc attcttctgc ttctagttat aaaggcagtg cttgcttctt 1740
ccagcacaga tctggatctc gaggagcttg gcgagatttt caggagctaa ggaagctaaa 1800
agccgccacc atgaaagcca tcttaatccc atttttatct cttctgattc cgttaacccc 1860
gcaatctgca ttcgctcaga gtgagccgga gctgaagctg gaaagtgtgg tgattgtcag 1920
tcgtcatggt gtgctgtctc caaccaaggc cacgcaactg atgcaggatg tcaccccaga 1980
cgcatggcca acctggccgg taaaactggg ttggctgaca ccgcgcggtg gtgagctaata 2040
cgcttatctc ggacattacc aacgccagcg tctggtagcc gacggattgc tggcgaaaaa 2100
gggctgcccg cagtctggtc aggtcgcgat tattgctgat gtcgacgagc gtaccgtaa 2160
aacaggcgaa gccttcgccg ccgggctggc acctgactgt gcaataaccg tacataacca 2220
ggcagatacg tccagtcctg atccgttatt taatcctcta aaaactggcg tttgccaaact 2280
ggataacgcg aacgtgactg acgcgacct cagcagggca ggagggtcaa ttgctgactt 2340
taccgggcat cggcaaacgg cgtttcgcga actggaacgg gtgcttaatt ttccgcaatc 2400
aaacttgtgc cttaaacgtg agaaacagga cgaaagctgt tcattaacgc aggcattacc 2460
atcggaatc aaggtgagcg ccgacaatgt ctcatiaacc ggtgcggtaa gcctcgcatc 2520
aatgctgacg gagatatttc tcctgcaaca agcacagga atgccggagc cggggtgggg 2580
aaggatcacc gattcacacc agtgaacac cttgctaagt ttgcataacg cgcaatttta 2640
tttgctacaa cgcacgccag aggttgcccg cagccgcgc acccgttat tagatttgat 2700
caagacagcg ttgacgcccc atccaccgca aaaacaggcg tatggtgtga cattaccac 2760
ttcagtgtg tttatcgccg gacacgatac taatctggca aatctcggcg gcgcactgga 2820
gctcaactgg acgcttcccg gtcagccgga taacacgccg ccagggtggtg aactggtgtt 2880
tgaacgctgg cgtcggctaa gcgataacag ccagtggatt cagggttctgc tggcttcca 2940
gactttacag cagatgcgtg ataaaacgcc gctgtcatta aatacgccgc ccggagaggt 3000
gaaactgacc ctggcaggat gtgaagagcg aaatgcgcag ggcattgtgt cgttggcagg 3060
ttttacgcaa atcgtgaatg aagcacgcat acccgcttgc agtttgtaag gtataaggca 3120
gttattggtg cccttaaacg cctggtgcta cgcctgaata agtgataata agcggatgaa 3180
tggcagaaat tcgccggatc tttgtgaagg aaccttactt ctgtggtgtg acataattgg 3240
acaaactacc tacagagatt taaaaaacct cccacacctc cccctgaacc tgaaacataa 3300
aatgaatgca attgttgttg ttaacttgtt tattgcagct tataatggtt acaataaaag 3360
caatagcatc acaaatttca caaataaagc atttttttca ctgcattcta gttgtggttt 3420
gtccaaactc atcaatgtat cttatcatgt ctggatcgat ccccggtac 3470

```

<210> 6

<211> 5421

<212> DNA



<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: SV40/APPA +
intron plasmid

<400> 6

```

cgagatttttc aggagctaag gaagctaaaa gccgccacca tgaaagccat cttaatccca 60
tttttatctc ttctgattcc gttaaccccg caatctgcat tcgctcagag tgagccggag 120
ctgaagctgg aaagtgtggt gattgtcagt cgtcatgggtg tgcgtgctcc aaccaaggcc 180
acgcaactga tgcaggatgt caccgccagac gcatggccaa cctggccgggt aaaactgggt 240
tggctgacac cgcgnggtgg tgagctaata gcctatctcg gacattacca acgccagcgt 300
ctggtagccg acggattgct ggcgaaaaag ggctgcccgc agtctggtca ggtcgcgatt 360
attgctgatg tcgacgagcg taccgcgtaaa acaggcgaag ccttcgccgc cgggctggca 420
cctgactgtg caataaccgt acatacccag gcagatacgt ccagtcccga tccgttattt 480
aatcctctaa aaactggcgt ttgccaaactg gataacgcga acgtgactga cgcgatcctc 540
agcagggcag gaggggtcaat tgctgacttt accgggcata ggcaaacggc gtttcgcgaa 600
ctggaacggg tgcttaattt tccgcaatca aacttgtgcc ttaaactgta gaaacaggac 660
gaaagctgtt cattaacgca ggcattacca tcggaactca aggtgagcgc cgacaatgtc 720
tcattaaccg gtgcggtaag cctcgcatca atgctgacgg agatatttct cctgcaacaa 780
gcacagggaa tgccggagcc ggggtgggga aggatcaccc attcacacca gtggaacacc 840
ttgctaagtt tgcataacgc gcaattttat ttgctacaac gcacgccaga ggttgcccgc 900
agccgcgcca ccccgttatt agatttgatc aagacagcgt tgacgcccca ccaccgcaa 960
aacaggcgta tgggtgtgaca ttaccactt cagtgtgtgt tatcgccgga cagataacta 1020
atctggcaaa tctcggcggc gcaactggagc tcaactggac gcttcccgggt cagccggata 1080
acacgccgcc aggtggtgaa ctggtgtttg aacgctggcg tcggctaagc gataacagcc 1140
agtggattca ggtttcgcgt gtcttccaga ctttacagca gatgcgtgat aaaacgccgc 1200
tgtcattaaa tacgccgccc ggagagggtga aactgaccct ggcaggatgt gaagagcgaa 1260
atgcgcaggg catgtgttcg ttggcagggt ttacgcaaat cgtgaatgaa gcacgcatac 1320
ccgcttgtag tttgtaaggc agttattggt gcccttaaac gcctggtgct acgcctgaat 1380
aagtgataat aagcggatga atggcagaaa ttcgccggat ctttgtgaag gaaccttact 1440
tctgtggtgt gacataattg gacaaactac ctacagagat ttaaagctct aaggtaaata 1500
taaaattttt aagtgtataa tgtgttaaac tactgattct aattgtttgt gtattttaga 1560
ttccaaccta tggaactgat gaatgggagc agtgggtggaa tgcccttaat gagggaaacc 1620
tgttttgctc agaagaaatg ccatctagtg atgatgaggg tactgctgac tctcaacatt 1680
ctactcctcc aaaaaagaag agaaaggtag aagaccccaa ggactttcct tcagaattgc 1740
taagtttttt gagtcatgct gtgttttagta atagaactct tgcttgcttt gctattttaca 1800
ccacaaagga aaaagctgca ctgctataca agaaaattat ggaaaaatat tctgtaacct 1860
ttataagtag gcataacagt tataatcata acatactgtt ttttcttact ccacacaggc 1920
atagagtgtc tgctattaat aactatgtc aaaaattgtg tacctttage tttttaattt 1980
gtaaaggggt taataaggaa tatttgatgt atagtgcctt gactagagat cataatcagc 2040
cataccacat ttgtagaggt tttacttgct ttaaaaaacc tcccacacct cccctgaac 2100
ctgaaacata aatgaatgc aattgttgtt gttaacttgt ttattgcagc ttataatggt 2160
tacaaataaa gcaatagcat cacaaatttc acaataaag catttttttc actgcattct 2220
agttgtgggt tgtccaaact catcaatgta tcttatcatg tctggatcga tccccgggta 2280
ccgagctcga attcgtaatc atggctcatg ctgtttcctg tgtgaaattg ttatccgctc 2340
acaattccac acaacatacg agccggaagc ataaagtgt aagcctgggg tgccaatga 2400
gtgagctaac tcacattaat tgcgttgccg tcactgcccc ctttccagtc gggaaacctg 2460

```



```

tcgtgccagc tgcattaatg aatcggccaa cgcgcgggga gaggcgggtt gcgtattggg 2520
cgctcttccg cttcctcgct cactgactcg ctgcgctcgg tcgttcgggt gcggcgagcg 2580
gtatcagctc actcaaaggc ggtaatacgg ttatccacag aatcagggga taacgcagga 2640
aagaacatgt gagcaaaagg ccagcaaaag gccaggaacc gtaaaaaggc cgcgttgctg 2700
gcgtttttcc ataggctccg cccccctgac gagcatcaca aaaatcgacg ctcaagtcag 2760
aggtggcgaa acccgacagg actataaaga taccaggcgt tccccctgg aagctccctc 2820
gtgcgctctc ctgttcggac cctgccgctt accggatacc tgtccgctt tctcccttcg 2880
ggaagcgtgg cgctttctca atgctcacgc tgtaggatc tcagttcggg taggtcgtt 2940
cgctccaagc tgggctgtgt gcacgaaccc cccgttcagc ccgaccgctg cgccttatcc 3000
ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact ggcagcagcc 3060
actggtaaca ggattagcag agcgaggat gtaggcgggt ctacagagtt cttgaagtgg 3120
tggcctaact acggctacac tagaaggaca gtatttggt tctgcgctct gctgaagcca 3180
gttaccttcg gaaaaagagt tggtagctct tgatccggca aacaaaccac cgctggtagc 3240
ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc tcaagaagat 3300
cctttgatct tttctacggg gtctgacgct cagtggaaacg aaaactcacg ttaagggatt 3360
ttggctcatga gattatcaaa aaggatcttc acctagatcc ttttaaatta aaaatgaagt 3420
tttaaataca tctaaagtat atatgagtaa acttggtctg acagttacca atgcttaatc 3480
agtgaggcac ctatctcagc gatctgtcta tttcgttcat ccatagttgc ctgactcccc 3540
gtcgtgtaga taactacgat acgggagggc ttaccatctg gccccagtgc tgcaatgata 3600
ccgcgagacc cacgctcacc ggctccagat ttatcagcaa taaaccagcc agccggaagg 3660
gccgagcgca gaagtggtec tgcaacttta tccgcctcca tccagtctat taattgttgc 3720
cggaagcta gagtaagtag ttcgccagtt aatagtttgc gcaacgttgt tgccattgct 3780
acaggcatcg tgggtgtcag ctcgtcgttt ggtatggctt cattcagctc cggttcccaa 3840
cgatcaaggc gagttacatg atcccccatg ttgtgcaaaa aagcggttag ctcttcgggt 3900
cctccgacg ttgtcagaag taagttggcc gcagtgttat cactcatggt tatggcagca 3960
ctgcataatt ctcttactgt catgccatcc gtaagatgct tttctgtgac tggtagtac 4020
tcaaccaagt cattctgaga atagtgtatg cggcgaccga gttgctcttg cccggcgta 4080
atacgggata ataccgcgcc acatagcaga actttaaaag tgctcatcat tggaaaacgt 4140
tcttcggggc gaaaactctc aaggatctta ccgctgttga gatccagttc gatgtaaccc 4200
actcgtgcac ccaactgatc ttcagcatct tttactttca ccagcgtttc tgggtgagca 4260
aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggc cgacacggaa atgttgaata 4320
ctcatactct tcctttttca atattattga agcatttatc agggttattg tctcatgagc 4380
ggatacatat ttgaatgtat ttagaaaaat aaacaaatag gggttccgcg cacatttccc 4440
cgaaaagtgc cacctgacgt ctaagaaacc attattatca tgacattaac ctataaaaaat 4500
aggcgtatca cgaggccctt tcgtctcgcg cgtttcgggt atgacgggtg aaacctctga 4560
cacatgcagc tcccggagac ggtcacagct tgtctgtaag cggatgccgg gagcagacaa 4620
gcccgtcagg gcgcgtcagc ggggtgttggc ggggtgtcggg gctggcttaa ctatgcggca 4680
tcagagcaga ttgtactgag agtgaccat atgcggtgtg aaataccgca cagatgcgta 4740
aggagaaaat accgcatcag gcgccattcg ccattcaggc tgcgcaactg ttgggaaggg 4800
cgatcgggtg gggcctcttc gctattacgc cagctggcga aagggggatg tgctgcaagg 4860
cgattaagtt gggtaacgcc agggttttcc cagtcacgac gttgtaaaac gacggccagt 4920
gccaaagctt acactttatg cttccggctc gtatgttgtg tggaaattgtg agcggataac 4980
aatttcacac aggaaacagc tatgaccatg attacgaatt cggcgcagca ccatggcctg 5040
aaataacctc tgaaagagga acttggttag gtacctctg aggcggaaag aaccagctgt 5100
ggaatgtgtg tcagttaggg tgtggaaagt cccaggctc cccagcaggc agaagtatgc 5160
aaagcatgca tctcaattag tcagcaacca ggtgtggaaa gtccccaggc tccccagcag 5220
gcagaagtat gcaaagcatg catctcaatt agtcagcaac catagtccc cccctaactc 5280
cgcccatccc gcccctaact ccgccagtt ccgccatc tccgccccat ggctgactaa 5340

```


tttttttttat ttatgcagag gccgaggccg cctcggcctc tgagctattc cagaagtagt 5400
gaggaggctc gaggagcttg g 5421

<210> 7

<211> 17732

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Lama2/APPA
transgene

<400> 7

tcgagagtat ctttgtcagc tgtgcctcca acaaaggggt actgttgccc acatagaaaag 60
atctaaacta attaatat cctcaccgcg caaatctttc agtcactaag ttagcacgat 120
tgttgaacaa gttctccaaa ggagagatac agatgagtgc gtataggggtg gacctggctg 180
ctgaggagac acctgcatct gactaagaag agccacgggtg ttagttgaat ggtgtggagt 240
aggggtgggtc tgtgggacag tagaaaatcg agaggcatgt gccgtttagt gaactgatgg 300
aagctacccc aaacgacaga gattgtcagt caggccaatc cgtttcgagt ttgatgggca 360
gccggacagt gagacagaca cacctactca gttggaggaa ggatgagaac aatggccagc 420
agggattgag agaccctgac aggcgcaagg ccctaacaca cacacctacc acctcacttg 480
acaaagctgc caaagaccaa agacttggtc tccattagaa atgacagctg gcttgacccg 540
acagcataat aagcagagtg tactctgatt ggagaacttt aatgtgtttc attcagtatt 600
ataaaaaggac agtattacag attttgttgt acactgctgt tacatgtggg gcagtgtgtc 660
tttaagtagg gtaaagtact ctttaaaaat gggtcctaga tattttttcc ttttaactcaa 720
gtctcttact gtttaaatga tttttatttt gtttaatatg gaggaaaaag aagcgtaaat 780
ggacaatata tatttagaga aagatggta gctgtcagaa aaatatgcaa atcaaaatca 840
caccaagact gcagcacacc cctgtcagat ggctgtgatc aagaaaataa atgacaatga 900
gtggtggtga agatgtacta aagggaaaca cacacacaca cacacacaca cacacacaca 960
cacactggag caaccactgt ggaaatcagt atgaatggtc ctcaaaaacc tgaagataga 1020
gcggggcgtg gtggcataca cttttattcc cagcactggg gaggcagagg caggtggatc 1080
tctgagttcc aggccagcct ggtctatagc acaggttcta ggacagccag ggctacacag 1140
aaaaaccctg ccttgattaa accaaaccaa accaaaccaa accaaaccaa accaaaccaa 1200
accaaaccaa accaaaccag accaaaccaa aacactgaag atagaacttc agtattccat 1260
tcctagatat atacccaatg gagactaagt cagcaagaca cctgcacagc catgttcact 1320
actacactgt tcaccacagc caggctgtgg aaccagcctg agtgtccatg ataaatgaat 1380
ggataggtaa ctttcaaggt aaatggactc tgctgtgtac atgcctcaca ttctgtttat 1440
tcatttttct ttatgagggt tccattcagg agtcacatgg tagttctatt ttcagtcttc 1500
tgaagatact acactgggtc ccacagttta cacttttatc agcagtgaat aagggttcct 1560
ctatccttac catcatttgt tgtaattttt cttgatgacc ctctttctga cagggatagg 1620
atgtaatatc agtgtgagga agtacaactt gttttctaag tatttattgg ccccttgcat 1680
ttcttctttt gaaaactgtc ggttcctgac atctgctcag gtattcattg gatgttgttt 1740
ctttggtgtt tgagttctta tgaattctag atgttaaate cctgcctgtg gttctctccc 1800
attctgtagg ctgcctctc accctggcaa ttgttgtcct tgttttgag aaacttttga 1860
cttcatggaa tctcatttgt cagttttccc tctctgcta tagcctgagc taatgcactg 1920
gtttttacag agccctgggtc tatgccttta tctcctctg gcagcttcgg agtttcattt 1980
cttacattta gatctttgat ccactttgaa caagttttgg agcagggtga gagatacgaa 2040


tctagttcca	ttcttccata	tgtgatccta	gtttacatag	catcgttggt	tgaagaggtt	2100
ttatttttatt	tttaaataat	gtgtcataaa	aaacgaggtg	gttgtagcag	tgtggatttg	2160
tttctttgtc	ctttgatcta	caggtcttgt	tttgtgtcag	tctcatgatg	ttttattgct	2220
atggctctgt	catacagtct	gaggtcaggt	attgtgatat	accttcagta	ttgctccctc	2280
agactcaggt	ttgctttggc	caggagtcat	cttactcagt	gctcttagag	ctcccccagc	2340
atgtagctgc	tactattctt	agttgataaa	tcaggaaact	ggggctcaga	gagattaact	2400
gtcttgaact	acttctgggg	aggtgaaacg	tggagacact	aaactgtgtt	taccctgtac	2460
tgctccagta	gctgtcgggt	gctgggctac	agcaaagcac	ctatactata	tattactcag	2520
gaggtggaaa	aactcagcct	cccttgggggt	tcccaagctc	ccaggtgtcc	agtcactgct	2580
ggaaacctca	tggagtctga	aaggaagggt	tgaggggtaca	tggggcagcg	atgaggagcc	2640
tggggctggg	atctcccaaa	cacctggata	tccagatgcc	actgggtcag	ggggagttag	2700
gaacagagtt	gggatgtcca	tggacctgtg	acaaggccag	ggccaggggg	aggataactc	2760
tggctttact	aatttgcgaa	agtccttagc	ttagcagcag	ttgtctggga	gcacagaggg	2820
gccttctgta	agaggctcag	gcagtccgc	tctgtaggcg	aagggtcttct	ccatgttccc	2880
catggtgggt	cttgatgaaa	gagacagtcc	ttggctccaa	actggtttat	tgattgttca	2940
ttgtggaaaa	tgggtgcaca	ccaccttctc	aggggtggacc	agagatcaaa	taccttttgc	3000
agggaggaat	atctgggaag	ggacgcttac	tggctaaacc	ctcagggcct	ctagatacat	3060
cattagcatg	gagaactctg	ttctgggcta	catgaccaca	ggccacattt	ccacaagcca	3120
catgtgggaa	gtgtggcaca	tgttctaggc	caggaatctg	gtagggagcg	tggagccacc	3180
taccatccca	ggtgggtgcc	tgggtgccag	ggacctgaa	cccgtcaac	cttaccaagt	3240
ttcctggcag	ggtccactgt	cctacacaga	agctggagga	ggtgtgaggg	ttgtgtcttt	3300
gtggaatgtc	ccatgctgct	tggggctcag	tttctccacc	tgtacctcat	tggtttgggt	3360
ataaaaagtg	gggatacttt	attattctct	gactcggtec	tgaggaaaaa	gcacgtggc	3420
agtcaggaa	ccacacctg	aggttcctgc	actgaaggga	ctccctaagt	ctctggagtc	3480
tctccccttc	acagagctgc	caaagtctag	gttcttttga	ggataacaga	gccatgcttg	3540
gtaagcagac	aacagcattt	gtttactcaa	ccttcttttg	tcagctccct	cttcataaac	3600
aagttgagac	accatgctgg	cttgaggaag	acttctaaag	ccagacaact	gtgcaaggaa	3660
gaagaagaag	gggcaagtgg	agttagcctg	gatgtagccc	tcaaagtctc	cagagaccag	3720
ccatgaaggc	tcaagtggag	ggcaagacct	gcagcagcca	agcatctggc	aggagaggat	3780
cctgggaacc	cctctaccat	gacacacatt	cttctgcag	gtcacactta	ataggccatt	3840
tcttatttgg	atctatcatg	gtgttctgtg	cgagattaat	gaggtgttat	gctgcgaaca	3900
gaaagttata	taaaaacaag	ttcccccccc	ttgtcactgc	tgctaagaat	gtagcagaaa	3960
ttgtctcaag	tgtctctcta	atcagaaaaca	ataaaggctc	ccttggattc	aagccctcca	4020
gtttctctct	tccttgetga	gccttggaca	cccatacaaa	cctcctggat	gctacagctc	4080
tgggcagaga	ctccaagggtg	gggagagact	gatggtacaa	aagcaaaaata	cttgtttggg	4140
ggtacaccca	ctcctctgcc	tgtgtgggtc	ctgcagtcag	tcctgcagac	aggccctcag	4200
tgggtcttcc	atgggcaaca	cgcagaggga	ggcaatggat	gggaataccc	acaccctggg	4260
tagtttacc	cggccatgct	ctctgctctt	catccctect	ctgccctctg	ccacggcttt	4320
ctctgcagga	atcatatctt	catattggcc	cacagggtgtt	ctcctcacc	tagctatgat	4380
gtttacttta	gagtgaacct	agcagggtctg	gtgggaatga	gttctagaag	gctcacggag	4440
atgctagggg	agaaacgtct	tctaactact	gaggttacta	agttcctggg	ggttgtctct	4500
gcctttccct	tgttaaagtc	accttgaagt	tagtgcagaa	gaaatcagag	cccagtcaca	4560
gagtaaatat	ggtcctgaag	atttcctttg	agtgcccaga	atccatgaca	tttcaagagc	4620
cctctttgta	ccttaagtea	tttgggggtg	tatcttctgc	ttgatgtatg	tgtgtgtgtt	4680
tatcaaagag	tgagatgggt	acataagagg	tgctctaaag	gacagagagg	atttgcaatt	4740
gtggcatgtg	acatcctcag	gccttgctct	ggtgccagga	ggaactgatg	cagaaaagag	4800
taagaggtca	tttcttggag	gctgtcacta	tagaggagat	cttacagtgc	attccctcct	4860
ccaggccctg	cctgaggata	gacatgtgct	gactgcaact	gaaacagagg	cttgggatgg	4920




agagtttaggt tcacagaagg gaggggtggga gatggatgct tgctgggttc tgggtctcat 4980
 caccagctcc tgaccacccg gtcagcccat gtgcttattc catagctttc ttttgctatg 5040
 tttactcagt gtggtgtttg ttgggaccca gcagaagcca gtcccaggct gacagctgtg 5100
 gatacacagg gcagcatgag ggtcctcagc ctgaagcagt caggctggca gaagagaaag 5160
 accagcacac attccttcaa ccaactatgt cttgaaaaac aaacatatta tatcacatat 5220
 attgcattta tgagacagct aaaatgtact cgggtagcat gactccagggt ggggatatct 5280
 gcaagtgcc a tgagtggcag agggacagcc aatgtgaggc aagaaggaat tctggctcaa 5340
 cacagcttag ctccctgggt ttggttcaaa ctttgagagt ttgaccacaa gcactttatt 5400
 tttgacatat ttaaacagag cacaactttg ggaaaaagtt ttcttatgaa aattatcaca 5460
 ataaagctta aggcattgact acattaaaat gcctttgcaa agtatatgtg cctctctcca 5520
 caagaatggt tctattgact gagaaataat gttcaggata aagatccagg aagaaaagat 5580
 cagggataag taaaataacta aactcctttg caaagtacat agaccctctt tcataacaat 5640
 gggttctatt gactgacaag cactgctcag gagttgggaa agagtctagc ataagcacga 5700
 tagcctggag actctagtga ggtctagtct tacagacagc aaaaatcacc aggttacaaa 5760
 ctacattcat ttccagtttt ctgatcagga acaggatga atcccttctg ttgaagagaa 5820
 aagtccatgt gtttaaaata tctggtttct ccagtgtat tagcgagaag acttgagccc 5880
 tatacaactc ccacctggag tgacatctg tcttcattgt atattacata cctagacacg 5940
 ctcatctcac agacttagga ctttgtcttc tgatctccat ttctgatccc acttccacct 6000
 ttgccttgat agtgtcattt tcttactgc cttggtgaca accatgttat cctctgtgta 6060
 tttgagtgtt accattttca gattttacct gtatgcaaga tcacacagtc tttgtctttc 6120
 tgtctggatg catgtctaac tctacacaac aacccttccc cgtcactcag atcttctctc 6180
 attaacacat acatggtgct gaagaggcta gggagcttcc cttcagtggg gagctagctg 6240
 gctattgggc ctttttgact gtccaggaag gcccccaatt gctgagacaa gaacttagat 6300
 tcttcattat tgactctaac tcatgtatca agcagaagct aatgaatagt tatcaacagg 6360
 atcagagggt ccagtgtgag acactttgac atgaaagaac ggaggaagga cagatggatg 6420
 cataaaagca ggaccactgc cccaggaagg tccctggaaac tgatgcaggg caaaggacag 6480
 gttataaacc aaatcttagg gagtcaggaa gagcacagag gagctcaacc aactgaccac 6540
 tgcttagggg ctaccaaccc aatcctccct gtgggaacag ctaagctatc agccaagggt 6600
 aataaacagg caggacctgt ggatgacatg gagagcatag ggaccctggg tccagccttt 6660
 agcacctgca ctctcaggat actccaccat tgtgtcttag agagcctagg gatactgggt 6720
 ccagcctttg gtaccttcac tctcagggtta ccccatcact gtgtcttgga gagcctaggc 6780
 accctgggtc cagccttcag tacctgcgct ctcaggacac cccaccattg tctcttgccc 6840
 cgtctcttct tccctctcct ccctttcatt gtctcttctc tgtttctttc ttgactctcc 6900
 tttccctca caccctcact ctagtctctc ccttccctct ctgcatcacc ctattctctc 6960
 tgtgttccct ccactttcct ttatctctca tgcctctctc ctccctcaa tacttgctac 7020
 ccactatact tcaggggcca gctctagtga caaagctgtt aatagcaaga ctctcagatc 7080
 tccaacggct cagaggagcc agaccacca agaactctct ccagggtcaa tttcagggtc 7140
 cttcgaaagc tttcagcaaa tgctcaggga acatgccact aacaagaaga tgcaaatcc 7200
 agttgagagt gggaaaggcc cttgcgtagg tcccatcttc caggccaagg tcagaggggc 7260
 tctgtgtaac ccgattgac agggctcaga acaatgtttt gtttttaagg tttatttatt 7320
 ttaggtgtta gtgtctttgc ttgcatgacc ttatgtgcat catgtgtgtg caggttcctg 7380
 atgacagtag aggagggtt tgaatccctg gggataggaa gttacaggaa attataagct 7440
 gctttgtggg tcttctagct ttcccaacag aagtgaatgc tcttcaccac tgagccatct 7500
 ctctaggccc aagagacatt gctttatgga tataattgtg tgtgtgtgtc aacattgagg 7560
 aaagggaat aaaaaaaaaa cttcagccgc taagggtgta cagtttctact aattgtact 7620
 tttagttgtg ataaaatggc aggtgcttca acatttatat atacaaaaac tccctgctg 7680
 gtggttcaac tgtgagaact ggggtaagtg ggtgagttct ctttttctgt ctctgtctct 7740
 gtctctctcc ttccattctt tcttaaagga aataaacatt gcagctgggt tatagctcat 7800

caatatggaa gttacagaag tgaaaaaagg cattgccttg gtgggtgggtg ttaccagctg 7860
 attttttggtt gtcctgcaag gaggtctggg gactggctgc tctgtctctg tctgtatgag 7920
 tgaggggaagt ctggggagca gattccctaa ccttcagcct ggcttggttc ctgagtgaac 7980
 ccagcctctc tggctcctagt agctttttcc aaacaggaat ctgagtgggtg acagggaaca 8040
 agtaccagcc cattgcttaa gtgccagggt tagtgagggc aggaagctgc catagctggg 8100
 attagtagtt gtattggatg taggaagtcc tatcctggga cagctaatacc ttaatgcttc 8160
 actggagatt ttcaatgaga aatttatccc acggcccata tggccccatc cttttgtctc 8220
 caacagccaa gtattttcca ttagaggaga cttcctgtac acttgatgga tgctcattcc 8280
 aaggtagctt ggggcagtca gtacagactt gggatgacct ctgacagcct aacctctccc 8340
 caacaagggc cctctatgtt tgctatgtaa tgtaatgtca gacattgtca ggagtgtccg 8400
 cagcacagcc tgcccagtggt gagggctctc atagggtttcc cactgtctta tctacacagg 8460
 gataacgagg aggtaagctg cagttcccag tctcacttca cagaggaaga gataacccca 8520
 tcccagggtca tgtagccagc agtggaaaga atgaggattt gaactcaggt cttccaagtc 8580
 ccattgatag catctcctca caagtccctt gccaccctca cgatgcctta gacacttgcc 8640
 tgccctttat actaaggaga tgcagggtaca aggggtttac ccatgtagca gctgaggcag 8700
 ctggggatag ataccagcag caggcctgat gtcaccactc taactccagc atccccagtc 8760
 tgtgttcctg gagtgtgaaa atccctactt aacaagattg tgcaacagtc cttggctctg 8820
 tgacccatag ctggaaacag gattctcatt gatttgtgga acatggtggc agccagccaa 8880
 aaagaggggtc tgcatacaga agacacgtgt ggcaaggcca cagcagactc tgactacctt 8940
 agcttacaga attacaaggt cataatgtcc tctgctttgg tcacctcatg ttaaggacag 9000
 gccctaataga agatggggca gaagactgaa ggaatggcca accaataact ggcccaactt 9060
 gagacccatc ctacaggcaa gcatcaattc ctgacactac taatgatact ctgttatgct 9120
 tgcagacaga agcctagcat aactatcctc cgagagggtcc acccagcaac tgactgaaac 9180
 agaaaaagat atccacaggc aaacagtggg tggagggtcag ggactattat gggagagctg 9240
 tgggaaggat taaaaaccct gaaggggata ggaacccac aggaagacca acagagtcaa 9300
 ctaagagacc tgtgggagct ctcagagact gagccaccaa ccaaagagca tacacaggcc 9360
 ggtccgaggc acctggcacg tgtgaagcag acatgcagct cagtctccat gtaggtcctc 9420
 caataagcgg tagcctgact gcagtatcca atccccaa ca gggctgcata gtctggcctc 9480
 agtgggggag gatgccccta atcctgcaga gacttgatga gtggagagct atccaggggg 9540
 aaccaccct ctctgagaag ggaatgggga tgggggaggg actctgtgaa gaggggacaa 9600
 ggacaaacaa gaacctcaaa taggtcaggc cctaaaggct tgctaagtag cagtggccca 9660
 gctctgtcct gttcctcagc ccaaggctca gctcccacct gtttctgtgt ttttctggct 9720
 tttcatgggc ctaggacttg gtgaccagtt caaacaatgg ggctgtgga agacacaata 9780
 tacaagacta gggacattcc tgttctgctg actatccata gcctgatgta ggtggaagga 9840
 cccaatcact ggatttctac ccttgacaaa ccttgacagc tgagggcctc tcagaaacct 9900
 atttcttcca ctgaaaaatg agactctcaa atgaacgtcg tgacaatcat caggcttatt 9960
 aaagaggtgt atctaacctg aatggcaagc agacagcagg caaatgtctg tatcaacctc 10020
 taggaaggac aagaactgct cactgctgcc cccaggagg ccatttgctg aaacagctgc 10080
 tctcctgctg gtgcacaggc cctgccttct cattgcagcc acagccccct cctgtctgaa 10140
 cctcctgtca ggtcactggg aaacagatca agatggaaca ggacagctcc tgatggtaaa 10200
 taaaaaacag tggatcatggc tattcatagg ggtttatgct tcttcagtc acactgtgaa 10260
 gagctgtggg catgaaccac agtggttcgag gtagagttgg ggttctgaaa ttcacagtgg 10320
 ggtgagctca gtaaagtga gctggagggtc actcgtgaga cacacagtcc tgctgcttct 10380
 gttcccaata tcctgaggag acgacacatc tactttgttc agaggccaca gtctagttga 10440
 cctgagagtt accagtttct tatttgtgtg tgtgtgtgtg tgtgtgtgtg tgtgtgtgtg 10500
 tgttgttcgt gtgtgagtgc aggtgcacat atgatagcgt acacgttgag gtcagaggat 10560
 aactatcagg cgttgtcccc tctactttt cctcggactc tggagaacaa acatgggtcc 10620
 ttattccagg ggagcaagtc gctgttggct gacacatctt gctcacatac attttaccta 10680

gacaaattggag	cctccatcag	agtattactt	tagctcctca	ccgatggcaa	tgcaccacct	10740
ctctacccac	ataggagttg	ggtctccaca	cacccccaca	cccccttcac	caaaacgttt	10800
tcagttactt	tatctggtaa	agttcatcag	agaatgaagc	cagtattaag	aacatggaat	10860
catttgggaa	cctggatcta	gcaatacccc	accctagatg	gagttgctga	gttttcacct	10920
cagattataa	ttcccccta	gcttctatgg	tttattctga	aaccagggga	actcgattcc	10980
tccttttgga	ccacagacat	cctggcttgt	gaattcacat	gtcatctact	gctaatccat	11040
tggtagtatg	tggctcacag	agacacacta	cagtcatggc	caatgtcaag	gtaggacaga	11100
tgtgaatcat	tccccagtc	ctgctgtttt	catgactaac	cctcctcagc	acagtgacca	11160
tgaacctact	tttccctcc	ttttattttt	agaattgctg	gaatttttcta	ttttgagaaa	11220
taatagcctt	gggcagcatt	aaacaaaatc	atctagaaaag	ctggtttaaa	atacagatgg	11280
ttgagtcagt	gaaagagtga	ggaatgtcat	tattggcccc	tcacagaggc	tggtctactc	11340
cagcagaggt	ggttgaagct	cttggacacg	ggtcaggtgc	ataggaaaagg	tngtctggga	11400
cactgagaac	cacaattgaa	caaacagaac	tgttggcttt	ttttttttta	aatgagttct	11460
caaaaaatga	ctggctagct	taggcaaata	cttcgagcca	acccaacaga	acattcttcc	11520
attgattcat	tctggatctt	ctttctagac	aatactgaac	tgaccctctg	ttggcagtcct	11580
caagtttgac	aacatagggc	tttgaacttg	gcacaaggtc	catcactgtc	acccaagcat	11640
cctgggtgac	ctttgggttg	gaatatcttg	gctaacctta	gatattttct	ttggagtatc	11700
tttagaacat	ccaggaaata	gggcttgatt	ctcatcctgg	gaccacaata	taagtcaccc	11760
tagaatccca	ggagatcgtg	cagagaaaca	aggatctctc	tcgtgtgcat	ccttcttcaa	11820
agcagtgagt	agtgactcca	ctaaactgag	ttcccatctg	agagtcacaca	ggaggccttg	11880
gggcaagaag	cagagggaag	gcactgtttg	tgttggtaaa	gttttgactc	taacaaattt	11940
gaagacatag	atgacattgt	gtcagactaa	caacaaccta	gactcatgtg	ggttctgttt	12000
agggatcaga	ttttattcat	caatgacttg	tcttagtgta	tagagaaagg	cttctacttg	12060
gagtgtaggc	tcaataatga	cagaagagat	agctatttcc	cctagggact	gtgctgctcc	12120
aagtttggtg	gagaaaaggca	gtggggaacc	tagatgtgct	ctctggggag	ggggtctgaa	12180
gctggcttca	tagaagggtg	gaagttttgc	tgaacatct	aaacagaatt	atagcttagg	12240
aaagtgagca	ggcaaggcag	ggaatgtgtt	gcatatgtat	atgtacatga	atatattatg	12300
ttatagatac	acacacattt	gaacctcatt	tgcagatgac	agaaaatagg	ttattttgcc	12360
tctcttaact	gctaagcaca	atgacttcca	gttccatcca	tttcttgaaa	tgccacaatt	12420
tcatttttca	ttgtggctga	ataaaattcc	attgcagact	gggccctact	tcatccactc	12480
ctgagggcag	gcatatcccc	tggctccatt	tcttacctat	tgtgaagaga	agtgcaactg	12540
tcttgttgaa	aggcaagcgt	gagagaggca	ggcactaatt	gtgggttttt	gtttcttctt	12600
cctgctatga	ctctccattt	gtcagaacca	aagatcgata	aaagccgcca	ccatgaaagc	12660
catcttaatc	ccatttttat	ctcttctgat	tccgttaacc	ccgcaatctg	cattcgctca	12720
gagtgagccg	gagctgaagc	tggaaagtgt	ggtgattgtc	agtcgtcatg	gtgtgcgtgc	12780
tccaaccaag	gccacgcaac	tgatgcagga	tgtacccca	gacgcatggc	caacctggcc	12840
ggtaaaactg	ggttggctga	caccgcgcgg	tggtgagcta	atcgcctatc	tcggacatta	12900
ccaacgccag	cgtctggtag	ccgacggatt	gctggcgaaa	aagggctgcc	cgcagtctgg	12960
tcaggtcgcg	attattgctg	atgtcgacga	gcgtaccctg	aaaacaggcg	aagccttcgc	13020
cgccgggctg	gcacctgact	gtgcaataac	cgtacatacc	caggcagata	cgtccagtcc	13080
cgatccgtta	tttaatcctc	taaaaactgg	cgtttgccaa	ctggataacg	cgaacgtgac	13140
tgacgcgata	ctcagcaggg	caggagggtc	aattgctgac	tttaccgggc	atcggcaaac	13200
ggcgtttctc	gaactggaac	gggtgcttaa	ttttccgcaa	tcaaacttgt	gccttaaacg	13260
tgagaaacag	gacgaaagct	gttcattaac	gcaggcatta	ccatcggaac	tcaagggtgag	13320
cgccgacaat	gtctcattaa	ccggtgcggt	aagcctcgca	tcaatgctga	cggagatatt	13380
tctcctgcaa	caagcacagg	gaatgccgga	gccgggggtg	ggaaggatca	ccgattcaca	13440
ccagtggaac	accttgctaa	gtttgcataa	cgcgcaattt	tatttgctac	aacgcacgcc	13500
agaggttgcc	cgcagccgcg	ccaccccggt	attagatttg	atcaagacag	cgttgacgcc	13560




ccatccaccg caaaaacagg cgtatggtgt gacattaccc acttcagtgc tgtttatcgc 13620
 cggacacgat actaatctgg caaatctcgg cggcgactg gagctcaact ggacgcttcc 13680
 cggtcagccg gataacacgc cgccaggtgg tgaactggtg tttgaacgct ggcgtcggct 13740
 aagcgataac agccagtggg ttcaggtttc gctggtcttc cagactttac agcagatgcg 13800
 tgataaaaacg ccgctgtcat taaatacgcg gcccgagag gtgaaactga ccctggcagg 13860
 atgtgaagag cgaaatgcgc agggcatgtg ttcgttggca ggttttacgc aaatcgtgaa 13920
 tgaagcacgc ataccgctt gcagtttgta aggtaccgg ggatcacaac ttgccctctg 13980
 aagaggaaga acagaaggat gccacaactc tcctgctggc tactctccag tggtttcac 14040
 ttacttctga tggcatttcc ctctagaaag tgctactatc atccacacat ttctacctga 14100
 gaccacccaa aggaccctcc caaattctct tcctctctga gtagtctcca cacctgttac 14160
 caccatccca gaattaaaat cctaactgca ctctggcggtg tgacttgccct cagtccttgc 14220
 aataagagtt gttggcagtg ccaggcggtg tggcgcacgc ctttaattcc agcacttggg 14280
 aggagaggc agggcgattt ctgagttcga ggcagcctg gtctacagag tgagttccag 14340
 gacagccagg gctatacaga gaaacctgt gtcgaaaaac caaaaaaaa aaaaaaagt 14400
 gttggcagag tgtgggttat ataccaggtg gagatttcaa atgagtggct gaagctgtag 14460
 ccagaaggaa cttagaggat agctcataac ttaaaaagaa atgtagagag tagcagaaac 14520
 attgagagag tgggcacaca gccactgtgt gaatgtggca gaacacaatc cagccagcta 14580
 tacatgcata agtgatatatt ggcgccatcc tgactgatga gacacaggaa aacagataga 14640
 cgggggttagg tggccatggc ctttctgcc tgctcttcc taagggtcat ctcaagacct 14700
 tatgctctct taactcttcc attgctactt agcttctaga tatcacctcc agattagtct 14760
 ctttgggtac atcagtgatc ctggtgatat ccagggtctc ctgattccat ctttgtcata 14820
 gaggtgcaa cttaaagaggt cttcttaata cttcacaccc tgatgccaaa aggaagacac 14880
 agaagttcac agaggtgaag tgattcatgt aggacataca gtgagcaagc atcagggtcc 14940
 ggattatctg actctactct aacttttatg taaatgtgct ttatgccatt aacactgtca 15000
 ttctgtgct tcagctctgg gagactccca agcactctta ggcacaagcc acaattaagg 15060
 gactctgaca ctctgcattg attaattagc atggtggtct ctatgtttcc agattcatga 15120
 ttgtttcact ttccatatag gctatgaagg gtgtgaggaa attttttggg gacagaattg 15180
 gaggcaatcc acctctctca ggaagcctct atctggaaaa gcttacaact cagggacagt 15240
 aactgtaggc ccagtccttg gtgtccaaaa tgggttttat ggtttgaatc tgcaaagcct 15300
 tccatgtgct caaaggtttg aacatggagc ctctctctgg taacactgta ttggaggctt 15360
 ttgagactgg atgctctttg gtcccatgtt ttgctacatc atctgtcaag atatgacca 15420
 ggcagctac cagctaccac agactatgcc tctccagctt tcatgttctc cccaccatga 15480
 tagacttgta tctcctaaaa atggaatcaa agcaaaactt tcctgcatta agttttttt 15540
 tttctgttaa gtgtttggtc acagggacaa gaaaacactc aatacagata attagtacca 15600
 gagttgaggt tcattgctct agcaagttgg atcaaat ttagggctttg gaactgattt 15660
 ataagagaca tgtagaagag tctgaagctg tgggctacag aagtgtcacc agtttttaag 15720
 aatagtttaa tacaccatgg gaattgtgaa aatcagaatg ctcacacaaa ggcagacagg 15780
 aaaacgtgag catgtggcgt gtgagagggc ataagaagga acctaggggg aaatgagcta 15840
 gaagccattc ggctacgtta gggaacgtgt gtggctgtgc ttggcccatg ccctggcaat 15900
 ctgaatgagg ccaaatttta aaggagtggc ctaactcgat tgtcagagaa aatatcaaga 15960
 cagaccacca ctcaggctat gccgtgtttg tgaccgacca gctactctta gccagctcta 16020
 ttgtgaaatt ccagagcaat tatcagagca tgaagatata tacagtttag tgaagtaagg 16080
 ggtgtgggtc cctaagtggg tgggtgcataa atctatgtag gtgatgccta agtgacactt 16140
 gataatccaa aatatcagca atgtggaatg tcttccaagg agacctgtag acacacattt 16200
 tagaactttg ctcatggctg taataaatag ctagctagaa atcatttctt gaagagggtta 16260
 gtctgagtta cggttccagg gcaaaccattc agtgatggca aggaaggcat tgcagtcagg 16320
 agccaaaggc cagctggtca cattgcatca agagtagaga gtcagagtgt gagtagaaag 16380
 aggatacagg ttataaaacc tcactgtcca ctctcagcaa tccattttct cctaaaaggc 16440




```

tttaccttct aaagatttta gtcttcaaaa ccagtaccag tagcctggga acaaaagttg 16500
aaacaaatga gcctttgtgg ggcatttcac acttaaaaca gggcatcacc taggaggagc 16560
cctgtgtgca gtaggaagtg tggcctctgt gtcaggaatg ctcaggctaa taaggggtcc 16620
tctatctgag ggaccctatg aagattcaac aagtagttgt gagaattccc tgtaaatgga 16680
tgctaccaat ttgacatttg tagacctgct attgtgtgct tctttattgg gctctcccat 16740
ctcccaactt tccaacccat attccacatt aatcccttcc accaccatgc aacactaggt 16800
aggagagaag gaaggttaga agagaaagtg ggtatagatc tatttagact acttctgtct 16860
gattaggggc aagtccaatc gtcattgtca ggatacctcc aaccagcaac cagcaaacca 16920
gcaaatcaga aacagcaaaa gcagccaaca aggcagcact aaccagcagg attggggctg 16980
gtagcgtggg agcagtcact actggtcttc tcatggcttt ggcattaata ctctctcaag 17040
aaattccgta attttttccc caccacctga aattccgtaa ttttaaagtc aaactatcta 17100
cagctggcaa aaatcacatc tctcctagag cacaagacaa atcatagtta ctggctattt 17160
gcaatctgaa gcatctcaat atcccacacc tgggattaaa acaaaaacat attcacatca 17220
cataactgtt ttttttttcc aattttttat taggtatttt ctttatttac atttcaaag 17280
ctatcccga agtcccctat accctcccac ctccctgctc ccctacacac ccactcccac 17340
tttttgaccc tggagttccc cgggtactggg gcatataaag tttgcaagac caaggggcct 17400
ctcttcccag tgatggccga ctaagccatc ttctgctaca tatgcagata gagacacgag 17460
ctctgggggt actagttagt tcatattgtt gttccacctc tagggctcga gacccttca 17520
gctccttggg tactttgtct agctcctcca ctgggggctc tgtgttttat ctaatagatg 17580
actgtgagca tccacttctg tatttgacag gcaactggcct agcgtcacat gagccagcta 17640
tatcagggtc ctttcagcaa aaccttgctg gcatgtgcaa tagtgtctgc gtttggtggt 17700
tgattatggg atggatccac tagttctaga gc
17732

```






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/55, 9/16, 1/19, 1/20, 1/21, A01H 5/00, A23K 1/00, 1/165, C12Q 1/44, 1/68, C12N 15/82 // (C12N 1/19, C12R 1:84) (C12N 1/21, C12R 1:01, 1:125, 1:19)		A2	(11) International Publication Number: WO 97/48812
(21) International Application Number: PCT/CA97/00414		(43) International Publication Date: 24 December 1997 (24.12.97)	
(22) International Filing Date: 13 June 1997 (13.06.97)		Suwon, Seoul 442-060 (KR). ZHOU, Luming [CA/US]; 1227 East Brickyard Road #203, Salt Lake City, UT 84106 (US). FORSBERG, Cecil, Wallace [CA/CA]; 44 Hands Drive, Guelph, Ontario N1G 3H3 (CA).	
(30) Priority Data: 60/019,735 14 June 1996 (14.06.96) US 08/744,779 6 November 1996 (06.11.96) US 08/862,531 23 May 1997 (23.05.97) US		(74) Agent: McKAY-CAREY, Mary, Jane; McKay-Carey & Company, 10155 - 102nd Street, 2125 Commerce Place, Edmonton, Alberta T5J 4G8 (CA).	
(71) Applicant (for all designated States except US): HER MAJESTY THE QUEEN IN RIGHT OF CANADA, represented by THE DEPARTMENT OF AGRICULTURE AND AGRI-FOOD CANADA [CA/CA]; Lethbridge Research Station, P.O. Box 3000, Main Lethbridge, Alberta T1J 4B1 (CA).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(72) Inventors; and (75) Inventors/Applicants (for US only): CHENG, Kuo, Joan [CA/CA]; 2015 - 6th Avenue South, Lethbridge, Alberta T1J 1C2 (CA). SELINGER, Leonard, Brent [CA/CA]; 114 Laval Court West, Lethbridge, Alberta T1K 4G3 (CA). YANKE, Lindsey, Jay [CA/CA]; 1520 - 23rd Avenue N. #18, Lethbridge, Alberta T1H 4X9 (CA). BAE, Hee, Dong [KR/KR]; Eung-Sup Shim, 114-5, Chi-dong, Paldal-gu,		Published Without international search report and to be republished upon receipt of that report.	

(54) Title: DNA SEQUENCES ENCODING PHYTASES OF RUMINAL MICROORGANISMS

(57) Abstract

Phytases derived from ruminal microorganisms are provided. The phytases are capable of catalyzing the release of inorganic phosphorus from phytic acid. Preferred sources of phytases include *Selenomonas*, *Prevotella*, *Treponema* and *Megasphaera*. A purified and isolated DNA encoding a phytase of *Selenomonas ruminantium* JY35 (ATCC 55785) is provided. Recombinant expression vectors containing DNAs encoding the phytases and host cells transformed with DNAs encoding the phytases are also provided. The phytases are useful in a wide range of applications involving the dephosphorylation of phytate, including, among other things, use in animal feed supplements.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	IS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DNA SEQUENCES ENCODING PHYTASES OF RUMINAL MICROORGANISMS

Field of the Invention

This invention relates to phytases derived from ruminal microorganisms.

Background of the Invention

Although the plant constituents of livestock feedstuffs are rich in phosphorus, inorganic phosphorus supplementation is required to obtain good growth performance of monogastric animals. Phytic acid (*myo*-inositol hexaphosphoric acid) generally occurs as a complex of calcium, magnesium and potassium salts and/or proteins, and is the predominant form of phosphorus in cereals, oil seeds, and legumes, and accounts for 1 to 3% of the seed dry weight and 60 to 90% of the total phosphorus present in seeds (Graf, 1986). However, monogastric animals (e.g., swine, poultry and fish) utilize phytate poorly or not at all because they are deficient in gastrointestinal tract enzymes capable of hydrolyzing phytate. Phytate passes largely intact through the upper gastrointestinal tract, where it may decrease the bioavailability of nutrients by chelating minerals (e.g., calcium and zinc), binding amino acids and proteins (Graf, 1986) and inhibiting enzymes. Phytate phosphorus in manure poses a serious pollution problem, contributing to eutrophication of surface waters in areas of the world where monogastric livestock production is intensive.

Production inefficiencies and phosphorus pollution caused by phytate may be effectively addressed by phytase supplementation of diets for monogastric animals. Phytases catalyze the hydrolysis of phytate to *myo*-inositol and inorganic phosphate, which are then absorbed in the small intestine. In addition to decreasing phosphorus supplementation requirements and reducing the amount of phytate pollutants released, phytases also diminish the antinutritional effects of phytate.

Phytases are produced in animal and plant (predominantly seeds) tissues and by a variety of microorganisms (U.S. Patent No. 3,297,548; Shieh and Ware, 1968; Ware and Shieh, 1967). Despite the array of potential phytase sources, only soil fungi (*Aspergillus niger* or *Aspergillus ficuum*) are currently used for commercial production of phytase. The phytase produced by *A. ficuum* possesses greater specific activity (100 units/mg of protein (wherein units are defined as μ moles of

phosphate released per minute)) and thermostability compared to those phytases that have been characterized from other microorganisms (European Patent Application No. 0,420,358 (van Gorcum *et al.*, 1991) and U.S. Patent No. 5,436,156 (van Gorcum *et al.*, issued July 25, 1995)). The *A. ficuum* phytase is an acid phytase and exhibits little activity above pH 5.5 (Howson and Davis, 1983; van Gorcum *et al.*, 1991). Consequently, activity is limited to a relatively small region of the monogastric digestive tract, in which the pH ranges from 2-3 (in the stomach) to 4-7 (in the small intestine).

Although the idea of phytase supplementation of monogastric diets was proposed more than 25 years ago (U.S. Patent No. 3,297,548, Ware and Shieh, 1967), the high cost of enzyme production has restricted the use of phytase in the livestock industry. In North America, supplemental phytase is generally more expensive than phosphorus supplements. In some circumstances, the cost of phytase utilization may be partially offset if the use of this enzyme also decreases the need for supplementation of a second nutrient such as calcium. The use of phytase in North America is likely to increase as swine and poultry populations increase and as public pressures force a reduction in pollution associated with livestock production. Higher costs of phosphorus supplements and legislation requiring the use of phytase have made the use of this supplement more common in Europe and parts of the Orient than in North America. Governments of the Netherlands, Germany, Korea and Taiwan have enacted or are enacting legislation to reduce the phosphorus pollution created by monogastric livestock production.

A more effective means of increasing phytase utilization is through cost reduction. The cost of phytase can be reduced by decreasing production costs and/or producing an enzyme with superior activity. Recent advances in biotechnology may revolutionize the commercial enzyme industry by offering alternative, cost effective methods of enzyme production. Application of recombinant DNA technology has enabled manufacturers to increase the yields and efficiency of enzyme production, and to create new products. The original source organism need no longer limit the production of commercial enzymes. Genes encoding superior enzymes can be transferred from organisms such as anaerobic bacteria and fungi, typically impractical for commercial production, into well characterized industrial

1 microbial production hosts (e.g., *Aspergillus* and *Bacillus* spp.). As well, these genes
2 may be transferred to novel plant and animal expression systems.

3 Unlike monogastric animals, ruminants (e.g., cattle, sheep) readily utilize the
4 phosphorus in phytic acid. It has been demonstrated that phytases are present in
5 the rumen, and it has been proposed that ruminants reared on high grain diets (rich
6 in phytate) do not require dietary phosphorus supplementation due to these ruminal
7 phytases. A single report has attributed this phytase production to ruminal
8 microorganisms (Raun *et al.*, 1956), but overall, the unique capacity of ruminants to
9 utilize phytate has largely been ignored. Raun *et al.* (1956) prepared microbial
10 suspensions by centrifugal sedimentation (Cheng *et al.*, 1955). Those microbial
11 suspensions were almost certainly contaminated with microscopic particles of plant
12 material. Since plants produce phytases, the study was inconclusive as to whether
13 plant phytases or microbial phytases produced the observed activity. Although Raun
14 *et al.* have raised the possibility that ruminal phytase production may be attributable
15 to ruminal microorganisms, this possibility has not been explored.

16 In view of the foregoing, there remains a need for low cost phytases having
17 biochemical characteristics well suited for use in animal feed supplements.

18 19 **Summary of the Invention**

20 The inventors have discovered that the rumen is a rich source of
21 microorganisms which produce phytases having biochemical characteristics (such
22 as temperature and pH stability, low metal ion sensitivity and high specific activity)
23 desirable for industrial applications such as animal feed supplementation and inositol
24 production. Ruminal microorganisms tolerate anaerobic conditions and may be
25 either facultative or obligate anaerobes. Ruminal microorganisms may be
26 prokaryotes (i.e. bacteria) or eukaryotes (i.e. fungi, protozoa). As used herein, the
27 term "ruminal microorganisms" includes microorganisms isolated from the digesta or
28 feces of a ruminant animal.

29 Ruminal bacterial species which have been identified as providing particularly
30 active phytases includes *Selenomonas ruminantium*, *Prevotella* sp, *Treponema*
31 *bryantii* and *Megaphaera elsdenii*. *Prevotella* and *Selenomonas* are Gram negative
32 anaerobic rods from the family Bacteroidaceae.

1 In accordance with the present invention, DNA sequences encoding novel and
2 useful phytases derived from ruminal microorganisms are provided.

3 A phytase gene (*phyA*) from *Selenomonas ruminantium* strain JY35 has been
4 cloned and sequenced, and the nucleotide sequence of the *phyA* gene is provided.
5 The invention extends to DNA sequences which encode phytases and which are
6 capable of hybridizing under stringent conditions with the *phyA* gene sequence. As
7 used herein, "capable of hybridizing under stringent conditions" means annealing to
8 a subject nucleotide sequence, or its complementary strand, under standard
9 conditions (ie. high temperature and/or low salt content) which tend to disfavor
10 annealing of unrelated sequences. As used herein, "conditions of low stringency"
11 means hybridization and wash conditions of 40 - 50°C, 6 X SSC and 0.1% SDS
12 (indicating about 50 - 80% homology). As used herein, "conditions of medium
13 stringency" means hybridization and wash conditions of 50 - 65°C, 1 X SSC and
14 0.1% SDS (indicating about 80 - 95% homology). As used herein, "conditions of high
15 stringency" means hybridization and wash conditions of 65 - 68°C, 0.1 X SSC and
16 0.1% SDS (indicating about 95-100% homology).

17 As used herein, the term "phytase" means an enzyme capable of catalyzing
18 the removal of inorganic phosphorus from a *myo*-inositol phosphate.

19 As used herein, the term "*myo*-inositol phosphate" includes, without limitation,
20 *myo*-inositol hexaphosphate, *myo*-inositol pentaphosphate, *myo*-inositol
21 tetraphosphate, *myo*-inositol triphosphate, *myo*-inositol diphosphate and *myo*-inositol
22 monophosphate.

23 As used herein, "phytate" means the salt of *myo*-inositol hexaphosphoric acid.

24 The invention extends to the *S. ruminantium* JY35 (ATCC 55785) organism
25 itself, and to methods for identifying and isolating this and other ruminal
26 microorganisms exhibiting phytase activity as well as methods for isolating, cloning
27 and expressing phytase genes from ruminal microorganisms exhibiting phytase
28 activity using part or all of the *phyA* gene sequence as a probe.

29 The invention further extends to methods for assaying phytase production by
30 a microorganism whereby false positive results caused by microbial acid production
31 are eliminated. Colonies of microorganisms are grown on a growth medium
32 containing phytate. The medium is contacted with an aqueous solution of cobalt

1 chloride and the medium is then examined for zones of clearing. Preferably, rather
2 than examining the medium immediately, the solution of cobalt chloride is removed
3 and the medium is contacted with aqueous solutions of ammonium molybdate and
4 ammonium vanadate and then examined for zones of clearing. False positive results
5 which occur when acid-forming microbes produce zones of clearing are avoided.

6 The invention extends to expression constructs constituting a DNA encoding
7 a phytase of the present invention operably linked to control sequences capable of
8 directing expression of the phytase in a suitable host cell.

9 The invention further extends to host cells which have been transformed with,
10 and express, DNA encoding a phytase of the present invention, and to methods of
11 producing such transformed host cells. As used herein "host cell" includes animal,
12 plant, yeast, fungal, protozoan and prokaryotic host cells.

13 The invention further extends to transgenic plants which have been
14 transformed with a DNA encoding a phytase of the present invention so that the
15 transformed plant is capable of expressing the phytase and to methods of producing
16 such transformed plants. As used herein, "transgenic plant" includes transgenic
17 plants, tissues and cells.

18 Phytases of the present invention are useful in a wide variety of applications
19 involving the dephosphorylation of phytate. Such applications include use in animal
20 feed supplements, feedstuff conditioning, human nutrition, and the production of
21 inositol from phytic acid. Phytases of the present invention may also be used to
22 minimize the adverse effects of phytate metal chelation. The high phytate content
23 of certain feedstuffs such as soy meal decreases their value as protein sources for
24 fish, monogastric animals, young ruminants and infants because the phytate
25 decreases the bioavailability of nutrients by chelating minerals, and binding amino
26 acids and proteins. Treatment of such feedstuffs with the phytases of the present
27 invention will reduce their phytate content by phytase mediated dephosphorylation,
28 rendering the feedstuffs more suitable for use as protein sources. Accordingly, the
29 invention extends to novel feed compositions comprising feedstuffs treated with a
30 phytase of the present invention, and feed additives containing a phytase of the
31 present invention. Such feed compositions and additives may also contain other
32 enzymes, such as, proteases, cellulase, xylanases and acid phosphatases. The

1 phytase may be added directly to an untreated, pelletized, or otherwise processed
2 feedstuff, or it may be provided separately from the feedstuff in, for instance, a
3 mineral block, a pill, a gel formulation, a liquid formulation, or in drinking water. The
4 invention extends to feed inoculant preparations comprising lyophilized
5 microorganisms which express phytases of the present invention under normal
6 growing conditions. With respect to these feed inoculant preparations, "normal
7 growing conditions" mean culture conditions prior to harvesting and lyophilization of
8 the microorganisms. The microorganisms express phytases during growth of the
9 microbial cultures in large-scale fermenters. The activity of phytases in the
10 microorganisms is preserved by lyophilization of the harvested microbial
11 concentrates containing the phytase.

12 The invention further extends to a method for improving an animal's utilization
13 of dietary phosphate by feeding the animal an effective amount of a phytase of the
14 present invention. As used herein "an effective amount" of a phytase means an
15 amount which results in a statistically significant improvement in phosphorus
16 utilization by the animal. Phytate phosphorus utilization may be evidenced by, for
17 instance, improved animal growth and reduced levels of phytate in animal manure.

18 19 **Brief Description of Drawings**

20 Figure 1 is a photograph showing the effect of counterstaining agar medium
21 containing phytate on zones of clearing produced by acid production or phytase
22 activity. Phytate agar was inoculated with *S. bovis* (top of left petri dish) and *S.*
23 *ruminantium* JY35 (bottom of left petri dish) and incubated for 5 d at 37°C. The
24 colonies were scraped off and the medium counterstained with cobalt chloride and
25 ammonium molybdate/ammonium vanadate solutions (right petri plate).

26 Figure 2 is a graph illustrating the growth (protein) and phytase production of
27 *S. ruminantium* JY35 in modified Scott and Dehority (1965) broth.

28 Figure 3A, 3B and C show transmission electron micrographs of cells from a
29 mid-exponential phase culture of *S. ruminantium* JY35 incubated for reaction product
30 deposition by phytase using sodium phytate as the substrate. Untreated control cells
31 are shown for comparison in Figures 3D, 3E and 3F.

Figure 4 is a graph illustrating the phytase pH profile for washed *S. ruminantium* JY35 cells in five different buffers.

Figure 5 is a graph illustrating the pH profile of *S. ruminantium* JY35 $MgCl_2$ cell extract in five different buffers.

Figure 6 is a graph illustrating the temperature profile of *S. ruminantium* JY35 $MgCl_2$ cell extract.

Figure 7 is a graph illustrating the effect of ions (10 mM) on *S. ruminantium* JY35 phytase activity (Ctr = control).

Figure 8 is a graph illustrating the effect of sodium phytate concentration on *S. ruminantium* JY35 phytase activity.

Figure 9 is a zymogram developed for confirmation of phytase activity. Concentrates (10 x) of *S. ruminantium* JY35 $MgCl_2$ extract (lanes B - E), low molecular weight markers (lane F, BioRad Laboratories Canada Ltd, Mississauga, Ontario) and *A. ficuum* phytase (Sigma, 1.6 U, lane A) were resolved by SDS-PAGE in a 10% polyacrylamide gel. Lanes A to E were stained for phytase activity and Lane F was stained with Coomassie brilliant blue.

Figure 10 is a photograph of a phytate hydrolysis plate assay for phytase activities of *E. coli* DH5 α transformed with pSrP.2 (top), pSrP.2 $\Delta SphI$ (bottom left), and pSrPf6 (bottom right). Zones of clearing were visible after incubating the plates at 37°C for 48 h.

Figure 11 is a Southern blot analysis using the 2.7-kb fragment from pSrP.2 as a probe against *SphI* digested pSrP.2 DNA (lane B) and *HindIII* digested genomic DNA isolated from *S. ruminantium* JY35 (lane C). Digoxigenin labelled *HindIII* digested Lambda DNA was run as a molecular weight standard in lane A.

Figure 12 is a physical map of pSrP.2. A 2.7-kb fragment, from a *Sau3A* partial digest of *S. ruminantium* JY35 genomic DNA, was cloned into the *BamHI* site of pUC18. This fragment contains the entire gene encoding the phytase from *S. ruminantium* JY35. The location of a *BamHI* site lost as a result of the ligation is indicated in square brackets.

Figure 13 is a schematic representation of the deletion analysis of the *S. ruminantium* phytase gene. The position of *phyA* is indicated by the horizontal arrow.

The hatched boxes indicate segments of the 2.7-kb *Sau*3A fragment carried by different plasmid derivatives. Phytase activity is indicated in the panel to the right.

Figure 14 is a zymogram developed for phytase activity. *E. coli* DH5 α (pSrP.2) cells (lane A), *E. coli* DH5 α (pSrP.2 Δ *SphI*) cells (lane B), and low molecular weight markers (lane C, BioRad Laboratories) were resolved by SDS-PAGE in a 10% polyacrylamide gel. Lanes A and B were stained for phytase activity and Lane C was stained with Coomassie brilliant blue.

Figure 15 is the nucleotide sequence of the *S. ruminantium* JY35 phytase gene (*phyA*) (SEQ ID NO. 1) and its deduced amino acid sequence (SEQ ID NO. 2). Nucleotide 1 corresponds to nt 1232 of the 2.7-kb insert of pSrP.2. The putative ribosome binding site is underlined and shown above the sequence as R.B.S. The signal peptidase cleavage site, predicted by the method of von Heijne (1986) is indicated by the \downarrow . The N-terminal amino acid sequence of the phytase secreted by *E. coli* (pSrPf6) is underlined.

Detailed Description of the Preferred Embodiment

The rumen is a complex ecosystem inhabited by more than 300 species of bacteria, fungi and protozoa. Screening these organisms for phytase activity requires the ability to discriminate the phytase activity of individual isolates. This may be accomplished through the assessment of pure cultures from a stock culture collection or separation and cultivation of individual cells through cultural techniques (e.g., streak plate, dilution and micromanipulation). Standard aseptic, anaerobic techniques described for bacteria, fungi and protozoa may be used to accomplish this goal.

Suitable enzyme assays are necessary for screening microbial isolates in ruminal fluid samples and from culture collections, and for cloning phytase genes. Assays for measuring phytase activity in solutions have been described in the literature. Sample solutions are typically assayed for phytase activity by measuring the release of inorganic phosphorus (P_i) from phytic acid (Raun *et al.*, 1956; van Hartingsveldt *et al.*, 1993). Phytase activity may also be detected on solid media. Microorganisms expressing phytase produce zones of clearing on agar media containing sodium or calcium phytate (Shieh and Ware, 1968; Howson and Davis,

1 1983). However, the solid media assays described in the literature were found to be
2 unsatisfactory for screening ruminal bacteria for phytase activity because of the false
3 positive reactions of acid-producing bacteria such as *Streptococcus bovis*. To
4 overcome this problem, a two-step counterstaining procedure was developed in
5 which petri dishes containing solid medium are flooded first with an aqueous cobalt
6 chloride solution and second with an aqueous ammonium molybdate/ammonium
7 vanadate solution. Following this treatment only clearing zones produced by enzyme
8 activity are evident (Figure 1).

9 Using the above solutions and solid medium assays, 345 isolates from the
10 Lethbridge Research Centre (Lethbridge, Alberta, Canada) culture collection were
11 screened for phytase activity (Table 1). A total of 29 cultures with substantial
12 phytase activity were identified, including 24 of the genus *Selenomonas* and 5 of the
13 genus *Prevotella*. Twelve of these cultures (11 *Selenomonas* isolates and 1
14 *Prevotella* isolate) had phytase activities substantially higher than the other positive
15 cultures (Table 2).

16 The phytase of *S. ruminantium* JY35 (deposited May 24, 1996 with the
17 American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland,
18 20852-1776, as ATCC 55785) was selected for further examination and compared
19 to a commercial phytase (Gist-brocades nv, Delft, The Netherlands) from *Aspergillus*
20 *ficuum* NRRL 3135 (van Gorcum *et al.*, 1991 and 1995). The phytase of *S.*
21 *ruminantium* JY35 (ATCC 55785) is constitutively expressed, exported from the cell
22 and associated with the cell surface. The pH (Figure 5) and temperature (Figure 6)
23 profiles of the *S. ruminantium* JY35 (ATCC 55785) phytase were comparable, if not
24 more suited to industrial production, than are those of the commercial *A. ficuum*
25 NRRL 3135 phytase. These results demonstrated the potential of ruminal and
26 anaerobic microbes as sources of phytases with characteristics superior to phytases
27 currently being produced by industry.

28 Microbial genes encoding selected enzymes can be cloned by a variety of
29 methods. Gene libraries (genomic DNA and/or cDNA) are constructed by standard
30 methods (Sambrook *et al.*, 1989; Ausubel *et al.*, 1990) and screened for the desired
31 gene. The screening methodology may utilize heterologous probes, enzyme activity

1 or results generated during purification of the gene product, such as N-terminal and
2 internal amino acid sequence data and antibodies.

3 Using the solid medium phytase assay developed to detect phytase activity
4 produced by ruminal microbes, a *S. ruminantium* JY35 (ATCC 55785) gene library
5 was screened for positive clones. Of 6000 colonies examined, a single colony was
6 identified as a phytase positive clone by a large zone of clearing around the colony.
7 This clone carried a 5.5-kb plasmid comprising a 2.7-kb *Sau*3A DNA fragment
8 inserted into cloning vector pUC18. The newly isolated 2.7-kb *Sau*3A DNA fragment
9 was used as a probe in Southern blot hybridizations. Under high stringency
10 conditions, a discrete band could be detected for *S. ruminantium* isolate JY35 (ATCC
11 55785), but not for *Prevotella* sp. 46/5², *E. coli* DH5 α or *A. ficuum* NRRL 3135.

12 Plasmid DNA isolated from the newly isolated clone and introduced into *E. coli*
13 cells by transformation produced ampicillin-resistant, phytase-positive CFUs.
14 Zymogram analysis of cell extracts from *E. coli* DH5 α cells carrying the 2.7-kb *Sau*3A
15 DNA fragment from *S. ruminantium* JY35 (ATCC 55785) revealed a single activity
16 band with an estimated molecular mass of 37 kDa. Deletion and DNA sequence
17 analyses were used to identify the gene (*phyA*) which encoded the phytase
18 responsible for the activity observed in recombinant *E. coli* clones. The N-terminal
19 amino acid sequence of the purified 37-kDa phytase expressed in *E. coli* cells
20 carrying *phyA* matched the N-terminal amino acid sequence of the mature phytase
21 predicted from the cloned *phyA* sequence. This indicated conclusively that the
22 nucleotide sequence encoding the phytase had been isolated. The nucleotide
23 sequence and deduced amino acid sequence are shown in Figure 15.

24 As with other genes, it is possible to use the characterized phytase coding
25 sequence in a variety of expression systems for commercial enzyme production.
26 Application of recombinant DNA technology has enabled enzyme manufacturers to
27 increase the volume and efficiency of enzyme production, and to create new
28 products. The original source organism need no longer limit the production of
29 commercial enzymes. Genes encoding superior enzymes can be transferred from
30 organisms such as anaerobic bacteria and fungi, typically impractical for commercial
31 production, into well characterized industrial microbial production hosts (e.g.,

1 *Aspergillus*, *Pichia*, *Trichoderma*, *Bacillus* spp.). As well, these genes may be
2 transferred to novel plant and animal expression systems.

3 Industrial strains of microorganisms (e.g., *Aspergillus niger*, *Aspergillus*
4 *ficuum*, *Aspergillus awamori*, *Aspergillus oryzae*, *Trichoderma reesei*, *Mucor miehei*,
5 *Kluyveromyces lactis*, *Pichia pastoris*, *Saccharomyces cerevisiae*, *Escherichia coli*,
6 *Bacillus subtilis* or *Bacillus licheniformis*) or plant hosts (e.g., canola, soybean, corn,
7 potato) may be used to produce phytase. All systems employ a similar approach to
8 gene expression. An expression construct is assembled to include the protein
9 coding sequence of interest and control sequences such as promoters, enhancers
10 and terminators. Other sequences such as signal sequences and selectable
11 markers may also be included. To achieve extracellular expression of phytase, the
12 expression construct of the present invention utilizes a secretory signal sequence.
13 The signal sequence is not included on the expression construct if cytoplasmic
14 expression is desired. The promoter and signal sequence are functional in the host
15 cell and provide for expression and secretion of the coding sequence product.
16 Transcriptional terminators are included to ensure efficient transcription. Ancillary
17 sequences enhancing expression or protein purification may also be included in the
18 expression construct.

19 The protein coding sequences for phytase activity are obtained from ruminal
20 microbial sources. This DNA may be homologous or heterologous to the expression
21 host. Homologous DNA is herein defined as DNA originating from the same species.
22 For example, *S. ruminantium* may be transformed with DNA from *S. ruminantium* to
23 improve existing properties without introducing properties that did not exist previously
24 in the species. Heterologous DNA is defined as DNA originating from a different
25 species. For example, the *S. ruminantium phyA* may be cloned and expressed in *E.*
26 *coli*.

27 It is well known in the biological arts that certain amino acid substitutions can
28 be made in protein sequences without affecting the function of the protein.
29 Generally, conservative amino acid substitutions are tolerated without affecting
30 protein function. Similar amino acids can be those that are similar in size and/or
31 charge properties, for example, aspartate and glutamate and isoleucine and valine
32 are both pairs of similar amino acids. Similarity between amino acid pairs has been

1 assessed in the art in a number of ways. For example, Dayhoff et al. (1978) in Atlas
2 of Protein Sequence and Structure, Volume 5, Supplement 3, Chapter 22, pages
3 345-352, which is incorporated by reference herein, provides frequency tables for
4 amino acid substitutions which can be employed as a measure of amino acid
5 similarity. Dayoff et al.'s frequency tables are based on comparisons of amino acid
6 sequences for proteins having the same function from a variety of evolutionary
7 different sources.

8 It is also well-known that often less than a full length protein has the function
9 of the complete protein, for example, a truncated protein lacking an N-terminal,
10 internal or a C-terminal protein often ha the biological and/or enzymatic activity of the
11 complete natural protein. Gene truncation experiments involving *phyA* have
12 confirmed that the truncated protein may retain the function of the intact protein.
13 *Exherichia coli* clones expressing PhyA missing N-terminal amino acids 1-37 or 1058
14 (SEQ ID NO. 2) showed phytase positive phenotypes. In contrast, no phytase
15 activity could be detected for a clone expressing PhyA missing acids 307-346 (SEQ
16 ID NO. 2). Those of ordinary skill in the art know how to make truncated protein and
17 proteins with internal deletions. In the present invention, the function of a truncated
18 phytase protein or an internally deleted phytase protein can be readily tested using
19 the assay described hereinbelow and n view of what is generally known in the art.

20 Substituted, internally-deleted and truncated rumina phytase derivatives which
21 retain substantially the same enzymatic activity as a phytase specifically disclosed
22 herein are considered equivalents of the exemplified phytase and are within the
23 scope of the present invention, particularly where the specific activity of the
24 substituted, internally-deleted or truncated phytase derivative is at least about 10%
25 of the specifically exemplified phytase. The skilled artisan can readily measure the
26 activity of a rumina phytase, truncated phytase, internally-deleted phytase or
27 substituted phytase using the assay procedures taught herein and in view of what
28 is generally known in the art.

29 This invention includes structurally variant phytases derived from a phytase
30 of a rumina microorganisms, particularly those derived from a phytase specifically
31 disclosed herein, that are substantially functionally equivalent to that phytase as
32 assayed as described herein in view of what is generally known in the art.

1 Structurally variant, functional equivalents of the phytases of this invention include
2 those phytase of rumina microorganisms having a contiguous amino acid sequence
3 as in the phytase amino acid sequence disclosed herein (SEQ ID NO. 2), particularly
4 those variant phytase which have a contiguous amino acid sequence of a phytase
5 of a rumina microorganism that is a contiguous sequence at least about 25 amino
6 acids in length.

7 The present invention also provides the starting material for the construction
8 of phytases with properties that differ from those of the enzymes isolated herein.
9 The genes can be readily mutated by known procedures (e.g., chemical, site
10 directed, random polymerase chain reaction mutagenesis) thereby creating gene
11 products with altered properties (e.g., temperature or pH optima, specific activity or
12 substrate specificity).

13 Various promoters (transcriptional initiation regulatory region) may be used
14 according to the present invention. The selection of the appropriate promoter is
15 dependent upon the proposed expression host. Choices of promoters may include
16 the promoter associated with the cloned protein coding sequence or promoters from
17 heterologous sources as long as they are functional in the chosen host. Examples
18 of heterologous promoters are the *E. coli* *tac* and *trc* promoters (Brosius *et al.*, 1985),
19 *Bacillus subtilis* *sacB* promoter and signal sequence (Wong, 1989), *aox1* and *aox2*
20 from *Pichia pastoris* (Ellis *et al.*, 1985), and oleosin seed specific promoter from
21 *Brassica napus* or *Arabidopsis thaliana* (van Rooijen and Moloney, 1994). Promoter
22 selection is also dependent upon the desired efficiency and level of peptide or
23 protein production. Inducible promoters such *tac* and *aox1* are often employed in
24 order to dramatically increase the level of protein expression. Overexpression of
25 proteins may be harmful to the host cells. Consequently, host cell growth may be
26 limited. The use of inducible promoter systems allows the host cells to be cultivated
27 to acceptable densities prior to induction of gene expression, thereby facilitating
28 higher product yields. If the protein coding sequence is to be integrated through a
29 gene replacement (omega insertion) event into a target locus, then promoter
30 selection may also be influenced by the degree of homology to the target locus
31 promoter.

1 Various signal sequences may be used according to the present invention.
2 A signal sequence which is homologous to the protein coding sequence to be
3 expressed may be used. Alternatively, a signal sequence which has been selected
4 or designed for improved secretion in the expression host may also be used. For
5 example, *B. subtilis* *sacB* signal sequence for secretion in *B. subtilis*, the
6 *Saccharomyces cerevisiae* α -mating factor or *P. pastoris* acid phosphatase *phoA*
7 signal sequences for *P. pastoris* secretion may be used. A signal sequence with a
8 high degree of homology to the target locus may be required if the protein coding
9 sequence is to be integrated through an omega insertion event. The signal
10 sequence may be joined directly through the sequence encoding the signal
11 peptidase cleavage site to the protein coding sequence, or through a short
12 nucleotide bridge consisting of usually fewer than ten codons.

13 Elements for enhancing expression transcription (promoter activity) and
14 translation have been identified for eukaryotic protein expression systems. For
15 example, positioning the cauliflower mosaic virus (CaMV) promoter 1000 bp on either
16 side of a heterologous promoter may elevate transcriptional levels by 10- to 400-fold.
17 The expression construct should also include the appropriate translational initiation
18 sequences. Modification of the expression construct to include the Kozak consensus
19 sequence for proper translational initiation may increase the level of translation by
20 10 fold.

21 Elements to enhance purification of the protein may also be included in the
22 expression construct. The product of oleosin gene fusions is a hybrid protein
23 containing the oleosin gene joined to the gene product of interest. The fusion protein
24 retains the lipophilic properties of oleosins and is incorporated in the oil body
25 membranes (van Rooijen and Moloney, 1994). Association with the oil bodies may
26 be exploited to facilitate purification of the recombinant oleosin fusion proteins (van
27 Rooijen and Moloney, 1994).

28 A selection marker is usually employed, which may be part of the expression
29 construct or separate from it (e.g., carried by the expression vector), so that the
30 marker may integrate at a site different from the gene of interest. Transformation of
31 the host cells with the recombinant DNA molecules of the invention is monitored
32 through the use of selectable markers. Examples of these are markers that confer

1 resistance to antibiotics (e.g., *bla* confers resistance to ampicillin for *E. coli* host cells,
2 *nptII* confers kanamycin resistance to *B. napus* cells) or that permit the host to grow
3 on minimal medium (e.g., *HIS4* enables *P. pastoris* GS115 His⁻ to grow in the
4 absence of histidine). The selectable marker will have its own transcriptional and
5 translational initiation and termination regulatory regions to allow for independent
6 expression of the marker. Where antibiotic resistance is employed as a marker, the
7 concentration of the antibiotic for selection will vary depending upon the antibiotic,
8 generally ranging from 10 to 600 µg of the antibiotic/mL of medium.

9 The expression construct is assembled by employing known recombinant
10 DNA techniques. Restriction enzyme digestion and ligation are the basic steps
11 employed to join two fragments of DNA. The ends of the DNA fragment may require
12 modification prior to ligation and this may be accomplished by filling in overhangs,
13 deleting terminal portions of the fragment(s) with nucleases (e.g., *ExoIII*), site
14 directed mutagenesis, and adding new base pairs by the polymerase chain reaction
15 (PCR). Polylinkers and adaptors may be employed to facilitate joining of select
16 fragments. The expression construct is typically assembled in stages employing
17 rounds of restriction, ligation and transformation of *E. coli*. There are numerous
18 cloning vectors available for construction of the expression construct and, the
19 particular choice is not critical to this invention. The selection of cloning vector will
20 be influenced by the gene transfer system selected for introduction of the expression
21 construct into the host cell. At the end of each stage, the resulting construct may be
22 analyzed by restriction, DNA sequence, hybridization and PCR analyses.

23 The expression construct may be transformed into the host as the cloning
24 vector construct, either linear or circular, or may be removed from the cloning vector
25 and used as is or introduced onto a delivery vector. The delivery vector facilitates
26 the introduction and maintenance of the expression construct in the selected host
27 cell type. The expression construct is introduced into the host cells by employing any
28 of a number of gene transfer systems (e.g., natural competence, chemically
29 mediated transformation, protoplast transformation, electroporation, biolistic
30 transformation, transfection, or conjugation). The gene transfer system selected
31 depends upon the host cells and vector systems used.

1 For instance, the expression construct can be introduced into *P. pastoris* cells
2 by protoplast transformation or electroporation. Electroporation of *P. pastoris* is
3 easily accomplished and yields transformation efficiencies comparable to spheroplast
4 transformation. *P. pastoris* cells are washed with sterile water and resuspended in
5 a low conductivity solution (e.g., 1 M sorbitol solution). A high voltage shock applied
6 to the cell suspension creates transient pores in the cell membrane through which
7 the transforming DNA (e.g., expression construct) enters the cells. The expression
8 construct is stably maintained by integration, through homologous recombination,
9 into the *aox1* (alcohol oxidase) locus.

10 Alternatively, an expression construct, comprising the *sacB* promoter and
11 signal sequence operably linked to the protein coding sequence, is carried on
12 pUB110, a plasmid capable of autonomously replicating in *B. subtilis* cells. The
13 resulting plasmid construct is introduced into *B. subtilis* cells by transformation.
14 *Bacillus subtilis* cells develop natural competence when grown under nutrient poor
15 conditions.

16 In a third example, *Brassica napus* cells are transformed by *Agrobacterium*-
17 mediated transformation. The expression construct is inserted onto a binary vector
18 capable of replication in *A. tumefaciens* and mobilization into plant cells. The
19 resulting construct is transformed into *A. tumefaciens* cells carrying an attenuated Ti
20 or "helper plasmid". When leaf disks are infected with the recombinant *A.*
21 *tumefaciens* cells, the expression construct is transferred into *B. napus* leaf cells by
22 conjugal mobilization of the binary vector::expression construct. The expression
23 construct integrates at random into the plant cell genome.

24 Host cells carrying the expression construct (i.e., transformed cells) are
25 identified through the use of the selectable marker carried by the expression
26 construct or vector and the presence of the gene of interest confirmed by a variety
27 of techniques including hybridization, PCR, and antibodies.

28 The transformant microbial cells may be grown by a variety of techniques
29 including batch and continuous fermentation on liquid or semi-solid media.
30 Transformed cells are propagated under conditions optimized for maximal product-
31 to-cost ratios. Product yields may be dramatically increased by manipulating of
32 cultivation parameters such as temperature, pH, aeration, and media composition.

Careful manipulation and monitoring of the growth conditions for recombinant hyper-expressing *E. coli* cells may result in culture biomass and protein yields of 150 g (wet weight) of cells/L and 5 g of insoluble protein/L, respectively. Low concentrations of a protease inhibitor (e.g., phenylmethylsulfonyl fluoride or pepstatin) may be employed to reduce proteolysis of the over-expressed peptide or protein. Alternatively, protease deficient host cells may be employed to reduce or eliminate degradation of the desired protein.

After selection and screening, transformed plant cells can be regenerated into whole plants and varietal lines of transgenic plants developed and cultivated using known methods. As used herein, "transgenic plant" includes transgenic plants, plant tissues and plant cells.

Following fermentation, the microbial cells may be removed from the medium through down-stream processes such as centrifugation and filtration. If the desired product is secreted, it can be extracted from the nutrient medium. In the case of intracellular production, the cells are harvested and the product released by rupturing cells through the application of mechanical forces, ultrasound, enzymes, chemicals and/or high pressure. Production of an insoluble product, such as occurs in hyper-expressing *E. coli* systems, can be used to facilitate product purification. The product inclusions can be extracted from disrupted cells by centrifugation and contaminating proteins may be removed by washing with a buffer containing low concentrations of a denaturant (e.g., 0.5 to 6 M urea, 0.1 to 1% sodium dodecyl sulfate or 0.5 to 4.0 M guanidine-HCl). The washed inclusions may be solubilized in solutions containing 6 to 8 M urea, 1 to 2% sodium dodecyl sulfate or 4 to 6 M guanidine-HCl. Solubilized product can be renatured by slowly removing denaturing agents during dialysis.

Phytase may be extracted from harvested portions or whole plants by grinding, homogenization, and/or chemical treatment. The use of seed specific lipophilic oleosin fusions can facilitate purification by partitioning the oleosin fusion protein in the oil fraction of crushed canola seeds, away from the aqueous proteins (van Rooijen and Moloney, 1994).

If necessary, various methods for purifying the product, from microbial, fermentation and plant extracts, may be employed. These include precipitation (e.g.,

1 ammonium sulfate precipitation), chromatography (gel filtration, ion exchange, affinity
2 liquid chromatography), ultrafiltration, electrophoresis, solvent-solvent extraction
3 (e.g., acetone precipitation), combinations thereof, or the like.

4 All or a portion of the microbial cultures and plants may be used directly in
5 applications requiring the action of phytase. Various formulations of the crude or
6 purified phytase preparations may also be prepared. The enzymes can be stabilized
7 through the addition of other proteins (e.g., gelatin, skim milk powder) and chemical
8 agents (e.g., glycerol, polyethylene glycol, reducing agents and aldehydes). Enzyme
9 suspensions can be concentrated (e.g., tangential flow filtration) or dried (spray and
10 drum drying, lyophilization) and formulated as liquids, powders, granules, pills,
11 mineral blocks and gels through known processes. Gelling agents such as gelatin,
12 alginate, collagen, agar, pectin and carrageenan may be used.

13 Further, complete dephosphorylation of phytate may not be achieved by
14 phytase alone. Phytases may not dephosphorylate the lower *myo*-inositol
15 phosphates. For instance, an *A. ficuum* phytase described in U.S. Patent No.
16 5,536,156 (van Gorcum *et. al.*, issued July 25, 1995) exhibits low or no phosphatase
17 activity against *myo*-inositol di-phosphate or *myo*-inositol mono-phosphate. Addition
18 of another phosphatase, such as an acid phosphatase, to a feed additive of the
19 present invention containing phytase will help dephosphorylate *myo*-inositol
20 di-phosphate and *myo*-inositol mono-phosphate.

21 Formulations of the desired product may be used directly in applications
22 requiring the action of a phytase. Liquid concentrates, powders and granules may
23 be added directly to reaction mixtures, fermentations, steeping grains, and milling
24 waste. The formulated phytase can be administered to animals in drinking water, in
25 a mineral block, as a salt, or as a powdered supplement to be sprinkled into feed
26 bunks or mixed with a ration. It may also be mixed with, sprayed on or pelleted with
27 other feed stuffs through known processes. Alternatively, a phytase gene with a
28 suitable promoter-enhancer sequence may be intergrated into an animal genome
29 and selectively expressed in an organ or tissue (e.g. salivary glands, pancreas or
30 epithelial cells) which secrete the phytase enzyme into the gastrointestinal tract,
31 thereby eliminating the need for the addition of supplemental phytase.

1 In a preferred formulation, phytases of the present invention may take the
2 form of microbial feed inoculants. Cultures of microorganisms expressing a native
3 phytase, such as *S. ruminantium* JY35 (ATCC 55785), or recombinant
4 microorganisms expressing a phytase encoded by a heterologous phytase gene are
5 grown to high concentrations in fermenters and then harvested and concentrated by
6 centrifugation. Food-grade whey and/or other cryoprotective agents are then
7 admixed with the cell concentrate. The resulting mixture is then cryogenically frozen
8 and freeze-dried to preserve phytase activity by standard lyophilization procedures.
9 The freeze-dried culture may be further processed to form a finished product by such
10 further steps as blending the culture with an inert carrier to adjust the strength of the
11 product.

12 All or a portion of the microbial cultures and plants as produced by the present
13 invention may be used in a variety of industrial processes requiring the action of a
14 phytase. Such applications include, without limitation, the manufacture of end
15 products such as inositol phosphate and inositol, production of feed ingredients and
16 feed additives for non-ruminants (e.g., swine, poultry, fish, pet food), in human
17 nutrition, and in other industries (soybean and corn processing, starch, and
18 fermentation) that involve feedstocks containing phytate. Degradation of phytate
19 makes inorganic phosphate and chelated metals available to animals and
20 microorganisms. The action of phytase increases the quality, value and utility of feed
21 ingredients and/or fermentation substrates that are high in phytate. The action of
22 phytases can also accelerate the steeping process and separation processes
23 involved in the wet milling of corn.

24 The phytase genes of the present invention can be used in heterologous
25 hybridization and polymerase chain reaction experiments, directed to isolation of
26 phytase encoding genes from other microorganisms. The examples herein are given
27 by way of illustration and are in no way intended to limit the scope of the present
28 invention. Efforts have been made to ensure the accuracy with respect to numbers
29 used (e.g., temperature, pH, amounts) but the possibility of some experimental
30 variance and deviations should be recognized.

Example 1

Isolation of ruminal bacteria

Ruminal fluid from a cannulated Holstein cow was collected in a sterile Whirlpak™ bag. Fluid may also be withdrawn from the rumen via an orogastric tube. Under a suitable anaerobic atmosphere (e.g., 90% CO₂ and 10% H₂), ten-fold serial dilutions of the rumen fluid were prepared and distributed over the surface of a solid growth medium (e.g., Scott and Dehority, 1965), and the plates were incubated at 39°C for 18 to 72 h. Isolated colonies were picked with a sterile loop and the cells were spread over the surface of fresh agar medium to produce isolated colonies. The cells from a single colony were confirmed by morphological examination to represent a pure culture and were cultured and stored in the Lethbridge Research Centre ("LRC") culture collection or used as a source of enzymatic activity or genetic material.

Example 2

Screening ruminal bacteria for phytase activity

A. Phytase assays

Sample solutions (culture filtrates, cell suspensions, lysates, washes or distilled water blanks) were assayed for phytase activity by incubating 150 µl of the solution with 600 µl of substrate solution [0.2% (w/v) sodium phytate in 0.1 M sodium acetate buffer, pH 5.0] for 30 min at 37°C. The reaction was stopped by adding 750 µl of 5% (w/v) trichloroacetic acid. Released orthophosphate in the reaction mixture was measured by the method of Fiske and Subbarow (1925). Freshly prepared colour reagent [750 µl of a solution containing 4 volumes of 1.5% (w/v) ammonium molybdate in a 5.5% (v/v) sulfuric acid solution and 1 volume of a 2.7% (w/v) ferrous sulfate solution] was added to the reaction mixture and the production of phosphomolybdate was measured spectrophotometrically at 700 nm. Results were compared to a standard curve prepared with inorganic phosphate. One unit ("Unit") of phytase was defined as the amount of enzyme required to release one µmole of inorganic phosphate (P_i) per min under the assay conditions.

An improved phytase plate assay was developed which eliminated false positive results caused by microbial acid production. Bacterial isolates were grown

under anaerobic conditions on modified Scott and Dehority (1965) agar medium containing 5% (v/v) rumen fluid, 1.8% (w/v) agar and 2.0% (w/v) sodium phytate for 5 d at 37°C. Colonies were washed from the agar surface and the petri plates were flooded with a 2% (w/v) aqueous cobalt chloride solution. After a 5-min incubation at room temperature the cobalt chloride solution was replaced with a freshly prepared solution containing equal volumes of a 6.25% (w/v) aqueous ammonium molybdate solution and 0.42% (w/v) ammonium vanadate solution. Following a 5-min incubation, the ammonium molybdate solution/ammonium vanadate solution was removed and the plates examined for zones of clearing. The effectiveness of this counterstaining technique is demonstrated in Figure 1. Prior to staining, zones of clearing were evident around colonies of phytase-producing *S. ruminantium* JY35 (ATCC 55785) and lactic acid-producing *S. bovis* grown on agar medium containing phytate (Figure 1, left petri plate). The false positive zones of clearing resulting from acid production by *S. bovis* colonies were eliminated by counterstaining the plates with cobalt chloride and ammonium molybdate/ammonium vanadate solutions (Figure 1, right petri plate).

B. Phytase activity of ruminal bacteria

The phytase activities of 345 rumen bacteria from the LRC culture collection were determined (Table 1). The anaerobic technique of Hungate (1950), as modified by Bryant and Burkey (1953), or an anaerobic chamber with a 90% CO₂ and 10% H₂ atmosphere was used to cultivate the microorganisms in the LRC culture collection. Phytase screening was performed on isolates grown anaerobically (100% CO₂) in Hungate tubes with 5 mL of modified Scott and Dehority medium (1965) containing 5% (v/v) rumen fluid, 0.2% (w/v) glucose, 0.2% (w/v) cellobiose and 0.3% (w/v) starch. After 18 to 24 h incubation at 39°C, whole cells or culture supernatants were assayed for phytase activity. Selenomonads were the predominant phytase producers (93% of the isolates tested had phytase activity, Table 1). *Prevotella* was the only other genus from which a significant number of positive cultures was identified (11 phytase positive isolates out of 40 tested). A total of 29 cultures with substantial phytase activity were identified. These included 24 of the genus *Selenomonas* and 5 of the genus *Prevotella*. Twelve of these cultures (11

Selenomonas and *Prevotella* isolate) had phytase activities substantially higher than the other positive cultures (Table 2). In all instances, the phytase activity was predominantly cell associated.

Example 3

Phytase activity of *Selenomonas ruminantium* JY35 (ATCC 55785)

A. Growth and phytase production

Phytase production during growth of *S. ruminantium* JY35 (ATCC 55785) was examined. *S. ruminantium* JY35 (ATCC 55785) was grown at 39°C in Hungate tubes with 5 mL of modified Scott and Dehority broth (1965) containing 5% (v/v) ruminal fluid. Growth (protein concentration) and phytase activity (cell associated) were monitored at intervals over a 24-h time period. Maximal growth and phytase activity of *S. ruminantium* JY35 (ATCC 55785) were achieved 8-10 h after inoculation (Figure 2). Cell growth was mirrored by increases in phytase activity.

B. Localization of phytase activity

S. ruminantium JY35 (ATCC 55785) phytase activity was determined to be predominantly cell associated. Little phytase activity was detected in culture supernatants and cell washes. The phytase activity of *S. ruminantium* JY35 (ATCC 55785) was localized by electron microscopy as described by Cheng and Costerton (1973). Cells were harvested by centrifugation, washed with buffer, embedded in 4% (w/v) agar, prefixed in 0.5% glutaraldehyde solution for 30 min and fixed for 2 hours in 5% (v/v) glutaraldehyde solution. Samples were washed five times with cacodylate buffer (0.1 M, pH 7.2) and treated with 2% (w/v) osmium tetroxide, washed five times with cacodylate buffer, dehydrated in a graded ethanol series, and embedded in Spurr's resin (J. B. EM Services Inc.). Ultrathin sections were cut with a Reichert model OM U3 ultramicrotome and stained with 2% (w/v) uranyl acetate and lead citrate. Specimens were viewed with Hitachi H-500 TEM at an accelerating voltage of 75 kV. A comparison of *S. ruminantium* JY35 (ATCC 55785) cells incubated with substrate for reaction product deposition with untreated cells clearly indicated that the phytase activity was associated with the cell outer membrane

surfaces (Figure 3). Deposition of electron dense material on the outer cell surfaces of treated cells was the result of phytase activity (Figures 3A, B and C).

C. Phytase pH optimum

Initial determinations of the pH optimum of the *S. ruminantium* JY35 (ATCC 55785) phytase were conducted with whole cells. Phytase activity was optimal over a pH range of 4.0 to 5.5 (Figure 4). A second pH curve was generated with a $MgCl_2$ cell extract (Figure 5). Cells from a 100-mL overnight culture were washed twice with sterile distilled water, resuspended in 0.3 volumes of a 0.2 M $MgCl_2$ aqueous solution and incubated overnight at 0°C. The solution was clarified by centrifugation and the resulting extract was used in phytase assays. Four buffers systems were used to cover the pH range; glycine (pH 1.5 - 3.0), formate (pH 3.0 - 4.0), acetate (pH 4.0 - 5.5) and succinate (pH 5.5 - 6.5).

D. Phytase temperature optimum

The temperature optimum of the *S. ruminantium* JY35 (ATCC 55785) phytase activity was determined at pH 5.0 (0.1 M sodium acetate buffer) with $MgCl_2$ cell extract. The enzyme retained over 50% of its activity over a temperature range of 37 to 55°C (Figure 6).

E. The effect of ions and substrate concentration on phytase activity

The effect of various ions (10 mM) and substrate concentration on whole cell phytase activity were determined at pH 5.0 (0.1 M sodium acetate buffer). Phytase activity was stimulated by the addition of Ca^{++} , Na^+ , K^+ and Mg^{++} , inhibited by Fe^{++} , Zn^{++} and Mn^{++} and unaffected by Co^{++} and Ni^{++} (Figure 7). The effect of substrate concentration on phytase activity in a *S. ruminantium* JY35 (ATCC 55785) $MgCl_2$ cell extract is presented in Figure 8.

F. Molecular Weight

The molecular size of the phytase in *S. ruminantium* JY35 (ATCC 55785) was determined by zymogram analysis. A ten-fold concentrated crude $MgCl_2$ released extract was mixed with 20 μ L of sample loading buffer (Laemmli, 1970) in a

1 microtube and the microtube was placed in a boiling water bath for 5 minutes. The
2 denatured MgCl_2 extracts were resolved by SDS-PAGE on a 10% separating gel
3 topped with a 4% stacking gel (Laemmli, 1970). Following electrophoresis, the
4 phytase was renatured by soaking the gel in 1% Triton X-100 for 1 h at room
5 temperature and 0.1 M sodium acetate buffer (pH 5.0) for 1 h at 4°C. Phytase
6 activity was detected by incubating the gel for 16 h in a 0.1 M sodium acetate buffer
7 (pH 5.0) containing 0.4% sodium phytate. The gel was treated with the cobalt
8 chloride and ammonium molybdate/ammonium vanadate staining procedure
9 described for the phytase plate assays in Example 2. A single dominant activity
10 band, corresponding to a molecular mass of approximately 35 to 45 kDa, was
11 observed (Figure 9).

13 Example 4

14 Cloning of a phytase gene (*phyA*) from *Selenomonas ruminantium* JY35 (ATCC 15 55785)

16 A. Isolation of phytase positive *Escherichia coli* clone

17 Genomic DNA libraries were prepared for *S. ruminantium* JY35 (ATCC 55785)
18 according to published procedures (Hu *et al.*, 1991; Sambrook *et al.*, 1989).
19 Genomic DNA was extracted from a fresh overnight culture of *S. ruminantium* JY35
20 (ATCC 55785) using a modification of the protocol described by Priefer *et al.* (1984).
21 *S. ruminantium* JY35 (ATCC 55785) genomic DNA was partially digested with *Sau*3A
22 and gel purified to produce DNA fragments in the 2- to 10-kb range. A genomic
23 library was constructed by ligating *Bam*HI-digested, dephosphorylated pUC18 with
24 *S. ruminantium* JY35 (ATCC 55785) *Sau*3A genomic DNA fragments. *Escherichia*
25 *coli* DH5 α competent cells (Gibco BRL, Mississauga, ON) were transformed with the
26 ligation mix and 6,000 clones carrying inserts were screened for phytase activity
27 (zones of clearing) on LB phytase screening agar [LB medium, 1.0 % sodium phytate
28 (filter sterilized), 100 mM HEPES (pH 6.0 - 6.5), and 0.2 % CaCl_2] containing
29 ampicillin (100 $\mu\text{g/mL}$). A phytase-positive clone SrP.2 was isolated and phytase
30 activity confirmed through enzyme assays (Figure 10). Very high levels of phytase
31 activity were found in the medium as well as associated with the *E. coli* cells (Table

3). Plasmid DNA isolated from clone SrP.2 carried a 5.5-kb plasmid, designated pSrP.2, consisting of pUC18 containing a 2.7-kb *Sau3A* insert.

B. Confirmation of the *Selenomonas ruminantium* JY35 (ATCC 55785) origin of the 2.7-kb insert

The *S. ruminantium* JY35 (ATCC 55785) origin of the 2.7-kb insert in pSrP.2 was confirmed by Southern blot hybridization (Sambrook *et al.*, 1989). Genomic DNA isolated from *S. ruminantium* JY35 (ATCC 55785) and digested with *EcoRI* or *HindIII* was resolved on a 0.8% agarose gel. After transfer to Zeta-probe[®] membrane (BioRad Laboratories), the hybridization was performed overnight at high stringency (2 x SSC; 65°C) with the 2.7-kb fragment from pSrP.2 labelled with digoxigenin (DIG DNA labeling and detection kit; Boehringer Mannheim Canada Ltd., Laval, PQ). The blots were washed twice in 2 x SSC at room temperature; 0.1% SDS for 5 minutes and twice 0.1 x SSC; 0.1% SDS for 20 minutes at 65°C. The blots were developed according to the protocol provided with the DIG DNA labeling and detection kit (Boehringer Mannheim Canada Ltd).

The probe reacted with a 14-kb *HindIII* (Figure 11) and a 23-kb *EcoRI* (data not shown) fragment of genomic DNA and confirmed that the 2.7-kb fragment was from *S. ruminantium* JY35 (ATCC 55785) and that a single homologous sequence exists in the genome. Single copies of a sequence homologous to the 2.7-kb fragment from *S. ruminantium* JY35 (ATCC 55785) also exist in the genomes of *S. ruminantium* HD86, HD141, and HD₄ (data not shown). However restriction fragment length polymorphisms were noted for *S. ruminantium* HD86 (9- and 23-kb *EcoRI* fragments) and *S. ruminantium* HD₄ (3-kb *EcoRI* fragment and a 20-kb *HindIII* fragment). The labelled 2.7-kb fragment from pSrP.2 failed to hybridize with genomic DNA isolated from *Prevotella* sp. 46/5², *E. coli* DH5α or *A. ficuum* NRRL 3135 (data not shown).

Example 5

Characterization of *Selenomonas ruminantium* phytase gene

A. Evidence for the cloning of a phytase gene

Escherichia coli DH5 α competent cells (Gibco BRL, Mississauga, ON) were transformed with plasmids pUC18 and pSrP.2. The resulting ampicillin-resistant transformants were tested for phytase activity on LB phytase screening agar. Only *E. coli* DH5 α cells transformed with pSrP.2 produced clearing zones on LB phytase screening agar.

B. Restriction and deletion analysis of pSrP.2

The phytase gene was localized on the 2.7-kb *Sau*3A insert by restriction endonuclease and deletion analyses (Ausubel *et al.*, 1990; Sambrook *et al.*, 1989). Cells carrying plasmid pSrP.2 Δ *Sph*I, constructed by the deletion of the 1.4-kb *Sph*I fragment from pSrP.2, lacked phytase activity (Figure 12 and Figure 13, Table 3).

C. Zymogram analysis

The molecular mass of the phytase produced by *E. coli* DH5 α (pSrP.2) was determined by zymogram analysis. One mL of an overnight culture was transferred to a 1.5-mL microtube. The cells were harvested by centrifugation and washed with 0.1 M sodium acetate buffer (pH 5.5). The cell pellet was resuspended in 80 μ L of sample loading buffer (Laemmli, 1970) and the microtube was placed in a boiling water bath for 5 minutes. The resulting cell extracts were resolved by SDS-PAGE on a 10% separating gel topped with a 4% stacking gel (Laemmli, 1970) and the gel was stained for phytase activity as described in Example 3F. A single dominant activity band, corresponding to a molecular mass of approximately 37 kDa, was observed (Figure 14, lane A). A corresponding activity band was not observed for *E. coli* DH5 α (pSrP.2 Δ *Sph*I) cells (Figure 14, lane B).

D. DNA sequence analysis of pSrP.2

The complete sequence of the 2.7-kb insert of pSrP.2 was determined. Samples were prepared for DNA sequence analysis on an Applied Biosystems Model 373A DNA sequencing system (Applied Biosystems, Inc., Mississauga, ON)

1 by using a Taq DyeDeoxy™ Terminator Cycle Sequencing Kit (Applied Biosystems,
2 Inc.). Template DNA was extracted from overnight cultures of *E. coli* DH5α (pSrP.2)
3 with the Wizards™ minipreps DNA purification system (Promega Corp., Madison,
4 WI). Overlapping sequences were generated by primer walking. The DNA
5 sequence data was analyzed using MacDNASIS DNA software (Hitachi Software
6 Engineering Co., Ltd., San Bruno, CA).

7 The sequence of the 2.7-kb DNA insert was determined and DNA structural
8 analysis identified an open reading frame (ORF2; bp 1493 to 2504) overlapping the
9 *SphI* site of the 2.7-kb *Sau3A* insert and large enough to encode the 37 kDa phytase.
10 Phytase activity was eliminated by deleting bp 1518 through to the end of the 2.7-kb
11 *Sau3A* fragment (pSrPr6, Table 3, Figure 13). This was accomplished by cloning the
12 PCR product of pSrP.2 bounded by sequencing primer SrPr6 (CGG GAT GCT TCT
13 GCC AGT AT, SEQ ID NO. 3 the reverse complement of bp 1518 to 1538) and M13
14 Forward primer (CGC CAG GGT TTT CCC AGT CAC GAC) into pGEM-T (Promega
15 Corp.). A PCR product subclone (pSrPf6) of pSrP.2, bounded by primer SrPf6 (bp
16 1232 to 1252, CGT CCA CGG AGT CAC CCT AC) SEQ ID NO. 4 and M13 Reverse
17 primer (AGC GGA TAA CAA TTT CAC ACA GGA), and containing ORF2 plus 252
18 bp upstream of the *SphI* cleavage site retained phytase activity (Table 3, Figure 13).

19 The sequence and translation of the *S. ruminantium* phytase gene (*phyA*) is
20 shown in Figure 15. Translation of ORF2 would result in the expression of a 346-
21 amino acid polypeptide with a predicted molecular weight of 39.6 kDa (Figure 15).
22 The first 31 residues were typical of a prokaryote signal sequence, encompassing
23 a basic N-terminus and central hydrophobic core (von Heijne, 1986). Application of
24 the method of von Heijne (1986) predicted the signal peptidase cleavage site most
25 probably occurs before Ala²⁸ or Pro³¹. This was confirmed by determining the N-
26 terminal amino acid sequence of gel purified from *E. coli* DH5α (pSrPf6) culture
27 supernatant (Figure 15). The secreted mature protein has a putative mass of 36.5
28 kDa.

29 A comparison of the *phyA* amino acid sequence with known protein
30 sequences from the MasDNASIS SWISSPROT database revealed no significant
31 similarities to any published sequences including *Aspergillus niger* phytase genes
32 *phyA* and *phyB*.

Example 6

Partial purification and characterization of *phyA* products expressed by *E. coli*.

Cell free supernatants, prepared from overnight cultures of *E. coli* (pSrPf6), were mixed 3:1 (v/v) with Ni²⁺-NTA agarose pre-equilibrated in 0.1 M Tris (pH 7.9), 0.3 M NaCl buffer. The mixture was incubated at room temperature for 0.5 h and washed 3 x with 0.1 M Tris (pH 7.9), 0.3 M NaCl buffer. The phytase activity was eluted from the resin with 1 volume 0.1 M sodium acetate (pH 5.0), 0.3 M NaCl. When resolved on SDS-polyacrylamide gels stained with Coomassie brilliant blue, over 70% of the eluted protein formed a single 37-kDa protein band. Zymogram and N-terminal amino acid sequence analyses confirmed that the 37-kDa band corresponded to the phytase encoded by the cloned *S. ruminantium* JY35 (ATCC 55785) *phyA*. The specific activity of Ni²⁺-NTA agarose-purified phytase ranged from 200 to 400 μ mol phosphate released/min/mg protein. This is 2 to 4 times higher than the specific activity reported for the purified *A. ficuum* NRRL 3135 phytase (van Gorcum et al., 1991, 1995; van Hartingsveldt et al., 1993).

Example 7

Overexpression of the *Selenomonas ruminantium phyA* gene

Isolation and characterization of *phyA* from *S. ruminantium* JY35 (ATCC 55785) enables the large scale production of protein PhyA in any of a number of prokaryotic (e.g., *E. coli* and *B. subtilis*) or eukaryotic (e.g., fungal - *Pichia*, *Saccharomyces*, *Aspergillus*, *Trichoderma*; plant - *Brassica*, *Zea*, *Solanum*; or animal - poultry, swine or fish) expression systems using known methods. Teachings for the construction and expression of *phyA* in *E. coli*, *P. pastoris*, and *B. napus* are provided below. Similar approaches may be adopted for expression of the *S. ruminantium* JY35 (ATCC 55785) phytase in other prokaryotic and eukaryotic organisms.

A. Cloning of the *Selenomonas ruminantium* *phyA* in an *Escherichia coli* - specific expression construct

An expression construct is constructed in which the region encoding the mature *PhyA* is transcriptionally fused with the *tac* promoter (Brosius et al., 1985). The promoter sequences may be replaced by those from other promoters that provide for efficient expression in *E. coli*. The expression construct is introduced into *E. coli* cells by transformation.

i. Construction of the *E. coli* expression vector

A number of *E. coli* expression vectors based on the *tac* or related promoters are commercially available. In this example the construct will be prepared with pKK223-3 available from Pharmacia Biotech Inc. (Uppsala, Sweden). The region of *phyA* encoding the mature *PhyA* (the peptide secreted following removal of the signal peptide) is amplified with oligonucleotide primers MATE2 (GC GAA TTC ATG GCC AAG GCG CCG GAG CAG AC) (SEQ ID NO. 5) and M13 Reverse. The oligonucleotide MATE2 (SEQ ID NO. 5) was designed to contain a suitable restriction site at its terminus to allow direct assembly of the amplified product with pKK223-3. The region of *phyA* amplified with MATE2 (SEQ ID NO. 5) and M13 Reverse is digested with *EcoRI* and *SmaI* and ligated into similarly cleaved pKK223-3.

ii. Transformation of *E. coli* and *PhyA* expression

The pKK223-3::*phyA* ligation mix is used to transform competent *E. coli* cells. Strains suitable for high levels of protein expression, such as SG13009, CAG926 or CAG929 (carrying *lacI* on a plasmid such as pREP4), are employed. Transformed cells are spread on LB agar containing ampicillin (100 µg/mL) and incubated overnight at 37°C. Ampicillin-resistant colonies are screened for the presence of the desired pKK223-3::*phyA* construct by extracting pDNA and subjecting the pDNA to agarose gel electrophoresis and restriction analysis. Positive clones may be further characterized by PCR and DNA sequence analysis.

Expression of the *S. ruminantium* JY35 (ATCC 55785) phytase by transformed *E. coli* cells is tested by growing the cells under vigorous aeration at 37°C in a suitable liquid medium (e.g., LB or 2xYT) containing the appropriate antibiotic selection until the optical density (at 600 nm) is between 0.5 and 1.0. The *tac* promoter is induced by adding isopropyl-β-D-thiogalactoside (IPTG) to a final

concentration between 0.1 and 2 mM. The cells are cultivated for an additional 2 to 4 h and harvested by centrifugation. Protein expression is monitored by SDS-PAGE, and western blot/immunodetection techniques. The expressed PhyA may be extracted by breaking (e.g., sonication or mechanical disruption) the *E. coli* cells. Protein inclusions of PhyA may be harvested by centrifugation and solubilized with 1 to 2 % SDS. The SDS may be removed by dialysis, electroelution or ultrafiltration. The phytase activity of prepared cell extracts may be assayed by standard methods described in Example 2.

B. Cloning of the *Selenomonas ruminantium* phyA in a *Pichia pastoris* - specific expression construct

An expression construct is constructed in which the region encoding the mature PhyA is translationally fused with the secretion signal sequences found on *P. pastoris* expression vectors (Pichia Expression Kit Instruction Manual, Invitrogen Corporation, San Diego, CA) in order to express the *S. ruminantium* phytase as a secreted product. The promoter and secretion signal sequences may be replaced by those from other promoters that provide for efficient expression in *Pichia*. The expression construct is introduced into *P. pastoris* cells by transformation.

i. Construction of the *P. pastoris* expression vector

A number of *P. pastoris* expression vectors based on the *aox1* promoters and α -Factor or *pho1* signal sequences are commercially available. In this example the construct will be prepared with pPIC9 available from Invitrogen Corporation. The region of *phyA* encoding the mature PhyA is amplified with oligonucleotide primers MATE (GC GAA TTC GCC AAG GCG CCG GAG CAG AC) (SEQ ID NO. 6) and M13 Reverse. The oligo MATE (SEQ ID NO. 6) was designed to contain a suitable restriction site at its terminus to allow direct assembly of the amplified product with pPIC9. The region of *phyA* amplified with MATE (SEQ ID NO. 6) and M13 Reverse is digested with *EcoRI* and ligated into similarly cleaved pPIC9.

ii. Transformation of *P. pastoris* and PhyA expression

The pPIC9::*phyA* ligation mix is used to transform competent *E. coli* DH5 α cells. Transformed cells are spread on LB agar containing ampicillin (100 μ g/mL) and incubated overnight at 37°C. Ampicillin-resistant colonies are screened for the

1 presence of the desired pPIC9::*phyA* construct by extracting pDNA and subjecting
2 the pDNA to agarose gel electrophoresis and restriction analysis. Positive clones are
3 further characterized by PCR and DNA sequence analysis. Plasmid DNA is
4 prepared from a 1 L culture of an *E. coli* clone carrying the desired pPIC9::*phyA*
5 construct. The pDNA is digested with *Bgl*I and analyzed by agarose gel
6 electrophoresis to confirm complete digestion of the vector. The digested pDNA is
7 extracted with phenol:chloroform, ethanol precipitated and resuspended in sterile
8 distilled H₂O to a final concentration of 1 µg/mL. In preparation for transformation,
9 *P. pastoris* GS115 or KM71 cells are grown for 24 h at 30°C in YPD broth. Cells from
10 100 µL of culture are harvested by centrifugation and resuspended in 100 µL of
11 transformation buffer (0.1M LiCl, 0.1M dithiothreitol, 45% polyethylene glycol 4000)
12 containing 10 µg salmon sperm DNA and 10 µg of linearized pPIC9::*phyA*. The
13 mixture is incubated for 1 h at 37°C, spread on *P. pastoris* minimal agar medium and
14 incubated for 2 to 5 d. Colonies growing on the minimal agar medium are streaked
15 for purity and analyzed for the presence of the integrated *phyA* by PCR and
16 Southern blot hybridization.

17 Expression of the *S. ruminantium* JY35 (ATCC 55785) phytase by
18 transformed *P. pastoris* cells is tested by growing the cells at 30°C under vigorous
19 aeration in a suitable liquid medium (e.g. buffered complex glycerol medium such as
20 BMGY) until a culture optical density (at 600 nm) (OD₆₀₀) of 2 to 6 is reached. The
21 cells are harvested and resuspended to an OD₆₀₀ of 1.0 in an inducing medium (e.g.,
22 buffered complex methanol medium, BMMY) and incubated for a further 3 to 5 days.
23 Cells and cell-free culture supernatant are collected and protein expression is
24 monitored by enzyme assay, SDS-PAGE, and western blot/immunodetection
25 techniques.

26
27 C. Cloning of the *Selenomonas ruminantium phyA* in a *Pichia pastoris* - specific
28 expression construct - A Further Example

29 An expression construct is constructed in which the region encoding the
30 mature PhyA is translationally fused with the secretion signal sequences found on
31 *P. pastoris* expression vectors (e.g., *Pichia* Expression Kit Instruction Manual,
32 Invitrogen Corporation, San Diego, CA) in order to express the *S. ruminantium*

1 phytase as a secreted product. The promoter and secretion signal sequences may
2 be replaced by those from other promoters that provide for efficient expression in
3 *Pichia*. The expression construct is introduced into *P. pastoris* cells by
4 transformation.

5 i. Construction of the *P. pastoris* expression vector

6 A number of *P. pastoris* expression vectors based on the *aox1* promoters and
7 α -Factor or *pho1* signal sequences are commercially available. In this example the
8 construct was prepared with pPICZ α A available from Invitrogen Corporation. The
9 region of *phyA* encoding the mature PhyA (i.e., the peptide secreted following
10 removal of the signal peptide) was amplified with oligonucleotide primers MATE (GC
11 GAA TTC GCC AAG GCG CCG GAG CAG AC SEQ ID NO. 6) and M13 Reverse.
12 The oligo MATE (SEQ ID NO. 6) was designed to contain an *EcoRI* restriction site
13 at its terminus to allow direct assembly of the amplified product with pPICZ α A. The
14 region of *phyA* amplified with MATE (SEQ ID NO. 6) and M13 Reverse was digested
15 with *EcoRI* and ligated into similarly cleaved pPICZ α A.

16 ii. Transformation of *P. pastoris*

17 The pPICZ α A::*phyA* ligation mix was used to transform competent *E. coli*
18 DH5 α cells. Transformed cells were spread on LB agar containing Zeocin (25
19 mg/mL) and incubated overnight at 37°C. Zeocin resistant colonies were screened
20 for the presence of the desired pPICZ α A::*phyA* construct by extracting pDNA and
21 subjecting the pDNA to agarose gel electrophoresis and restriction analysis. Positive
22 clones were further characterized by PCR and DNA sequence analysis. Plasmid
23 DNA was prepared from a 1 L culture of an *E. coli* clone carrying the desired
24 pPICZ α A::*phyA* construct. The pDNA is digested with *BglII* and analyzed by
25 agarose gel electrophoresis to confirm complete digestion of the vector. The
26 digested pDNA was extracted with phenol:chloroform, ethanol precipitated and
27 resuspended in sterile distilled H₂O to a final concentration of 1 μ g/ μ L.

28 In preparation for transformation, 50 mL of YPD broth were inoculated with *P.*
29 *pastoris* GS115 cells and incubated at 28°C and 250 RPM for 1 day. Subsequently,
30 5 mL of the 1 d culture was used to inoculate 50 mL of fresh YPD broth. The culture
31 was propagated overnight at 28°C and 250 RPM. The following morning, 5 mL of
32 this culture was used to inoculate 50 mL of fresh YPD broth. This culture was

1 incubated at 28°C and 250 RPM until the culture OD₆₀₀ reached approximately 1.2
2 (~ 6 h). The yeast cells from 20 ml of fresh culture were harvest by centrifugation,
3 washed once with and resuspended in 1 mL of room temperature 10 mM Tris, 1 mM
4 EDTA, 0.1 M LiCl, 0.1 M dithiothreitol buffer (pH 7.4). After a 1 h incubation at 30°C,
5 the cell suspension was washed once with 1 mL ice cold water and once with 1 mL
6 ice cold 1 M sorbitol. The cells were resuspended in 160 µL of ice cold 1 M sorbitol
7 (to obtain cell concentrations approaching 10¹⁰ cells/mL). Linearized
8 pPICZαA::phyA (5 to 10 µg) was mixed with 80 µL of cells, loaded into prechilled
9 electroporation cuvettes (0.2 cm inter-electrode distance) and incubated on ice for
10 5 min. A high voltage pulse (1.5 kV, 25 µF, 200 Ohms) was applied to the cuvette
11 with a Bio-Rad Gene Pulser™. Immediately following the pulse, 1 mL of ice cold 1M
12 sorbitol was added to the cuvette which was incubated subsequently for 2 h at 30°C.
13 The cell suspension was spread (100 to 200 µL per plate) on YPD agar medium
14 containing Zeocin (100 µg/mL) and incubated for 2 to 4 d at 30°C. Colonies growing
15 on the selective medium were streaked for purity and analyzed for the presence of
16 the integrated *phyA* by PCR and/or Southern blot hybridization.

17 iii. *Pichia pastoris* expression of the *S. ruminantium* JY35 phytase gene

18 Expression of the *S. ruminantium* JY35 phytase by transformed *P. pastoris*
19 cells was tested by growing transformed cells grown overnight in buffered complex
20 glycerol medium (e.g., buffered complex glycerol medium, BMGY, *Pichia* Expression
21 Kit Instruction Manual) at 28°C and 250 RPM and transferring them into inducing
22 medium (e.g., buffered complex methanol medium, BMMY). The cells harvested
23 from the BMGY medium were washed once with BMMY medium, resuspended in
24 BMMY to an OD₆₀₀ of 1.0 and incubated for a further 3 to 5 days at 28°C and 250
25 RPM. Methanol (0.005 volumes) was added every 24 h. Cells and cell free culture
26 supernatants were collected and assayed for phytase activity.

27 Sixteen *P. pastoris* pPICZαA::MATE transformants were tested for phytase
28 activity following 96 h growth in BMMY medium. The most active transformant,
29 named clone 17, was selected for further study. Growth and phytase production by
30 *P. pastoris* pPICZαA::MATE clone 17 and a negative clone (*P. pastoris* pPICZαA)
31 were monitored over a period of 9 d. Starter cultures were prepared by growing the
32 isolates overnight (28°C, 250 RPM) in 10 mL of BMGY (glycerol) medium. The cells

1 were harvested and duplicate cultures were prepared by resuspending the cells in
2 50 mL BMMY (methanol) medium to an approximate OD_{600} of 2.5. The resulting
3 cultures were transferred into 500 mL flasks and incubated at 28°C and 250 RPM.
4 Methanol was added every 24 h to a final concentration of 0.5%. Optical density and
5 phytase activity were measured over the time course of the experiment. The results
6 are presented in Table 4. Phytase activity was detected only in cultures carrying the
7 *S. ruminantium phyA* gene. These cultures produced up to 22.5 units of phytase
8 activity per mL after 210.5 h cultivation.

9 Phytase activity in shake flask cultures was increased through modification of
10 the induction protocol and medium composition. The phytase activity of clone 17 was
11 dramatically improved by increasing the initial cell density ($OD_{610} = 36.0$) of the
12 induced culture. After nearly 4 d growth (91.5 h), phytase activities greater than 40
13 and 20 units/mL were observed for whole culture and cell free supernatant samples,
14 respectively. The optical densities (OD_{610}) of these cultures were between 62 and
15 69. Experimental results suggest that the greater the culture biomass at the time
16 of methanol induction, the greater the yields of recombinant phytase. Biomass yields
17 as high as 150 g/L (dry weight) or optical densities of 1500 have been reported for
18 *Pichia* cultivated under optimal growth conditions in a tightly controlled fermentor
19 system operating with oxygen enrichment.

20 *Pichia* phytase yields were also increased by adding Tween-80 to the
21 medium. Surfactants have been shown previously to affect phytase production by
22 *Aspergillus carbonarius* (Al-Asheh and Duvnjak, 1994). The effect of incorporating
23 0, 0.02, 0.1 or 0.5 % Tween-80 on phytase yields of BMMY cultures of *P. pastoris*
24 pPICZ α A::MATE clone 17 is illustrated in Table 5. The cells from 2 d YPD cultures
25 were harvested and resuspended in BMMY ($OD_{610} = 8.3$). Triplicate flasks for each
26 concentration of Tween-80 were prepared and incubated at 28°C and 250 RPM.
27 Methanol (0.005 volumes) was added on a daily basis to the flasks. Phytase activity
28 increased more rapidly in cultures containing higher concentrations of Tween-80.
29 Furthermore, a larger proportion of the phytase activity was found in the supernatant
30 when higher Tween-80 concentrations were used. Phytase yields as high as 298
31 units/mL of shake flask culture have been achieved with a 9 d culture of clone 17
32 cultivated in BMMY medium amended with 0.5% Tween-80.

Cellular and supernatant proteins were analyzed by SDS-PAGE to confirm the production of PhyA by *P. pastoris*. The presence of a 37 kDa protein band was readily apparent when as little as 5 μ L of supernatant was resolved on a 12% SDS-PAGE gel. The 37 kDa band was visible in the cellular protein sample but represented less than 10% of that found in the corresponding amount of supernatant. In addition to PhyA, supernatants from clone 17 contained very few additional proteins (a useful characteristic of *Pichia* expression). The recombinant PhyA protein comprised over 95% (estimated from SDS-PAGE gels) of the secreted protein. The 37 kDa protein band was not present in the supernatant or cells of a negative control culture (*P. pastoris* pPICZ α A).

Shake flask experiments with recombinant *P. pastoris* cells expressing the *S. ruminantium* phytase (PhyA) have demonstrated the potential of this protein production system. Significant gains in phytase yields will be obtained by cultivating and inducing clone 17 in a fermentor. Additional gains in phytase yields may be achieved by increasing gene copy number through further screening of independent transformants or the use of multicopy vector systems. Spontaneous multiple plasmid integration events occur in *Pichia* at a frequency between 1/10 and 1/100 transformants. It is not unrealistic to expect that a 10 fold gain in phytase yield (e.g., 3,000 units/mL) may be readily achieved through manipulation of phytase gene copy number and control of fermentation parameters. This would result in production levels comparable to commercial *A. ficuum* phytase production systems. Yields for these systems are believed to be around 3,000,000 units (μ mol Pi released/min) of phytase activity per L of culture.

iv. The Activity of recombinant the *S. ruminantium* phytase (PhyA) on grain substrates

The liberation of phosphate from corn by the recombinant *S. ruminantium* JY35 phytase produced by *Pichia pastoris* was examined. Feed corn was ground and sieved through a mesh to obtain a particle size between 1 - 3 mm. Ground corn (0.5 g) was weighed into sterile 15 mL Falcon tubes to which 2 mL of 0.1 M sodium acetate buffer (pH 5.0) was added. After addition of phytase, the reaction mixtures were incubated at 37°C. Phosphate release was determined by measuring supernatant phosphate. In order to measure the background phosphate, reaction

1 mixtures were prepared and terminated immediately through the addition of 5% (w/v)
2 TCA. All experiments were conducted in triplicate.

3 Incubation of corn in a sodium acetate buffer resulted in the release of
4 increasing amounts of phosphorus over time (Table 6). Although the addition of
5 phytase activity significantly increased the amount of phosphorus released, the rate
6 of phosphorus release decreased with time.

7 The concentration of phytase added to the incubation mixture also influenced
8 the amount of phosphorus released. Raising phytase concentrations from 0.08 units
9 to 0.48 units per g of corn resulted in increased levels of phosphorus in the
10 supernatant (Table 7). It should be noted that increasing the phytase concentration
11 from 0.32 to 0.48 units produced only a marginal increase in phosphorus released.
12

13 D. Cloning of the *Selenomonas ruminantium* phyA in a *Brassica napus* seed -
14 specific expression construct

15 Transformation and gene expression methods have been developed for a
16 wide variety of monocotyledonous and dicotyledonous crop species. In this example,
17 a *S. ruminantium* JY35 (ATCC 55785) phytase expression construct is constructed
18 in which the region encoding the mature PhyA is translationally fused with an oleosin
19 coding sequence in order to target seed oil body specific expression of the *S.*
20 *ruminantium* phytase. The promoter and/or secretion signal sequences may be
21 replaced by those from other promoters that provide for efficient expression in *B.*
22 *napus* or other transformable plant species. The expression construct is introduced
23 into *B. napus* cells by *Agrobacterium*-mediated transformation.

24 i. Construction of the *B. napus* expression vector

25 A number of expression vectors functional in *B. napus* are described in the
26 literature (Gelvin et al., 1993). In this example, the construct is prepared by replacing
27 the *E. coli* β -glucuronidase CDS of pCGBPGUS (van Rooijen and Moloney, 1994)
28 with a fragment encoding the *phyA* mature CDS. This is accomplished by
29 subcloning the pCGBPGUS *Pst*I *Kpn*I fragment, containing the oleosin
30 promoter::oleosin CDS:: β -glucuronidase CDS::NOS region, on to *Pst*I *Kpn*I- digested
31 pUCBM20 (Boehringer Mannheim Canada, Laval, PQ). This plasmid is called
32 pBMOBPGUS. The region of *phyA* encoding the mature PhyA is amplified with

oligonucleotide primers MATN (GA GGA TCC ATG GCC AAG GCG CCG GAG CAG AC) (SEQ ID NO. 7) and M13 Reverse. The oligonucleotide MATN (SEQ ID NO. 7) was designed to contain a suitable restriction site at its terminus to allow direct assembly of the amplified product with digested pBMOBPGUS. The *phyA* fragment amplified with MATN (SEQ ID NO. 7) and M13 Reverse is digested with *NcoI* *SstI* and ligated into similarly cleaved pBMOBPGUS to generate plasmid pBMOBP*phyA*. The *B. napus* expression vector, pCGOBP*phyA*, is constructed by replacing the *PstI* *KpnI* fragment from pCGOBPGUS with the *PstI* *KpnI* fragment from pBMOBP*phyA*, containing the oleosin promoter::oleosin CDS::phyA CDS::NOS fragment.

ii. Transformation of *B. napus* and stable PhyA expression

Transgenic *B. napus* is prepared as described by van Rooijen and Moloney (1994). *Agrobacterium tumefaciens* strain EHA101 is transformed by electroporation with pCGOBP*phyA*. Cotyledonary petioles of *B. napus* are transformed with *A. tumefaciens* EHA101 (pCGOBP*phyA*). Transgenic plants are regenerated from explants that root on hormone-free MS medium containing 20 µg/mL kanamycin. Young plants are assayed for NPTII activity, grown to maturity and allowed to self pollenate and set seed. Seeds from individual transformants are pooled and part of the seed sample is assayed for the presence of phytase activity and compared to seeds from untransformed plants. Second generation plants (T2) are propagated from the seeds of clones with the highest levels of phytase activity. Seeds from the T2 plants homozygous for NPTII (hence also for *phyA*) are selected and used for mass propagation of plants (T3) capable of producing the highest amounts of phytase.

Example 8

Identification of Related Phytase Genes in Other Microorganisms

To identify a phytase gene related to *phyA*, hybridization analysis can be used to screen nucleic acids from one or more ruminal isolates of interest using *phyA* (SEQ ID NO. 1) or portions thereof as probes by known techniques (Sambrook, 1989; Ausubel, 1990) as described in example 4B. Related nucleic acids may be cloned by employing known techniques. Radioisotopes (i.e., ³²P) may be required

1 when screening organisms with complex genomes in order to increase the sensitivity
2 of the analysis. Polymerase chain reaction (PCR) amplification may also be used to
3 identify genes related to *phyA*. Related sequences found in pure or mixed cultures
4 are preferentially amplified by PCR (and variations of such as Reverse Transcription
5 - PCR) with oligonucleotides primers designed using SEQ ID NO. 1. Amplified
6 products may be visualized by agarose gel electrophoresis and cloned using known
7 techniques. A variety of materials, including cells, colonies, plaques, and extracted
8 nucleic acids (e.g., DNA, RNA), may be examined by these techniques for the
9 presence of related sequences. Alternatively, known immunodetection techniques
10 employing antibodies specific to PhyA (SEQ ID NO. 2) can be used to screen whole
11 cells or extracted proteins of interest for the presence of related phytase(s).
12

Table 1. Phytase activity among rumen bacteria.

Phytase Activity	Microorganism	Number of isolates tested
Very Strong	<i>Prevotella</i> sp.	1
	<i>Selenomonas ruminantium</i>	11
Strong	<i>Prevotella ruminicola</i>	4
	<i>S. ruminantium</i>	13
Moderate	<i>Bacillus</i> sp.	1
	<i>Megasphaera elsdenii</i>	7
	<i>P. ruminicola</i>	6
	<i>S. ruminantium</i>	37
	<i>Treponema</i> sp.	1
Negative	<i>Anaerovibrio lipolytica</i>	2
	<i>Bacillus</i> sp.	4
	<i>Butyrivibrio fibrisolvens</i>	47
	<i>Clostridium</i> sp.	1
	<i>Coprococcus</i> sp.	3
	<i>Enterococcus</i> sp.	4
	<i>Eubacterium</i> sp.	7
	<i>Fibrobacter succinogenes</i>	8
	<i>Fusobacterium</i> sp.	3
	<i>Lachnospira multiparus</i>	4
	<i>Lactobacillus</i> sp.	20
	<i>M. elsdenii</i>	7
	<i>Peptostreptococcus</i> sp.	1
	<i>P. ruminicola</i>	41
	<i>Ruminobacter amylophilus</i>	4
	<i>Ruminococcus albus</i>	7
	<i>Ruminococcus flavefaciens</i>	10
	<i>S. ruminantium</i>	4
	<i>Streptococcus bovis</i>	48
	<i>Streptococcus milleri</i>	1
	<i>Staphylococcus</i> sp.	6
	<i>Succinovibrio dextrisolvans</i>	12
	<i>Treponema</i> sp.	12
	Unknown	8
Total isolates screened		345

Table 2. Phytase activity of selected rumen bacterial isolates.

Isolate	Phytase activity (mU*/mL)
<i>Selenomonas ruminantium</i> JY35	646
<i>Selenomonas ruminantium</i> KJ118	485
<i>Selenomonas ruminantium</i> BS131	460
<i>Selenomonas ruminantium</i> HD141	361
<i>Selenomonas ruminantium</i> HD86	286
<i>Selenomonas ruminantium</i> JY135	215
<i>Selenomonas ruminantium</i> D	69
<i>Selenomonas ruminantium</i> HD16	52
<i>Selenomonas ruminantium</i> BS114	47
<i>Selenomonas ruminantium</i> JY4	27
<i>Prevotella</i> sp. 46/5 ²	321
<i>Prevotella ruminicola</i> JY97	68
<i>Prevotella ruminicola</i> KJ182	61
<i>Prevotella ruminicola</i> JY106	49
<i>Megasphaera elsdenii</i> JY91	5

*U= μ moles, P_i released/min

Table 3. Overexpression of *S. ruminantium*¹ phytase in recombinant *E. coli* DH5 α .

Strain	Sample Composition	Units ² /mL	Specific Activity (Units/mg protein)
<i>E. coli</i> (pSrP.2)	cells	0.30 (0.08) ³	1.56 (0.41)
	supernatant	0.308 (0.21)	2.64 (1.51)
<i>E. coli</i> (pSrPf6)	cells	0.91 (0.41)	6.42 (0.64)
	supernatant	5.10 (0.58)	22.83 (1.67)
<i>E. coli</i> (pSrP.2 <i>SphI</i>)	cells	ND ⁴	ND
	supernatant	ND	ND

¹*S. ruminantium* JY35 is a crescent shaped-rod, an obligate anaerobe, produces propionic acid from the fermentation of glucose, ferments lactose, does not ferment glycerol, does not ferment mannitol (see also Bergey's Manual of Systematic Bacteriology, ed. John G. Holt, Williams and Wilkins, Baltimore, 1984)

²Units = μ moles P_i released/min

³Numbers in parenthese are standard errors

⁴ND = not detected

Table 4. Growth and phytase activity of *P. pastoris* cells transformed with pPICZ α A (negative control) or pPICZ α A::MATE (clone 17).

Culture	Time (h)	Optical Density (610 nm)	Phytase activity (μ mol/min/mL)	
			Culture	Supernatant
<i>P. pastoris</i> (pPICZ α A)	0.0	2.6	0.0	0.0
	20.5	10.1	0.0	0.0
	42.5	17.8	0.0	0.0
	68.0	17.0	0.0	0.0
	91.0	28.5	0.0	0.0
	138.5	39.3	0.0	0.0
	210.5	46.7	0.0	0.0
<i>P. pastoris</i> (pPICZ α A::MATE)	0.0	2.5	0.0	0.0
	20.5	11.3	1.9	0.1
	42.5	13.9	4.4	1.5
	68.0	12.9	8.0	2.7
	91.0	15.7	4.7	0.5
	138.5	18.3	12.6	5.3
	210.5	18.7	22.5	12.5

Table 5. The effect of Tween-80 concentration on growth and phytase activity of *P. pastoris* cells transformed with pPICZ α A::MATE (clone 17).

Time (d)	Sample (% Tween-80)	Optical Density (610 nm)	Phytase Activity (μ mol/min/mL)		Supernatant/ Culture Activity
			Culture	Supernatant	
2	0.0	24.3	4.1	2.2	0.55
	0.02	24.4	4.8	2.7	0.57
	0.1	25.1	5.2	3.2	0.61
	0.5	24.4	4.9	3.2	0.65
4	0.0	31.2	6.9	4.7	0.69
	0.02	31.0	8.2	5.5	0.67
	0.1	31.8	10.3	6.9	0.67
	0.5	29.2	10.3	9.1	0.88
8	0.0	32.8	10.6	5.9	0.55
	0.02	30.4	14.8	9.8	0.67
	0.1	33.9	20.2	17.2	0.86
	0.5	33.8	22.1	18.9	0.86

Table 6. The effect of incubation period and recombinant *S. ruminantium* JY35 phytase (2 units/g of corn) on phosphate release from corn.

Sample	Length of incubation (h)	Phosphate concentration (μmoles/mL)
No phytase	1	0.85
	2	1.72
	3	2.56
	4	3.77
	5	4.35
Phytase	1	4.76
	2	6.83
	3	7.72
	4	8.41
	5	8.49

Table 7. The effect of recombinant *S. ruminantium* JY35 phytase concentration on phosphate release from corn.

Phytase activity (units/g of corn)	Phosphate concentration (μmoles/g of corn)
0.08	11.8
0.16	14.8
0.24	22.5
0.32	23.0
0.40	23.2
0.48	23.8
0.56	23.8
0.64	23.6
0.72	23.8

REFERENCES

- Al-Asheh, S. And Z. Duvnjak. 1994. The effect of surfactants on the phytase production and the reduction of the phytic content in canola meal by *Aspergillus carbonarius* during a solid state fermentation process. *Biotechnol. Lett.* 16:183-188.
- Ausubel, F.A., R. Brent, R.E. Kingston, D.D. Moore, J.G. Sneiderman, J.A. Smith and K. Struhl. (eds.) 1990. Current protocols in molecular biology. Green Publishing and Wiley-Interscience, New York.
- Brosius, J., M. Erfle and J. Storella. 1985. Spacing of the -10 and -35 regions in the *tac* promoter. *J. Biol. Chem.* 260:3539-3541.
- Bryant, M.P. and L.A. Burkey. 1953. Cultural methods and some characteristics of some of the numerous groups of bacteria in the bovine rumen. *J. Dairy Sci.* 36:205-217.
- Cheng, E.W., G. Hall and W. Burroughs. 1955. A method for the study of cellulose digestion by washed suspensions of rumen microorganisms. *J. Dairy Sci.* 38:1255-1230.
- Cheng, K.-J. and J.W. Costerton, 1973. Localization of alkaline phosphatase in three Gram-negative rumen bacteria. *J. Bacteriol.* 116:424-440.
- Dayhoff, M.O., R.M. Schwartz and B.C. Orcutt. 1978. A model of evolutionary change in proteins. In: Atlas of Protein Sequence and Structure. Volume 5, Supplement 3, Chapter 22, pp 345-352.
- Ellis, S.B., P.F. Brust, P.J. Koutz, A.F. Waters, M.M. Harpold and R.R. Gingeras. 1985. Isolation of alcohol oxidase and two other methanol regulated genes from the yeast, *Pichia pastoris*. *Mol. Cell. Biol.* 5:1111-1121.

- 1 Fiske, C.H. and Y. Subbarow. 1925. The colorimetric determination of phosphorus.
2 J. Biol. Chem. 66:376-400.
3
- 4 Gelvin, S.B., R.A. Schilperoort and D.P.S. Verma (eds.). 1993. Plant Molecular
5 Biology Manual. Kluwer Academic Publishers, Boston, MA.
6
- 7 Graf, E. (ed.). 1986. Phytic acid, chemistry and applications. Pilatus Press.
8 Minneapolis, MN. 344 pp.
9
- 10 Howson, S.J. and R.P. Davis. 1983. Production of phytate-hydrolysing enzyme by
11 some fungi. Enzyme Microb. Technol. 5:377-382.
12
- 13 Hu, Y.J., D.C. Smith, K.-J. Cheng and C.W. Forsberg. 1991. Cloning of a xylanase
14 gene from *Fibrobacter succinogenes* 135 and its expression in *Escherichia*
15 *coli*. Can.J. Microbiol. 37:554-561.
16
- 17 Hungate, R.E. 1950. The anaerobic mesophilic cellulolytic bacteria. Bacteriol. Rev.
18 14:1-49.
19
- 20 Laemmli, U.K. 1970. Cleavage of the structural proteins during assembly of the
21 head of bacteriophage T4. Nature 227:680-685.
22
- 23 Priefer, U., R. Simon and A. Puhler. 1984. Cloning with cosmids. In: Puhler, A. and
24 K.N. Timmis (eds) Advanced molecular genetics. Springer-Verlag, New York.
25 pp.190-201.
26
- 27 Raun, A., E. Cheng and W. Burroughs. 1956. Phytate phosphorus hydrolysis and
28 availability to rumen microorganisms. Agric. Food Chem. 4:869-871.
29
- 30 Sambrook, J., E.F. Fritsch and T. Maniatis. 1989. Molecular cloning. A laboratory
31 manual. 2nd. edn. Cold Spring Harbor Laboratory Press. Cold Spring Harbor,
32 NY.

- 1 Scott, H.W. and B.A. Dehority. 1965. Vitamin requirements of several cellulolytic
2 bacteria. J. Bacteriol. 89:1169-1175.
3
- 4 Shieh, T.R. and J.H. Ware. 1968. Survey of microorganisms for the production of
5 extracellular phytase. Appl. Microbiol. 16:1348-1351.
6
- 7 van Gorcom, R.F.M. and C.A.M.J Van Den Hondel. 1993. Cloning, characterization
8 and overexpression of the phytase gene (*phyA*) of *Aspergillus niger*. Gene
9 127:87-94.
10
- 11 van Hartingsveldt W., C.M.J. van Zeijl, M.G. Hartevelde, R.J. Gouka, M.E.G.
12 Suykerbuyk, R.G.M. Luiten, P.A. Van Paridon, G.C.M. Selten, A.E. Veenstra,
13 van Rooijen, G.J.H. and M. M. Moloney. 1994. Plant seed oil-bodies as
14 carriers for foreign proteins. Bio/Technology 13:72-77.
15
- 16 von Heijne, G. 1986. A new method for predicting signal sequence cleavage sites.
17 Nucleic Acids Res. 14:4683-4690.
18
- 19 Wong, S.-L. 1989. Development of an inducible and enhancible expression and
20 secretion system in *Bacillus subtilis*. Gene 83:215-223.
21

22 All publications mentioned in this specification are indicative of the level of skill
23 of those skilled in the art to which this invention pertains. All publications are herein
24 incorporated by reference to the same extent as if each individual publication was
25 specifically indicated to be incorporated by reference.

26 Although the foregoing invention has been described in some detail by way
27 of illustration and example for purposes of clarity and understanding, it will be
28 obvious that certain changes and modifications may be practised within the scope
29 of the claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Cheng, Kuo-Joan
Selinger, Leonard B.
Yanke, Lindsey J.
Bae, Hee-Dong
Zhou, Lu Ming
Forsberg, Cecil W.
- (ii) TITLE OF INVENTION: DNA sequences encoding phytases of
ruminant microorganisms.
- (iii) NUMBER OF SEQUENCES: 7
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: McKay-Carey & Company
 - (B) STREET: 2125, 10155-102 St.
 - (C) CITY: Edmonton
 - (D) STATE: Alberta
 - (E) COUNTRY: CA
 - (F) ZIP: T5J 4G8
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: May 23, 1997
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Mary Jane McKay-Carey
 - (B) REGISTRATION NUMBER:
 - (C) REFERENCE/DOCKET NUMBER: 37003W00
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (403) 424-0222
 - (B) TELEFAX: (403) 421-0834

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1401 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: circular
- (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Selenomonas ruminantium*

(B) STRAIN: JY35

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: Genomic DNA library

(B) CLONE: pSrP.2

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 231..1268

(C) IDENTIFICATION METHOD: experimental

(D) OTHER INFORMATION: /codon_start= 231

/function= "Dephosphorylation of phytic acid"

/product= "Phytase"

/evidence= EXPERIMENTAL

/gene= "phyA"

/number= 1

/standard_name= "myo-inositol hexaphosphate
phosphohydrolase"

/citation= ([1])

(ix) FEATURE:

(A) NAME/KEY: sig_peptide

(B) LOCATION: 231..311

(C) IDENTIFICATION METHOD: experimental

(D) OTHER INFORMATION: /codon_start= 1

/function= "phytase secretion"

/product= "Signal peptide"

/evidence= EXPERIMENTAL

/citation= ([1])

(ix) FEATURE:

(A) NAME/KEY: mat_peptide

(B) LOCATION: 312..1268

(C) IDENTIFICATION METHOD: experimental

(D) OTHER INFORMATION: /codon_start= 312

/product= "Phytase"

/evidence= EXPERIMENTAL

/number= 2

/citation= ([1])

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGTCCACGGA GTCACCCTAC TATACGACGT ATGTGAAGTT CACGTCGAAG TTCTAGGGAA 60

TCACCGATTC GTGCAGGATT TTACCACTTC CTGTTGAAGC GGATGAGAAG GGGAACCGCG 120

AAGCGGTGGA AGAGGTGCTG CACGACGGAC GATCGCGCTG AATGAATCAG TGCTTCCTAA 180

CTATTGGGAT TCCGCGCAGA CGCGCGGATG GAGTAAAGGA GTAAGTTGTT ATG AAA	236
Met Lys	
-27	
TAC TGG CAG AAG CAT GCC GTT CTT TGT AGT CTC TTG GTC GGC GCA TCC	284
Tyr Trp Gln Lys His Ala Val Leu Cys Ser Leu Leu Val Gly Ala Ser	
-25 -20 -15 -10	
CTC TGG ATA CTG CCG CAG GCC GAT GCG GCC AAG GCG CCG GAG CAG ACG	332
Leu Trp Ile Leu Pro Gln Ala Asp Ala Ala Lys Ala Pro Glu Gln Thr	
-5 1 5	
GTG ACG GAG CCC GTT GGG AGC TAC GCG CGC GCG GAG CGG CCG CAG GAC	380
Val Thr Glu Pro Val Gly Ser Tyr Ala Arg Ala Glu Arg Pro Gln Asp	
10 15 20	
TTC GAG GGC TTT GTC TGG CGC CTC GAC AAC GAC GGC AAG GAG GCG TTG	428
Phe Glu Gly Phe Val Trp Arg Leu Asp Asn Asp Gly Lys Glu Ala Leu	
25 30 35	
CCG CGT AAT TTC CGC ACG TCG GCT GAC GCG CTG CGC GCG CCG GAG AAG	476
Pro Arg Asn Phe Arg Thr Ser Ala Asp Ala Leu Arg Ala Pro Glu Lys	
40 45 50 55	
AAA TTC CAT CTC GAC GCC GCG TAT GTA CCG TCG CGC GAG GGC ATG GAT	524
Lys Phe His Leu Asp Ala Ala Tyr Val Pro Ser Arg Glu Gly Met Asp	
60 65 70	
GCA CTC CAT ATC TCG GGC AGT TCC GCA TTC ACG CCG GCG CAG CTC AAG	572
Ala Leu His Ile Ser Gly Ser Ser Ala Phe Thr Pro Ala Gln Leu Lys	
75 80 85	
AAC GTT GCC GCG AAG CTG CGG GAG AAG ACG GCT GGC CCC ATC TAC GAT	620
Asn Val Ala Ala Lys Leu Arg Glu Lys Thr Ala Gly Pro Ile Tyr Asp	
90 95 100	
GTC GAC CTA CGG CAG GAG TCG CAC GGC TAT CTC GAC GGT ATC CCC GTG	668
Val Asp Leu Arg Gln Glu Ser His Gly Tyr Leu Asp Gly Ile Pro Val	
105 110 115	
AGC TGG TAC GGC GAG CGC GAC TGG GCA AAT CTC GGC AAG AGC CAG CAT	716
Ser Trp Tyr Gly Glu Arg Asp Trp Ala Asn Leu Gly Lys Ser Gln His	
120 125 130 135	
GAG GCG CTC GCC GAC GAG CGG CAC CGC TTG CAC GCA GCG CTC CAT AAG	764
Glu Ala Leu Ala Asp Glu Arg His Arg Leu His Ala Ala Leu His Lys	
140 145 150	
ACG GTC TAC ATC GCG CCG CTC GGC AAG CAC AAG CTC CCC GAG GGC GGC	812
Thr Val Tyr Ile Ala Pro Leu Gly Lys His Lys Leu Pro Glu Gly Gly	
155 160 165	
GAA GTC CGC CGC GTA CAG AAG GTG CAG ACG GAA CAG GAA GTC GCC GAG	860
Glu Val Arg Arg Val Gln Lys Val Gln Thr Glu Gln Glu Val Ala Glu	
170 175 180	

GCC GCG GGG ATG CGC TAT TTC CGC ATC GCG GCG ACG GAT CAT GTC TGG	908
Ala Ala Gly Met Arg Tyr Phe Arg Ile Ala Ala Thr Asp His Val Trp	
185 190 195	
CCA ACG CCG GAG AAC ATC GAC CGC TTC CTC GCG TTT TAC CGC ACG CTG	956
Pro Thr Pro Glu Asn Ile Asp Arg Phe Leu Ala Phe Tyr Arg Thr Leu	
200 205 210 215	
CCG CAG GAT GCG TGG CTC CAT TTC CAT TGT GAA GCC GGT GTC GGC CGC	1004
Pro Gln Asp Ala Trp Leu His Phe His Cys Glu Ala Gly Val Gly Arg	
220 225 230	
ACG ACG GCG TTC ATG GTC ATG ACG GAT ATG CTG AAG AAC CCG TCC GTA	1052
Thr Thr Ala Phe Met Val Met Thr Asp Met Leu Lys Asn Pro Ser Val	
235 240 245	
TCG CTC AAG GAC ATC CTC TAT CGC CAG CAC GAG ATC GGC GGC TTT TAC	1100
Ser Leu Lys Asp Ile Leu Tyr Arg Gln His Glu Ile Gly Gly Phe Tyr	
250 255 260	
TAC GGG GAG TTC CCC ATC AAG ACG AAG GAT AAA GAT AGC TGG AAG ACG	1148
Tyr Gly Glu Phe Pro Ile Lys Thr Lys Asp Lys Asp Ser Trp Lys Thr	
265 270 275	
AAA TAT TAT AGG GAA AAG ATC GTG ATG ATC GAG CAG TTC TAC CGC TAT	1196
Lys Tyr Tyr Arg Glu Lys Ile Val Met Ile Glu Gln Phe Tyr Arg Tyr	
280 285 290 295	
GTG CAG GAG AAC CGC GCG GAT GGC TAC CAG ACG CCG TGG TCG GTC TGG	1244
Val Gln Glu Asn Arg Ala Asp Gly Tyr Gln Thr Pro Trp Ser Val Trp	
300 305 310	
CTC AAG AGC CAT CCG GCG AAG GCG TAAAGCGCA GGCGGCGGCT CGGAGTCAGG	1298
Leu Lys Ser His Pro Ala Lys Ala	
315	
GAAATGGCGC TGCCAGCACG GGACGCGCGG CGGCGGATGC TGCGCCGGTC AGGGATGATT	1358
GACGACAGCC AGAGAAGAAA GGATGGTTTT ATGAGGTGGA TCC	1401

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 346 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2

Met	Lys	Tyr	Trp	Gln	Lys	His	Ala	Val	Leu	Cys	Ser	Leu	Leu	Val	Gly
-27		-25					-20					-15			

Ala Ser Leu Trp Ile Leu Pro Gln Ala Asp Ala Ala Lys Ala Pro Glu
 -10 -5 1 5
 Gln Thr Val Thr Glu Pro Val Gly Ser Tyr Ala Arg Ala Glu Arg Pro
 10 15 20
 Gln Asp Phe Glu Gly Phe Val Trp Arg Leu Asp Asn Asp Gly Lys Glu
 25 30 35
 Ala Leu Pro Arg Asn Phe Arg Thr Ser Ala Asp Ala Leu Arg Ala Pro
 40 45 50
 Glu Lys Lys Phe His Leu Asp Ala Ala Tyr Val Pro Ser Arg Glu Gly
 55 60 65
 Met Asp Ala Leu His Ile Ser Gly Ser Ser Ala Phe Thr Pro Ala Gln
 70 75 80 85
 Leu Lys Asn Val Ala Ala Lys Leu Arg Glu Lys Thr Ala Gly Pro Ile
 90 95 100
 Tyr Asp Val Asp Leu Arg Gln Glu Ser His Gly Tyr Leu Asp Gly Ile
 105 110 115
 Pro Val Ser Trp Tyr Gly Glu Arg Asp Trp Ala Asn Leu Gly Lys Ser
 120 125 130
 Gln His Glu Ala Leu Ala Asp Glu Arg His Arg Leu His Ala Ala Leu
 135 140 145
 His Lys Thr Val Tyr Ile Ala Pro Leu Gly Lys His Lys Leu Pro Glu
 150 155 160 165
 Gly Gly Glu Val Arg Arg Val Gln Lys Val Gln Thr Glu Gln Glu Val
 170 175 180
 Ala Glu Ala Ala Gly Met Arg Tyr Phe Arg Ile Ala Ala Thr Asp His
 185 190 195
 Val Trp Pro Thr Pro Glu Asn Ile Asp Arg Phe Leu Ala Phe Tyr Arg
 200 205 210
 Thr Leu Pro Gln Asp Ala Trp Leu His Phe His Cys Glu Ala Gly Val
 215 220 225
 Gly Arg Thr Thr Ala Phe Met Val Met Thr Asp Met Leu Lys Asn Pro
 230 235 240 245
 Ser Val Ser Leu Lys Asp Ile Leu Tyr Arg Gln His Glu Ile Gly Gly
 250 255 260
 Phe Tyr Tyr Gly Glu Phe Pro Ile Lys Thr Lys Asp Lys Asp Ser Trp
 265 270 275

Lys Thr Lys Tyr Tyr Arg Glu Lys Ile Val Met Ile Glu Gln Phe Tyr
280 285 290

Arg Tyr Val Gln Glu Asn Arg Ala Asp Gly Tyr Gln Thr Pro Trp Ser
295 300 305

Val Trp Leu Lys Ser His Pro Ala Lys Ala
310 315

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "oligonucleotide SrPr6"
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Selenomonas ruminantium
 - (B) STRAIN: JY35
- (vii) IMMEDIATE SOURCE:
 - (A) LIBRARY: Genomic DNA library
 - (B) CLONE: pSrP.2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CGGGATGCTT CTGCCAGTAT

20

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "oligonucleotide SrPf6"
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: Selenomonas ruminantium

(B) STRAIN: JY35

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Genomic DNA library
- (B) CLONE: pSrP.2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CGTCCACGGA GTCACCCTAC

20

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide MATE2"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Selenomonas ruminantium
- (B) STRAIN: JY35

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Genomic DNA library
- (B) CLONE: pSrP.2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GCGAATTCAT GGCCAAGGCG CCGGAGCAGA C

31

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide MATE"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Selenomonas ruminantium*
- (B) STRAIN: JY35

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Genomic DNA library
- (B) CLONE: pSrP.2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GCGAATTCCG CAAGGCGCCG GAGCAGAC

28

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "oligonucleotide MATN"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Selenomonas ruminantium*
- (B) STRAIN: JY35

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: Genomic DNA library
- (B) CLONE: pSrP.2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GAGGATCCAT GGCCAAGGCG CCGGAGCAGA C

31

1 WE CLAIM:

- 2 1. A purified and isolated DNA encoding a phytase of a ruminal microorganism.
- 3
- 4 2. A purified and isolated DNA according to claim 1 wherein said ruminal
- 5 microorganism is a prokaryote.
- 6
- 7 3. A purified and isolated DNA according to claim 1 wherein said ruminal
- 8 microorganism is of the genus *Selenomonas*, *Prevotella*, *Treponema* or
- 9 *Megasphaera*.
- 10
- 11 4. A purified and isolated DNA according to claim 1 wherein said ruminal
- 12 microorganism is *Selenomonas ruminantium*, *Prevotella ruminicola*,
- 13 *Treponema bryantii* or *Megasphaera elsdenii*.
- 14
- 15 5. A purified and isolated DNA according to claim 1 wherein said ruminal
- 16 microorganism is *Selenomonas ruminantium*.
- 17
- 18 6. A purified and isolated DNA according to claim 1 wherein said ruminal
- 19 microorganism is *Selenomonas ruminantium* JY35 (ATCC 55785).
- 20
- 21 7. A purified and isolated DNA according to claim 1, said DNA being capable of
- 22 hybridizing under stringent conditions with a probe comprising at least 25
- 23 continuous nucleotides of nucleotide sequence SEQ ID NO. 1.
- 24
- 25 8. A purified and isolated DNA according to claim 1, said phytase comprising
- 26 amino acid sequence SEQ ID NO. 2.
- 27
- 28 9. A purified and isolated DNA according to claim 1, said DNA comprising
- 29 nucleotide sequence SEQ ID NO. 1.
- 30
- 31 10. A purified and isolated DNA according to claim 1, said DNA comprising
- 32 nucleotides 312-1268 of SEQ ID NO. 1.

- 1 11. A purified and isolated DNA according to claim 1, wherein said encoded
2 phytase has the following characteristics:
 - 3 a) a molecular mass of about 37 kDa;
 - 4 b) is active within a pH range of about 3.0 to 6.0; and
 - 5 c) is active within a temperature range of about 4 to 55°C.
- 6
- 7 12. A purified and isolated DNA according to claim 11 wherein said encoded
8 phytase is active within a temperature range of about 20 to 55°C.
- 9
- 10 13. A purified and isolated DNA according to claim 11 wherein said encoded
11 phytase is active within a temperature range of about 35 to 40°C.
- 12
- 13 14. A purified and isolated DNA according to claim 11, wherein the encoded
14 phytase has the following additional characteristic:
 - 15 d) a specific activity at least two fold higher than that of *Aspergillus ficuum*
16 NRRI 3135 PhyA as measured by the release of inorganic phosphate.
- 17
- 18 15. An expression construct capable of directing the expression of a phytase in
19 a suitable host cell, said expression construct comprising a DNA encoding a
20 phytase of a ruminal microorganism operably linked to control sequences
21 compatible with said host cell.
- 22
- 23 16. An expression construct according to claim 15 wherein said ruminal
24 microorganism is *Selenomonas ruminantium*.
- 25
- 26 17. An expression construct according to claim 15 wherein said encoded phytase
27 comprises amino acid sequence SEQ ID NO. 2.
- 28
- 29 18. A host cell transformed with a DNA encoding a phytase of a ruminal
30 microorganism so that the host cell can express the phytase encoded by said
31 DNA.
- 32

- 1 19. A transformed host cell according to claim 18 wherein said ruminal
2 microorganism is *Selenomonas ruminantium*.
3
- 4 20. A transformed host cell according to claim 18 wherein said encoded phytase
5 comprises amino acid sequence SEQ ID NO. 2.
6
- 7 21. A transformed host cell according to claim 18 wherein said host cell is a
8 eukaryote.
9
- 10 22. A transformed host cell according to claim 18 wherein said host cell is a
11 prokaryote.
12
- 13 23. A transformed host cell according to claim 18 wherein said host cell is a
14 *Pichea pastoris* cell.
15
- 16 24. A transformed host cell according to claim 18 wherein said host cell is a
17 *Bacillus subtilis* cell.
18
- 19 25. A transformed host cell according to claim 18 wherein said host cell is an *E.*
20 *coli* cell
21
- 22 26. *Selenomonas ruminantium* JY35 (ATCC 55785).
23
- 24 27. A transgenic plant transformed with a DNA encoding a phytase of a ruminal
25 microorganism so that the phytase encoded by said DNA can be expressed
26 by said plant.
27
- 28 28. A transgenic plant according to claim 27 wherein said ruminal microorganism
29 is *Selenomonas ruminantium*.
30
- 31 29. A transgenic plant according to claim 27 wherein said encoded phytase
32 comprises amino acid sequence SEQ ID NO. 2.

- 1 30. A phytase of a ruminal microorganism.
- 2
- 3 31. A phytase according to claim 30 wherein said ruminal microorganism is
- 4 *Selenomonas ruminantium*.
- 5
- 6 32. A phytase according to claim 30 wherein said phytase has the following
- 7 characteristics:
- 8 a) a molecular mass of about 37 kDa;
- 9 b) is active within a pH range of about 3.0 to 6.0; and
- 10 c) is active within a temperature range of about 4 to 55°C.
- 11
- 12 33. A phytase according to claim 32 having the following additional characteristic:
- 13 d) a specific activity at least two fold higher than that of *Aspergillus ficuum*
- 14 NRRI 3135 PhyA as measured by the release of inorganic phosphate.
- 15
- 16 34. A phytase according to claim 30, comprising a contiguous amino acid
- 17 sequence residing within amino acid sequence SEQ ID NO. 2.
- 18
- 19 35. A phytase according to claim 30 comprising amino acid sequence SEQ ID
- 20 NO. 2.
- 21
- 22 36. A feed composition comprising a feedstuff treated with a phytase of a ruminal
- 23 microorganism.
- 24
- 25 37. A feed composition according to claim 36 wherein said ruminal microorganism
- 26 is *Selenomonas ruminantium*.
- 27
- 28 38. A feed composition according to claim 36 wherein said phytase comprises
- 29 amino acid sequence SEQ ID NO. 2.
- 30

- 1 39. A feed composition according to claim 36 containing a sufficient amount of
2 said phytase to provide up to about 2000 Units (μ moles phosphate
3 released/minute) of phytase activity per kg feed composition.
4
- 5 40. A feed composition according to claim 36 containing a sufficient amount of
6 said phytase to provide up to about 1000 Units of phytase activity per kg feed
7 composition.
8
- 9 41. A feed composition according to claim 36 containing a sufficient amount of
10 said phytase to provide from about 50 to 800 Units of phytase activity per kg
11 feed composition.
12
- 13 42. A feed composition according to claim 36 containing a sufficient amount of
14 said phytase to provide from about 300 to 800 Units of phytase activity per kg
15 feed composition.
16
- 17 43. A feed additive comprising a preparation of lyophilized microorganisms, said
18 microorganisms expressing a phytase of a ruminal microorganism under
19 normal growing conditions.
20
- 21 44. A feed additive according to claim 43 wherein said microorganism is
22 *Selenomonas ruminantium*.
23
- 24 45. A feed additive according to claim 43 wherein said microorganism is a
25 recombinant microorganism transformed with a DNA encoding said phytase
26 of said ruminal microorganism.
27
- 28 46. A feed additive according to claim 45 wherein said ruminal microorganism is
29 *Selenomonas ruminantium*.
30
- 31 47. A feed additive according to claim 45 wherein said expressed phytase
32 comprises amino acid sequence SEQ ID NO. 2.

- 1 48. A feed additive for treatment of a feedstuff, said feed additive comprising a
2 phytase of a ruminal microorganism.
3
- 4 49. A feed additive according to claim 48 wherein said microorganism is
5 *Selenomonas ruminantium*.
6
- 7 50. A feed additive according to claim 48 wherein said phytase comprises amino
8 acid sequence SEQ ID NO. 2.
9
- 10 51. A method for producing a phytase, comprising:
11 (a) transforming at least one host cell with a DNA encoding a phytase of
12 a ruminal microorganism so that said host cell can express said phytase; and
13 (b) growing a culture of said host cells under conditions conducive to the
14 expression of said phytase by said host cells.
15
- 16 52. A method according to claim 51 comprising the further step of:
17 (c) extracting said phytase from said culture.
18
- 19 53. A method according to claim 51 wherein said ruminal microorganism is
20 *Selenomonas ruminantium*.
21
- 22 54. A method according to claim 51 wherein said phytase comprises amino acid
23 sequence SEQ ID NO. 2.
24
- 25 55. A method for producing a transgenic plant, comprising:
26 (a) transforming a plant with a DNA encoding a phytase of a ruminal
27 microorganism so that said plant can express said phytase; and
28 (b) growing said plant under conditions conducive to the expression of said
29 phytase by said plant.
30
- 31 56. A method according to claim 55 wherein said ruminal microorganism is
32 *Selenomonas ruminantium*.

- 1 57. A method according to claim 55 wherein said phytase comprises amino acid
2 sequence SEQ ID NO. 2:
3
- 4 58. A method for improving dietary phytate utilization by an animal, comprising
5 feeding said animal a diet which includes an effective amount of a phytase of
6 a ruminal microorganism.
7
- 8 59. A method according to claim 58 wherein said ruminal microorganism is
9 *Selenomonas ruminantium*.
10
- 11 60. A method according to claim 58 wherein said phytase comprises amino acid
12 sequence SEQ ID NO. 2.
13
- 14 61. A method according to claim 58 wherein said diet includes drinking water and
15 said phytase is included in said drinking water.
16
- 17 62. A method according to claim 58 wherein said phytase is provided in a mineral
18 block for consumption by said animal.
19
- 20 63. A method according to claim 58 wherein said phytase is provided in a pill for
21 consumption by said animal.
22
- 23 64. A method according to claim 58 wherein said phytase is provided in a gel
24 formulation for consumption by said animal.
25
- 26 65. A method according to claim 58 wherein said phytase is sprayed in a liquid
27 formulation onto a feedstuff for consumption by said animal.
28
- 29 66. A method according to claim 58 wherein said phytase is provided in a
30 pelletized feedstuff for consumption by said animal.
31

- 1 67. A method according to claim 58 wherein a feedstuff for consumption by said
2 animal is treated with a preparation of lyophilized microorganisms, said
3 microorganisms expressing said phytase under normal growing conditions.
4
- 5 68. A method for assaying phytase activity of a microorganism, comprising the
6 steps of:
7 (a) providing a growth medium upon which colonies of microorganisms
8 have been grown, said medium containing a source of phytate;
9 (b) contacting said medium with an aqueous solution of cobalt chloride;
10 and
11 (c) examining said medium for zones of clearing,
12
- 13 whereby false positive results caused by microbial acid production are
14 eliminated.
15
- 16 69. A method according to claim 68 wherein after step (b), said medium is
17 contacted with an aqueous solution of ammonium molybdate and an
18 aqueous solution of ammonium vanadate;
19
- 20 70. A method according to claim 68 wherein said medium is contacted with said
21 aqueous solution of cobalt chloride for at least about 5 minutes.
22
- 23 71. A method according to claim 68 wherein said medium is contacted with said
24 aqueous solutions of ammonium molybdate and ammonium vanadate for at
25 least about 5 minutes.
26
- 27 72. A method according to claim 68 wherein said medium is contacted with said
28 aqueous solutions of ammonium molybdate and ammonium vanadate
29 simultaneously.
30
- 31 73. A method according to claim 68 wherein the concentration of said aqueous
32 solution of cobalt chloride is about 2% (weight/volume).

- 1 74. A method according to claim 69 wherein the concentration of said aqueous
2 solution of ammonium molybdate is about 6% (weight/volume) and the
3 concentration of said aqueous solution of ammonium vanadate is about 0.5%
4 (weight/volume).
5
- 6 75. A method for identifying a nucleic acid molecule from an organism, said
7 nucleic acid molecule encoding a phytase, said method comprising the steps
8 of:
9
- 10 (a) isolating nucleic acid molecules from said organism;
11
- 12 (b) performing nucleic acid hybridization under conditions of moderate to
13 high stringency with said nucleic acid molecules and a labelled
14 hybridization probe having a nucleotide sequence comprising at least
15 25 continuous nucleotides of SEQ ID NO: 1.
16
- 17 76. A method according to claim 75 wherein said hybridization conditions are of
18 moderate stringency.
19
- 20 77. A method according to claim 75 wherein said hybridization conditions are of
21 high stringency.

1/10




Figure 1




Figure 2

Fig.3A



Fig. 3D




Fig.3B




Fig. 3E



Fig. 3C




Fig. 3F



3/10




Figure 4




Figure 5

4/10




Figure 6




Figure 7

5/10




Figure 8




Figure 9

6/10




Figure 10

7/10




Figure 11




Figure 12

8/10




Figure 13




Figure 14

9/10

Figure 15

1 CGTCCACGGA GTCACCCTAC TATACGACGT ATGTGAAGTT CACGTCGAAG TTCTAGGGAA 60
 61 TCACCGATTG GTGCAGGATT TTACCACTTC CTGTTGAAGC GGATGAGAAG GGGAAACGCG 120
 121 AAGCGGTGGA AGAGGTGCTG CACGACGGAC GATCGCGCTG AATGAATCAG TGCTTCCTAA 180
 181 CTATTGGGAT TCCGCGCAGA CGCGCGGATG GAGTAAAGGA GTAAGTTGTT ATG AAA TAC 239
 1 M K Y 3
 240 TGG CAG AAG CAT GCC GTT CTT TGT AGT CTC TTG GTC GGC GCA TCC CTC TGG 290
 4 W Q K H A V L C S L L V G A S L W 20
 291 ATA CTG CCG CAG GCC GAT GCG GCC AAG GCG CCG GAG CAG ACG GTG ACG GAG 341
 21 I L P Q A D A A K A P E O T V T E 37
 342 CCC GTT GGG AGC TAC GCG CGC GCG GAG CGG CCG CAG GAC TTC GAG GGC TTT 392
 38 P V G S Y A R A E R P Q D F E G F 54
 393 GTC TGG CGC CTC GAC AAC GAC GGC AAG GAG GCG TTG CCG CGT AAT TTC CGC 443
 55 V W R L D N D G K E A L P R N F R 71
 444 ACG TCG GCT GAC GCG CTG CGC GCG CCG GAG AAG AAA TTC CAT CTC GAC GCC 494
 72 T S A D A L R A P E K K F H L D A 88
 495 GCG TAT GTA CCG TCG CGC GAG GGC ATG GAT GCA CTC CAT ATC TCG GGC AGT 545
 89 A Y V P S R E G M D A L H I S G S 105
 546 TCC GCA TTC ACG CCG GCG CAG CTC AAG AAC GTT GCC GCG AAG CTG CGG GAG 596
 106 S A F T P A Q L K N V A A K L R E 122
 597 AAG ACG GCT GGC CCC ATC TAC GAT GTC GAC CTA CGG CAG GAG TCG CAC GGC 647
 123 K T A G P I Y D V D L R Q E S H G 139
 648 TAT CTC GAC GGT ATC CCC GTG AGC TGG TAC GGC GAG CGC GAC TGG GCA AAT 698
 140 Y L D G I P V S W Y G E R D W A N 156
 699 CTC GGC AAG AGC CAG CAT GAG GCG CTC GCC GAC GAG CGG CAC CGC TTG CAC 749
 157 L G K S Q H E A L A D E R H R L H 173
 750 GCA GCG CTC CAT AAG ACG GTC TAC ATC GCG CCG CTC GGC AAG CAC AAG CTC 800
 174 A A L H K T V Y I A P L G K H K L 190
 801 CCC GAG GGC GGC GAA GTC CGC CGC GTA CAG AAG GTG CAG ACG GAA CAG GAA 851
 191 P E G G E V R R V Q K V Q T E Q E 207
 852 GTC GCC GAG GCC GCG GGG ATG CGC TAT TTC CGC ATC GCG GCG ACG GAT CAT 902
 208 V A E A A G M R Y F R I A A T D H 224
 903 GTC TGG CCA ACG CCG GAG AAC ATC GAC CGC TTC CTC GCG TTT TAC CGC ACG 953
 225 V W P T P E N I D R F L A F Y R T 241
 954 CTG CCG CAG GAT GCG TGG CTC CAT TTC CAT TGT GAA GCC GGT GTC GGC CGC 1004
 242 L P Q D A W L H F H C E A G V G R 258
 1005 ACG ACG GCG TTC ATG GTC ATG ACG GAT ATG CTG AAG AAC CCG TCC GTA TCG 1055
 259 T T A F M V M T D M L K N P S V S 275
 1056 CTC AAG GAC ATC CTC TAT CGC CAG CAC GAG ATC GGC GGC TTT TAC TAC GGC 1106
 276 L K D I L Y R Q H E I G G F Y Y G 292

10/10

1107 GAG TTC CCC ATC AAG ACG AAG GAT AAA GAT AGC TGG AAG ACG AAA TAT TAT 1157
293 E F P I K T K D K D S W K T K Y Y 309

1158 AGG GAA AAG ATC GTG ATG ATC GAG CAG TTC TAC CGC TAT GTG CAG GAG AAC 1208
310 R E K I V M I E Q F Y R Y V Q E N 326

1209 CGC GCG GAT GGC TAC CAG ACG CCG TGG TCG GTC TGG CTC AAG AGC CAT CCG 1259
327 R A D G Y Q T P W S V W L K S H P 343

1260 GCG AAG GCG TAA AAGCGCAGGC GCGCGCTCGG AGTCAGGGAA ATGGCGCTGC 1311
344 A K A " 346

1312 CAGCACGGGA CGCGCGGCGG CGGATGCTGC GCGGTCAGG GATGATTGAC GACAGCCAGA 1371

1372 GAAGAAAGGA TGGTTTTATG AGGTGGATCC 1401

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/55, 9/16, 1/19, 1/20, 1/21, A01H 5/00, A23K 1/00, 1/165, C12Q 1/44, 1/68, C12N 15/82 // (C12N 1/19, C12R 1:84) (C12N 1/21, C12R 1:01, 1:125, 1:19)		A3	(11) International Publication Number: WO 97/48812 (43) International Publication Date: 24 December 1997 (24.12.97)
(21) International Application Number: PCT/CA97/00414 (22) International Filing Date: 13 June 1997 (13.06.97)		(74) Agent: McKAY-CAREY, Mary, Jane; McKay-Carey & Company, 10155 - 102nd Street, 2125 Commerce Place, Edmonton, Alberta T5J 4G8 (CA). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(30) Priority Data: 60/019,735 14 June 1996 (14.06.96) US 08/744,779 6 November 1996 (06.11.96) US 08/862,531 23 May 1997 (23.05.97) US		Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant (for all designated States except US): HER MAJESTY THE QUEEN IN RIGHT OF CANADA, represented by THE DEPARTMENT OF AGRICULTURE AND AGRI-FOOD CANADA [CA/CA]; Lethbridge Research Station, P.O. Box 3000, Main Lethbridge, Alberta T1J 4B1 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): CHENG, Kuo, Joan [CA/CA]; 2015 - 6th Avenue South, Lethbridge, Alberta T1J 1C2 (CA). SELINGER, Leonard, Brent [CA/CA]; 114 Laval Court West, Lethbridge, Alberta T1K 4G3 (CA). YANKE, Lindsey, Jay [CA/CA]; 1520 - 23rd Avenue N. #18, Lethbridge, Alberta T1H 4X9 (CA). BAE, Hee, Dong [KR/KR]; Eung-Sup Shim, 114-5, Chi-dong, Paldal-gu,		(88) Date of publication of the international search report: 26 March 1998 (26.03.98)	
(54) Title: DNA SEQUENCES ENCODING PHYTASES OF RUMINAL MICROORGANISMS			
(57) Abstract <p>Phytases derived from ruminal microorganisms are provided. The phytases are capable of catalyzing the release of inorganic phosphorus from phytic acid. Preferred sources of phytases include <i>Selenomonas</i>, <i>Prevotella</i>, <i>Treponema</i> and <i>Megasphaera</i>. A purified and isolated DNA encoding a phytase of <i>Selenomonas ruminantium</i> JY35 (ATCC 55785) is provided. Recombinant expression vectors containing DNAs encoding the phytases and host cells transformed with DNAs encoding the phytases are also provided. The phytases are useful in a wide range of applications involving the dephosphorylation of phytate, including, among other things, use in animal feed supplements.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

Patent Application No

PCT/CA 97/00414

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/55 C12N9/16 C12N1/19 C12N1/20 C12N1/21
 A01H5/00 A23K1/00 A23K1/165 C12Q1/44 C12Q1/68
 C12N15/82 //(C12N1/19,C12R1:84),(C12N1/21,C12R1:01,1:125,1:19)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H A23K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	EP 0 420 358 A (GIST BROCADES NV) 3 April 1991 see page 2, line 10-12 see page 4, line 47-55 see page 7; table 1 see page 11, line 30-50; example 2 see page 19; example 9 see page 28; claims 22,23,28-31 --- -/-	30 1,2,11, 12,14, 15,18, 21,22, 24,36, 39-42, 48,51, 52,58, 69,71, 72,74

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

22 January 1998

Date of mailing of the international search report

03.02.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Macchia, G

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/CA 97/00414

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	EP 0 449 375 A (GIST BROCADES NV ;MOGEN INT (NL)) 2 October 1991 see page 2, line 11-13 see page 3, line 45-54	30 1,27,36, 51,52, 55,58, 69,71, 72,74
X A	see page 7, line 28-55 see page 12, line 40-58; example 10 --- WO 94 03072 A (PANLABS INC ;NEVALAINEN HELENA K M (AU); ALKO LTD (FI); PALOHEIMO) 17 February 1994 see page 1, paragraph 2	30 43,45, 69,71, 72,74
X A A	see page 2, paragraph 1 see page 28, paragraph 1-2 see page 53, paragraph 6 --- PUNJ M.L. ET AL.: "Utilization of phytin phosphorus by rumen microorganisms" THE INDIAN VETERINARY JOURNAL, vol. 46, no. 10, 1969, pages 881-886, XP002044300 see page 885 --- WO 93 16175 A (GIST BROCADES NV) 19 August 1993 see page 5, line 16 - page 6, line 7 -----	30 12,13 69,71, 72,74

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA 97/00414

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/CA 97/00414

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-67 75-77

Isolated nucleic acid encoding a phytase of a ruminal microorganism, as in Seq.ID:1. Expression construct comprising it, transformed host cells and transgenic plants thereof. *Selenomonas ruminantium* JY35 (ATCC 55785). Phytase of a ruminal microorganism as in Seq.ID:2. Feed composition comprising feedstuff treated with said phytase, feed additive comprising microorganisms expressing said phytase. Method for producing said phytase. Method for producing a transgenic plant expressing said phytase. Method for improving dietary phytate utilization by an animal, via use of said phytase. Method to identify a nucleic acid encoding a phytase by means of at least part of Seq.ID:1.

2. Claims: 68-74

A method for assaying phytase activity of a microorganism.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/CA 97/00414

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0420358 A	03-04-91	AU 636673 B	06-05-93
		AU 6501190 A	28-04-91
		BG 60108 A	15-10-93
		CA 2042054 A	28-03-91
		CN 1051058 A	01-05-91
		DE 420358 T	12-10-95
		EP 0779037 A	18-06-97
		ES 2072834 T	01-08-95
		JP 4506007 T	22-10-92
		LT 1527 A,B	26-06-95
		LV 10310 A,B	20-10-94
		WO 9105053 A	18-04-91
		PL 167790 B	30-11-95
		PL 168470 B	29-02-96
		US 5436156 A	25-07-95
EP 0449375 A	02-10-91	AU 649447 B	26-05-94
		AU 7765691 A	21-10-91
		AU 632941 B	14-01-93
		AU 7776691 A	21-10-91
		CA 2056396 A	24-09-91
		EP 0449376 A	02-10-91
		IL 97645 A	18-03-97
		JP 6501838 T	03-03-94
		JP 6502296 T	17-03-94
		WO 9114782 A	03-10-91
		WO 9114772 A	03-10-91
		US 5543576 A	06-08-96
		US 5593963 A	14-01-97
WO 9403072 A	17-02-94	AU 4787993 A	03-03-94
		CA 2141431 A	17-02-94
		EP 0655890 A	07-06-95
		FI 950409 A	30-03-95
		JP 9505201 T	27-05-97
WO 9316175 A	19-08-93	AU 3628493 A	03-09-93
		EP 0626010 A	30-11-94
		FI 943707 A	10-08-94

THIS PAGE BLANK (USPTO)

PCTD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/04, 63/00, C12N 13/00	A1	(11) International Publication Number: WO 99/17610 (43) International Publication Date: 15 April 1999 (15.04.99)
(21) International Application Number: PCT/US98/20199 (22) International Filing Date: 24 September 1998 (24.09.98) (30) Priority Data: 08/942,939 2 October 1997 (02.10.97) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 12th floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US). (72) Inventors: ROTHMAN, Stephen, S.; 98 Acacia Avenue, Berkeley, CA 94708 (US). GOLDFINE, Ira, D.; 640 Goodhill Road, Kentfield, CA 94909 (US). GERMAN, Michael, S.; 1543 33rd Avenue, San Francisco, CA 94122 (US). (74) Agent: FRANCIS, Carol, L.; Bozicevic, Field & Francis LLP, Suite 200, 285 Hamilton Avenue, Palo Alto, CA 94301 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PROTEIN DELIVERY BY SECRETORY GLAND EXPRESSION (57) Abstract Secretory gland cells, particularly pancreatic, hepatic, and salivary gland cells, are genetically altered to operatively incorporate a gene which expresses a protein which has a desired therapeutic effect on a mammalian subject. The expressed protein is secreted directly into the bloodstream to obtain therapeutic levels of the protein thereby treating the patient in need of the protein. The transformed secretory gland cells provide long term or short term therapies for diseases associated with a deficiency in a particular protein or which are amenable to treatment by overexpression of a protein.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROTEIN DELIVERY BY SECRETORY GLAND EXPRESSION

Field of the Invention

This invention relates generally to the field of genetic transformation of cells *in vivo*, more particularly to *in vivo* transformation of secretory gland cells by introduction of the transforming nucleic acid into a secretory gland via a secretory gland duct.

Background of the Invention

The ability to replace defective or absent genes has attracted wide attention as a method to treat a variety of human diseases (Crystal 1995 *Science* 270:404), Lever et al. 1995 *Gene Therapy*, Pearson Professional, New York p. 1-91; Friedmann 1996 *Nature Med.* 2:144). Although originally intended as a means of correcting inherited disorders in certain populations of somatic cells, gene-based therapy can be a useful means to supply exogenous gene products to the circulatory system for the treatment of a wide range of systemic disorders that involve deficiencies in circulating proteins, such as hormones, growth factors, and clotting proteins (Lever et al. 1995 *supra*; Buckel 1996 *TIPS* 17:450), as well as a means of administering other polypeptide drugs. The success of this application depends upon developing effective methods to both manufacture the desired protein *in vivo* and then secrete it into blood (Crystal 1995 *supra*; Lever et al. 1995 *supra*).

Currently, DNA-based therapy (*i.e.*, gene therapy) is carried out in a variety of ways but involves two general protocols. In the first method, referred to as *ex vivo* gene therapy, cells are extracted from an individual and subjected to genetic manipulation. After genetic material has been properly inserted into the cells, the cells are implanted back into the individual from which they were removed. Persistent, *in vivo* expression of the newly implanted genetic material after transplantation of the transformed cells has been successful (see Morgan et al., *Science* 237:1476 (1987); and Gerrard et al., *Nat. Genet.* 3:180 (1993)). In the second approach to DNA-based therapy, referred to as *in vivo* gene therapy, cells within a living organism are transformed *in situ* with exogenous genetic material.

Several different methods for transforming cells can be used in accordance with either the *ex vivo* or *in vivo* transfection procedures. For example, various mechanical methods can be used to deliver the genetic material, including the use of fusogenic lipid vesicles (liposomes incorporating cationic lipids such as lipofection; see Felgner et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:7413-7417 (1987)); direct injection of DNA (Wolff, et al., *Science* (1990) 247:1465-1468); and pneumatic delivery of DNA-coated gold particles with a device referred to as the gene gun (Yang et al., *Proc. Natl. Acad. Sci. U.S.A.* 1990; 87:1568-9572). Morsy et al. reviews several of the different

techniques useful in transformation of cells *ex vivo* or *in vivo* and provides citations of numerous publications in each area (Morsy et al., *JAMA* 270:2338-2345 (1993)).

One method of particular interest for delivery of genetic material involves use of recombinant viruses to infect cells *in vivo* or *ex vivo*. In these methods, a virus containing the desired genetic material is allowed to infect target cells within the subject. Upon infection, the virus injects its genetic material into the target cells. The genetic material is then expressed within the target cell, providing for expression of the desired genetic material. However, it would be preferable to avoid introduction of the desired genetic material by viral infection for a number of reasons. For example, viral infection results in delivery of viral DNA in addition to the desired genetic material, which may in turn result in undesirable cellular effects such as, adverse immune reactions, productive viral replication, and adverse integration events.

There is a need in the field for a method for delivery of genetic material into a cell *in vivo* to provide for expression of the introduced polynucleotide and secretion of the gene product it encodes into the bloodstream. The present invention addresses this problem.

Summary of the Invention

Secretory gland cells are genetically altered to operatively incorporate a gene which is expressed by the genetically altered secretory gland cell to produce a polypeptide which is subsequently secreted into the bloodstream. Specifically, the invention involves introduction of a nucleotide of interest into a secretory gland via the duct system (*e.g.*, by retrograde ductal administration) to transform a secretory gland cell.

In one embodiment the invention features genetic alteration of cells of two secretory glands (*e.g.*, the liver and the pancreas).

In another embodiment, the invention features transformation of pancreatic cells with insulin-encoding nucleic acid to provide for expression and secretion of insulin at levels sufficient to maintain a substantially euglycemic state in a subject having a diabetic syndrome.

A primary object is to provide a non-invasive method of protein delivery (*i.e.*, the method involves introduction of the nucleic acid of interest from outside the body (*i.e.*, from the duct system of particular glands) wherein cells of a secretory gland, preferably the pancreas, salivary gland, or liver of a mammal are genetically modified to express a biologically active and therapeutically useful polypeptide, which polypeptide is secreted into the circulatory system of the individual.

Another object is to produce genetically transformed secretory gland cells which cells have incorporated into their genome genetic material which, when expressed, produces a biologically active and therapeutically useful protein which is secreted into the circulatory system.

An advantage of the present invention is that both long and short term therapy can be provided for diseases wherein individuals are suffering from the disease due to a deficiency in a particular protein, or by supplying an exogenous protein having a desired activity (e.g. antimicrobial activity).

5 These and other objects, advantages and features of the present invention will become apparent to those persons skilled in the art upon reading the details of the methodology and compositions as more fully set forth below.

Brief Description of the Drawing

10 Fig. 1 is a map of the pFGH construct, which contains the human growth hormone (hGH) genomic sequence.

Fig. 2 is a map of the pFGH.CMV construct, which contains the hGH genomic sequence operably linked to the CMV promoter.

15 Fig. 3 is a map of the pFGH.chymo construct, which contains the hGH genomic sequence operably linked to the chymotrypsin B promoter.

Fig. 4 is a graph showing the levels of tissue expression of hGH expression in the pancreas of rats after retrograde injection with either a control containing no DNA or a test sample containing a hGH construct.

20 Fig. 5 is a graph showing the serum levels of hGH in rats after retrograde pancreatic injection with either a control containing no DNA or a test sample containing a human growth hormone construct.

Fig. 6 is a graph showing the correlation between pancreatic tissue expression and serum levels of hGH.

25 Fig. 7 is a graph showing tissue expression of hGH following transformation of salivary gland cells by intraductal injection.

Fig. 8 is a graph showing regulation of plasma levels of hGH expressed from transformed salivary gland cells.

Fig. 9 is a graph showing regulation of plasma levels of hGH expressed from transformed pancreatic cells.

30 Fig. 10 is a map of the pBAT16.hInsG1.M2 construct, which contains DNA encoding an altered form of human insulin.

Fig. 11 is a graph showing the glucose response in streptozotocin-treated rats having pancreatic cells transformed with either human insulin (open bars) or green fluorescent protein (GFP; striped bars).

Fig. 12, is a graph showing the blood glucose levels in control rats (mock-treated: closed squares), streptozotocin-treated rats (open squares), and streptozotocin rats treated by transformation of pancreatic cells with DNA encoding human insulin (closed circles).

Fig. 13 is a graph showing expression of hGH in the plasma of control rats (no DNA) and of rats in which hGH-encoding DNA was introduced into the liver by intraductal injection.

Fig. 14 is a graph showing the relative amounts of hGH in the pancreatic tissue of rats that received either pFGH (control), pFGH.chymo, pFGH.RSV, pFGH.RSV, or pFGH.CMV by intraductal administration to the pancreas.

Fig. 15 is a graph showing the relative amounts of hGH in the pancreatic tissue of rats that received either no DNA (mock-transformed), pFGH.CMV, pFGH.CMV premixed with lipofectin, or pFGH.CMV premixed with adenovirus.

Fig. 16 is a graph showing the relative levels of plasma hGH in rats that received either pFGH (control), pFGH.chymo, pFGH.RSV, pFGH.RSV, or pFGH.CMV by intraductal administration to the pancreas.

Fig. 17 is a graph showing the relative amounts of plasma hGH in rats that received either no DNA (mock-transformed), pFGH.CMV, pFGH.CMV premixed with lipofectin, or pFGH.CMV premixed with adenovirus by intraductal administration to the pancreas..

Fig. 18 is a graph showing the relative levels of plasma hGH in rats that received no DNA (control), received hGH-encoding DNA via intraductal delivery to the liver, received hGH-encoding DNA via intraductal delivery to the pancreas, or received hGH-encoding DNA via intraductal delivery to both the liver and pancreas.

Fig. 19 is a graph showing the relative levels of hGH expression in pancreas tissue following administration of DNA to both pancreas and liver or to pancreas alone. The graph shows tissue levels of hGH after administration of a control (no DNA) to both pancreas and liver (left-most bar); administration of pFGH.CMV to both pancreas and liver (center bar); and pFGH.CMV to pancreas alone (right-most bar). Adenovirus was admixed with the construct as an adjuvant.

Fig. 20 is graph showing the relative levels of hGH expression in salivary gland tissue in rats that received either no DNA (control rats), pFGH.CMV, pFGH.CMV premixed with lipofectin, or pFGH.CMV premixed with adenovirus.

Fig. 21 is a graph showing stimulation of hGH secretion into the plasma of rats that received hGH-encoding DNA by intraductal injection into both the pancreas and liver.

Fig. 22 is a graph showing the blood glucose levels of streptozotocin-treated rats (diabetic) that received either no DNA (open squares) or received human insulin-encoding DNA by intraductal injection into the pancreas (closed squares) over a three day period.

Fig. 23 is a graph showing the plasma insulin levels of streptozotocin-treated rats (diabetic) that received either no DNA (open squares) or received human insulin-encoding DNA by intraductal injection into the pancreas (closed squares) over a three day period.

Fig. 24 is a graph showing the blood glucose levels (over a six day period) of streptozotocin-treated rats (diabetic) that received either no DNA (open squares) or received human insulin-encoding DNA by intraductal injection into the pancreas (closed squares).

Description of the Preferred Embodiments

Before the present method of genetically transforming secretory gland cells and methods for protein delivery are described, it is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, secretory glands, vectors and reagents described as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a secretory gland cell" includes a plurality of such cells and reference to "the transformation vector" includes reference to one or more transformation vectors and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs. Although any methods, devices and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, the preferred methods, devices and materials are now described.

All publications mentioned herein are incorporated herein by reference for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are described in the publications which might be used in connection with the presently described invention. The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such a disclosure by virtue of prior invention.

Definitions

By "secretory gland" is meant an aggregation of cells specialized to secrete or excrete materials not related to their ordinary metabolic needs. Secretory glands include salivary glands,

pancreas, mammary glands, thyroid gland, thymus gland, pituitary gland, liver, and other glands well known in the art.

By "exocrine gland" is meant a ducted gland or portion of a ducted gland that releases its products externally relative to the body, *e.g.*, either into the internal cavities such as the ocular and nasal cavities, the lumen of the gastrointestinal tract, or onto the surface of the body.

By "salivary gland" is meant a gland of the oral cavity which secretes saliva, including the glandulae salivariae majores of the oral cavity (the parotid, sublingual, and submandibular glands) and the glandulae salivariae minores of the tongue, lips, cheeks, and palate (labial, buccal, molar, palatine, lingual, and anterior lingual glands).

By "pancreas" is meant a large, elongated, racemose gland situated transversely behind the stomach, between the spleen and the duodenum. The pancreas is composed of an endocrine portion (the pars endocrina) and an exocrine portion (the pars exocrina). The pars endocrina, which contains the islets of Langerhans, produces and secretes proteins, including insulin, directly into the bloodstream. The pars exocrina contains secretory units and produces and secretes a pancreatic juice, which contains enzymes essential to protein digestion, into the duodenum.

By "retrograde ductal injection" is meant the administration of a liquid or other material into the fluid contents of the duct system of an exocrine gland in a direction opposite to the normal flow of that fluid, either at the external orifice of the duct system or through its wall. "Retrograde ductal injection" can be a single, discontinuous administration or continuous administration (*i.e.*, perfusion).

By "transformation" is meant a genetic change induced in a cell following incorporation of new DNA (*i.e.*, DNA exogenous to the cell). Where the cell is a mammalian cell, the genetic change may be achieved by introduction of the DNA into the genome of the cell.

By "transfection" is meant the transformation of a cell with DNA from a virus.

By "transformed cell" is meant a cell into which (or, where the introduced DNA is incorporated into the genome, into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA molecule encoding a protein of interest.

By "nucleic acid of interest" is meant any DNA or RNA molecule which encodes a polypeptide or other molecule which is desirable for administration to a mammalian subject for expression of the product encoded by the nucleic acid of interest and delivery of the encoded product into the blood stream of the mammalian subject. The nucleic acid is generally operatively linked to other sequences which are needed for its expression such as a promoter. The term "DNA of interest" is used as shorthand herein to refer to the nucleic acid of interest.

By "construct" is meant a nucleic acid molecule which contains the nucleic acid of interest (*e.g.*, the DNA of interest), generally operably linked to a promoter for expression of the polypeptide

encoded by the nucleic acid of interest. "Constructs" as used herein is generally meant to refer a nucleic acid molecule that facilitates expression of a polypeptide encoded by the nucleic acid to be introduced into a secretory gland cell.

By "vector" is meant any compound, biological or chemical, that facilitates transformation
5 of a secretory gland cell with a DNA of interest. Exemplary biological vectors include viruses, particularly attenuated and/or replication-deficient viruses. Exemplary chemical vectors include lipid complexes and naked DNA constructs.

By "naked DNA" or "naked nucleic acid" or DNA sequence and the like is meant a nucleic acid molecule that is not contained within a viral particle, bacterial cell or other encapsulating means
10 that facilitates delivery of nucleic acid into the cytoplasm of the target cell. Naked nucleic acid can be associated with means for facilitating delivery of the nucleic acid to the site of the target cell (*e.g.*, means that facilitate travel into the target cell of the nucleic acid through the alimentary canal, protect the nucleic acid from stomach acid, and/or serve to penetrate intestinal mucus) and/or to the surface of the target epithelial cell.

By "promoter" is meant a minimal DNA sequence sufficient to direct transcription.
15 "Promoter" is also meant to encompass those promoter elements sufficient for promoter-dependent gene expression controllable for cell-type specific, tissue-specific or inducible by external signals or agents; such elements may be located in the 5' or 3' regions of the native gene.

By "secretory gland specific promoter" is meant a promoter which directs expression of an
20 operably linked DNA sequence when bound by transcriptional activator proteins, or other regulators of transcription, which are unique to a specific type of secretory gland cell. For example, by "salivary gland specific promoter" is meant a secretory gland specific promoter which directs expression in a salivary gland cell. A salivary amylase promoter is an example of a salivary gland specific promoter. By "pancreas specific promoter" is meant a secretory gland specific promoter
25 which directs expression in a pancreatic cell. Examples of pancreas specific promoters include a pancreatic amylase promoter and an insulin promoter.

By "operably linked" is meant that a DNA sequence and a regulatory sequence(s) are connected in such a way as to permit gene expression when the appropriate molecules (*e.g.*, transcriptional activator proteins) are bound to the regulatory sequence(s).

By "operatively inserted" is meant that the DNA of interest introduced into the cell is
30 positioned adjacent a DNA sequence which directs transcription and translation of the introduced DNA (*i.e.*, facilitates the production of, *e.g.*, a polypeptide encoded by a DNA of interest).

By "mammalian subject" or "mammalian patient" is meant any mammal to which intravenous protein delivery is desired, including human, bovine, equine, canine, and feline subjects.

By "euglycemia" or "euglycemic state" is meant a state associated with a level of blood glucose that is normal or nearly normal, particularly relative to the levels of blood glucose in a subject having a disease or condition associated with hyperglycemia. In humans, euglycemia correlates with blood glucose levels in the range of 70 mg/dl to 130 mg/dl.

5 The terms "synergistic," "synergistic effect," and the like are used herein to describe improved effects (e.g., an increase in tissue expression levels in one or more secretory glands, an increased responsiveness to hormonal stimulation to elicit secretion of a polypeptide of interest, or a decrease in an undesirable phenotype) by combining one or more aspects of the invention (e.g., by transformation of more than one secretory gland in a single subject, or by transformation of a
10 secretory gland(s) with multiple constructs encoding the same or different polypeptides).

Overview of the Invention

The present invention features methods for genetically altering a secretory gland cell (*i.e.*, secretory gland cell transformation) and methods of delivering a protein using the methods of
15 genetically altering secretory gland cells. More specifically the invention features methods for delivery of a protein or other product encoded by a nucleic acid sequence of interest to a mammalian subject by expression of a DNA of interest in cells within a secretory gland of a mammalian patient (*i.e.*, by *in vivo* gene therapy). Preferably, the transformed secretory gland cells expressing the protein encoded by the DNA of interest secrete a therapeutically effective amount of the protein into
20 the bloodstream of the mammalian patient. Preferably, the secretory gland into which the DNA of interest is introduced and expressed is the pancreas, a salivary gland, or the liver. In short, the invention features a delivery system that involves introduction of a nucleic acid sequence encoding a product of interest (e.g., a protein) into a secretory gland cell (e.g., a salivary gland cell, hepatocyte, or pancreatic cell, particularly exocrine cells of salivary gland, liver, or pancreas), expression of the
25 encoded protein, and delivery of the protein into the blood stream by secretion of the protein by the transformed secretory gland cell.

The present invention preferably uses either naked DNA or DNA premixed with adjuvants (e.g., lipofectin or viral particles). It is not necessary to incorporate the DNA into viral particles in order to achieve transformation of secretory gland cells and provide expression of the polypeptide of
30 interest at physiologic/therapeutic levels in the bloodstream.

An important feature of the invention is the use of exocrine cells of glands of the gastrointestinal tract (*i.e.*, pancreas, liver, salivary gland) to produce and secrete therapeutic proteins into blood. While it is well understood that exocrine cells secrete into the lumen of the glands' ducts (*i.e.* in an exocrine direction), with the exception of the liver (*i.e.*, the hepatocytes secrete cellular
35 products in both directions, e.g. blood proteins into blood and bile salts into the intestinal lumen), it

is not widely appreciated that exocrine cells can also secrete significant amounts of protein into the systemic circulation. For example, exocrine proteins such as α -amylase (salivary glands), pepsinogen (gastric glands), various digestive enzymes from the exocrine pancreas, salivary gland kallikreins and nerve growth factor (Liebow, 1988 Pancreas 3:343-351) are normal constituents of blood. In the pancreas, substantial quantities of digestive enzymes are released into the circulation (Saito et al., 1973 Jpn. J. Physiol. 23:477-95; Isenman et al. 1997 Proc. Natl. Acad. Sci (USA) 74:4068-4072; Papp et al. 1980 Acta Physiol. Acad. Sci. Hung. 56:401-410; Geokas et al., 1980 Am. J. Physiol. 238:238-246; Miyasaka et al. 1981 Am. J. Physiol. 241:170-175; Grendell et al. 1982 Am. J. Physiol. 243:54-59). Endocrine secretion can be greatly enhanced by common secretory stimulants (Saito et al., *supra*; Isenman et al. *supra*; Miyasaka et al. *supra*; Grendell et al. *supra*). As much as 20-25% of the total secreted product can be released into blood as a consequence of stimulation (Grendell et al. *supra*). The present invention takes advantage of the discovery that exocrine gland cells can be transformed with a desired DNA sequence and secrete the encoded polypeptide into the bloodstream rather than only or primarily into the gastrointestinal tract.

In addition to the advantages described above, the invention also permits access to the cells of secretory glands without invasive procedures. For example, it is possible to cannulate either the collecting duct of a major salivary gland through its orifice in the mouth, or the common bile or pancreatic duct by means of endoscopic retrograde cholangiopancreatography (ERCP). These are common diagnostic procedures performed on awake patients. The non-invasive methods of the invention allow delivery of the DNA of interest in a safe manner that substantially avoids the inflammatory and immunological responses associated with other means of DNA delivery.

The invention also takes advantage of the protein-producing capacity of secretory gland cells. This advantage is particularly useful for the production of hormones such as hGH and insulin, which have short half-lives in blood and are cleared quickly. The cells of the exocrine glands are the body's major protein synthesizing and secreting systems. For example, the human exocrine pancreas manufactures and secretes approximately 20 g of protein daily. According to the present invention, even a small proportion of protein synthesized by secretory glands provides enough secreted product to provide therapeutic protein levels for the treatment of most diseases of circulating proteins.

The invention will now be described in further detail.

Constructs

Any nucleic acid construct having a eukaryotic promoter operably linked to a DNA of interest can be used in the invention. The constructs containing the DNA sequence (or the corresponding RNA sequence) which may be used in accordance with the invention may be any eukaryotic expression construct containing the DNA or the RNA sequence of interest. For example,

a plasmid or viral construct (e.g. adenovirus) can be cleaved to provide linear DNA having ligatable termini. These termini are bound to exogenous DNA having complementary, like ligatable termini to provide a biologically functional recombinant DNA molecule having an intact replicon and a desired phenotypic property. Preferably the construct is capable of replication in both eukaryotic and prokaryotic hosts, which constructs are known in the art and are commercially available.

The exogenous (*i.e.*, donor) DNA used in the invention is obtained from suitable cells, and the constructs prepared using techniques well known in the art. Likewise, techniques for obtaining expression of exogenous DNA or RNA sequences in a genetically altered host cell are known in the art (see, for example, Kormal et al., *Proc. Natl. Acad. Sci. USA*, **84**:2150-2154, 1987; Sambrook et al., Molecular Cloning: a Laboratory Manual, 2nd Ed., 1989, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; each of which are hereby incorporated by reference with respect to methods and compositions for eukaryotic expression of a DNA of interest).

Preferably, the DNA construct contains a promoter to facilitate expression of the DNA of interest within a secretory gland cell. Preferably the promoter is a strong, eukaryotic promoter such as a promoter from cytomegalovirus (CMV), mouse mammary tumor virus (MMTV), Rous sarcoma virus (RSV), or adenovirus. More specifically, exemplary promoters include the promoter from the immediate early gene of human CMV (Boshart et al., *Cell* **41**:521-530, 1985) and the promoter from the long terminal repeat (LTR) of RSV (Gorman et al., *Proc. Natl. Acad. Sci. USA* **79**:6777-6781, 1982). Of these two promoters, the CMV promoter is preferred as it provides for higher levels of expression than the RSV promoter.

Alternatively, the promoter used may be a tissue-specific promoter. For example, where the secretory gland is the pancreas, the promoter used in the vector is preferably a pancreas specific promoter, *e.g.*, an insulin promoter or a pancreas α -amylase promoter; where the secretory gland is a salivary gland, the tissue-specific promoter may be a salivary α -amylase promoter or mumps viral gene promoter. Both pancreatic and salivary α -amylase genes have been identified and characterized in both mice and humans (see, for example, Jones et al., *Nucleic Acids Res.*, **17**:6613-6623; Pittet et al., *J. Mol. Biol.*, **182**:359-365, 1985; Hagenbuchle et al., *J. Mol. Biol.*, **185**:285-293, 1985; Schibler et al., *Oxf. Surv. Eukaryot. Genes*, **3**:210-234, 1986; and Sierra et al., *Mol. Cell. Biol.*, **6**:4067-4076, 1986 for murine pancreatic and salivary α -amylase genes and promoters; Samuelson et al., *Nucleic Acids Res.*, **16**:8261-8276, 1988; Groot et al., *Genomics*, **5**:29-42, 1989; and Tomita et al., *Gene*, **76**:11-18, 1989 for human pancreatic and salivary α -amylase genes and their promoters; Ting et al., *Genes Dev.* **6**:1457-65, 1992 for human salivary α -amylase AMY1C promoter sequences).

The constructs of the invention may also include sequences in addition to promoters which enhance secretory gland specific expression. For example, where pancreas specific expression of the

DNA of interest is desired, the construct may include a PTF-1 recognition sequence (Cockell et al., *Mol. Cell. Biol.*, 9:2464-2476, 1989). Sequences which enhance salivary gland specific expression are also well known in the art (see, for example, Robins et al., *Genetica* 86:191-201, 1992).

5 Other components such as a marker (*e.g.*, an antibiotic resistance gene (such as an ampicillin resistance gene) or β -galactosidase) to aid in selection of cells containing and/or expressing the construct, an origin of replication for stable replication of the construct in a bacterial cell (preferably, a high copy number origin of replication), a nuclear localization signal, or other elements which facilitate production of the DNA construct, the protein encoded thereby, or both.

10 For eukaryotic expression, the construct should contain at a minimum a eukaryotic promoter operably linked to a DNA of interest, which is in turn operably linked to a polyadenylation sequence. The polyadenylation signal sequence may be selected from any of a variety of polyadenylation signal sequences known in the art. Preferably, the polyadenylation signal sequence is the SV40 early polyadenylation signal sequence. The construct may also include one or more introns, which can increase levels of expression of the DNA of interest, particularly where the DNA
15 of interest is a cDNA (*e.g.*, contains no introns of the naturally-occurring sequence). Any of a variety of introns known in the art may be used (*e.g.*, the human β -globin intron, which is inserted in the construct at a position 5' to the DNA of interest).

The DNA of interest may be inserted into a construct so that the therapeutic protein is expressed as a fusion protein (*e.g.*, a fusion protein having β -galactosidase or a portion thereof at the
20 N-terminus and the therapeutic protein at the C-terminal portion). Production of a fusion protein can facilitate identification of transformed cells expressing the protein (*e.g.*, by enzyme-linked immunosorbent assay (ELISA) using an antibody which binds to the fusion protein).

The Nucleic Acid (DNA) of Interest

25 The DNA of interest can be any DNA encoding any protein for which intravenous therapy is desirable. For example, intravenous protein therapy is appropriate in treating a mammalian subject having an inherited or acquired disease associated with a specific protein deficiency (*e.g.*, diabetes, hemophilia, anemia, severe combined immunodeficiency). Such protein deficient states are amenable to treatment by replacement therapy, *i.e.*, expression of a protein to restore the normal bloodstream
30 levels of the protein to at least normal levels.

Alternatively, the DNA of interest may encode a polypeptide that is either normally present in a healthy mammalian subject or which is foreign to the mammalian subject, and which polypeptide is effective in treatment of a condition by expression or over-expression of the polypeptide. For example, the DNA of interest can encode antimicrobial, antiparasitic, antifungal, or
35 antiviral polypeptides for treatment of a mammalian subject having a viral (*e.g.*, human

immunodeficiency virus (HIV), Epstein-Barr virus (EBV), herpes simplex virus (HSV), bacterial, fungal, and/or parasitic infection, particularly where the infection is chronic, *i.e.*, persisting over a relatively long period of time. The methods of the invention may also be used to enhance expression of a protein present in a normal mammal, or to express a protein not normally present in a normal mammal, in order to achieve a desired effect (*e.g.*, to enhance a normal metabolic process). For example, a secretory gland of a dairy cow may be transformed with DNA encoding bovine growth hormone (BGH) in order to enhance levels of BGH in the bloodstream and enhance milk production.

The DNA of interest is preferably obtained from a source of the same species as the mammalian subject to be treated (*e.g.* human to human), but this is not an absolute requirement. DNA obtained from a species different from the mammalian subject can also be used, particularly where the amino acid sequences of the proteins are highly conserved and the xenogeneic protein is not highly immunogenic so as to elicit a significant, undesirable antibody response against the protein in the mammalian host.

Exemplary, preferred DNAs of interest include DNA encoding insulin, growth factors (*e.g.*, growth hormone, insulin-like growth factor-1 (IGF-I), platelet-derived growth factor (PDGF), epidermal growth factor (EGF), acidic fibroblast growth factor, basic fibroblast growth factor, or transforming growth factor β), cytokines (*e.g.*, interferon (INF) (*e.g.*, INF- α 2b, INF- α 2a, INF- α N1, INF- β 1b, INF- γ), interleukin (*e.g.* IL-2, IL-8), or tumor necrosis factor (TNF) (*e.g.* TNF- α , TNF- β)), clotting factors (*e.g.*, clotting factor VIII), hormones (*e.g.* GP-1), antimicrobial polypeptides (*e.g.*, antibacterial, antifungal, antiviral, and/or antiparasitic polypeptides), enzymes (*e.g.*, adenosine deaminase), filgrastim (Neupogen), hemoglobin, erythropoietin, insulinotropin, imiglucrase, sarbramostim, antigens, tissue plasminogen activator (tPA), urokinase, streptokinase, endothelial, soluble CD4, and antibodies and/or antigen-binding fragments (*e.g.* FAbs) thereof (*e.g.*, orthoclone OKT-c (anti-CD3), GPIIb/IIIa monoclonal antibody). Preferably, the mammalian subject is a human subject and the DNA expressed encodes a human protein.

Table 1 provides a list of exemplary proteins and protein classes which can be delivered to the bloodstream of a mammalian subject via the method of secretory gland cell transformation of the invention.

TABLE 1: Exemplary Proteins and Protein Classes for Expression in and Secretion by Secretory Gland Cells

SPECIFIC EXEMPLARY PROTEINS	
insulin	interferon- α 2B
human growth hormone (hGH)	transforming growth factor (TGF)
erythropoietin (EPO)	ciliary neurite transforming factor (CNTF)
clotting factor VIII	insulin-like growth factor-1 (IGF-1)
bovine growth hormone (BGH)	granulocyte macrophage colony stimulating factor (GM-CSF)
platelet derived growth factor (PDGF)	interferon- α 2A
clotting factor VIII	brain-derived neurite factor (BDNF)
thrombopoietin (TPO)	insulintropin
IL-1	tissue plasminogen activator (tPA)
IL-2	urokinase
IL-1 RA	streptokinase
superoxide dismutase (SOD)	adenosine deaminase
catalase	calcitonin
fibroblast growth factor (FGF) (acidic or basic)	arginase
neurite growth factor (NGF)	phenylalanine ammonia lyase
granulocyte colony stimulating factor (G-CSF)	γ -interferon
L-asparaginase	pepsin
uricase	trypsin
chymotrypsin	elastase
carboxypeptidase	lactase
sucrase	intrinsic factor
calcitonin	parathyroid hormone(PTH)-like hormone
Ob gene product	cholecystokinin (CCK)
glucagon	insulinotrophic hormone

EXEMPLARY CLASSES OF PROTEINS	
enzymes (e.g., proteases, phospholipases, etc.)	pituitary hormones
protease inhibitors	growth factors
cytokines	somatomedians
chemokines	immunoglobulins
gonadotrophins	interleukins
chemotactins	interferons
lipid-binding proteins	

Numerous proteins that are desirable for intravenous protein therapy are well known in the art and the DNA encoding these proteins has been isolated. For example, the sequence of the DNAs encoding insulin, human growth hormone, intrinsic factor, clotting factor VIII, and erythropoietin are available from Genbank and/or have been described in the scientific literature (e.g., human clotting factor VIII gene: Gitschier et al., *Nature* 312:326-330, 1984; Wood et al., *Nature* 312:330-337, 1984; human intrinsic factor: Hewitt et al., *Genomics* 10:432-440, 1991). Moreover, proteins commonly used in treatments can be used in the procedures of the present invention. Such proteins are disclosed in, for example, the Physicians' Desk Reference (1994 Physicians' Desk Reference,

48th Ed., Medical Economics Data Production Co., Montvale, NJ; incorporated by reference) and can be dosed using methods described in Harrison's Principles of Internal Medicine and/or the AMA "Drug Evaluations Annual" 1993, all incorporated by reference.

Where the DNA encoding a protein of interest has not been isolated, this can be accomplished by various, standard protocols well known to those of skill in the art (see, for example, Sambrook et al., *ibid*; Suggs et al., *Proc. Natl. Acad. Sci. USA* **78**:6613-6617, 1981; USPN 4,394,443; each of which are incorporated herein by reference with respect to identification and isolation of DNA encoding a protein of interest). For example, genomic or cDNA clones encoding a specific protein can be isolated from genomic or cDNA libraries using hybridization probes designed on the basis of the nucleotide or amino acid sequences for the desired gene. The probes can be constructed by chemical synthesis or by polymerase chain reaction (PCR) using primers based upon sequence data to amplify DNA fragments from pools or libraries (USPNs 4,683,195 and 4,683,202). Nucleotide substitutions, deletions, additions, and the like can also be incorporated into the polynucleotides, so long as the ability of the polynucleotide to hybridize is not substantially disrupted. (Sambrook et al. *ibid*). The clones may be expressed or the DNA of interest can be excised or synthesized for use in other constructs. If desired, the DNA of interest can be sequenced using methods well known in the art.

It may also be desirable to produce altered forms of the therapeutic proteins that are, for example, protease resistant or have enhanced activity relative to the wild-type protein. For example, where the therapeutic protein is a hormone, it may be desirable to alter the protein's ability to form dimers or multimeric complexes. For example, insulin may be modified so as to prevent its dimerization has a more rapid onset of action relative to wild-type, dimerized insulin.

Vectors for Delivery of the DNA of Interest to the Secretory Gland Cell

The vectors for delivery of the DNA of interest can be either viral or non-viral, or may be composed of naked DNA admixed with an adjuvant such as viral particles (e.g., adenovirus) or cationic lipids or liposomes. An "adjuvant" is a substance that does not by itself produce the desired effect, but acts to enhance or otherwise improve the action of the active compound. The precise vector and vector formulation used will depend upon several factors such as the secretory gland targeted for gene transfer.

Non-viral vectors

The DNA of interest may be administered using a non-viral vector. "Non-viral vector" as used herein is meant to include naked DNA, chemical formulations containing naked DNA (e.g., a formulation of DNA and cationic compounds (e.g., dextran sulfate)), and naked DNA mixed with an adjuvant such as a viral particle (*i.e.*, the DNA of interest is not contained within the viral particle,

but the transforming formulation is composed of both naked DNA and viral particles (*e.g.*, adenovirus particles) (see, *e.g.*, Curiel et al. 1992 Am. J. Respir. Cell Mol. Biol. 6:247-52). Thus "non-viral vector" can include vectors composed of DNA plus viral particles where the viral particles do not contain the DNA of interest within the viral genome.

5 In one preferred embodiment, the formulation comprises viral particles which are mixed with the naked DNA construct prior to administration. Preferably, about 10^8 to 10^{10} viral particles (preferably about 1×10^{10} to 5×10^{10} , more preferably about 3×10^{10} particles) are mixed with the naked DNA construct (about 5 μ g to 50 μ g DNA, more preferably about 8 μ g to 25 μ g DNA) in a total volume of about 100 μ l. Preferably the viral particles are adenovirus particles (Curiel et al.,
10 1992 *supra*).

Alternatively or in addition, the DNA of interest can be complexed with polycationic substances such as poly-L-lysine or DEAC-dextran, targeting ligands, and/or DNA binding proteins (*e.g.*, histones). DNA- or RNA-liposome complex formulations comprise a mixture of lipids which bind to genetic material (DNA or RNA) and facilitate delivery of the nucleic acid into the cell.
15 Liposomes which can be used in accordance with the invention include DOPE (dioleoyl phosphatidyl ethanol amine), CUDMEDA (N-(5-cholestrum-3- β -ol 3-urethanyl)-N',N'-dimethylethylene diamine).

For example, the naked DNA can be administered in a solution containing Lipofectin™ (LTI/BRL) at a concentrations ranging from about 2.5 % to 15% volume:volume, preferably about
20 6% to 12% volume:volume. Preferred methods and compositions for formulation of DNA for delivery according to the method of the invention are described in USPN 5,527,928, incorporated herein by reference.

The DNA of interest can also be administered as a chemical formulation of DNA or RNA coupled to a carrier molecule (*e.g.*, an antibody or a receptor ligand) which facilitates delivery to
25 host cells for the purpose of altering the biological properties of the host cells. By the term "chemical formulations" is meant modifications of nucleic acids to allow coupling of the nucleic acid compounds to a carrier molecule such as a protein or lipid, or derivative thereof. Exemplary protein carrier molecules include antibodies specific to the cells of a targeted secretory gland or receptor ligands, *i.e.*, molecules capable of interacting with receptors associated with a cell of a
30 targeted secretory gland.

Viral vectors

In general, viral vectors used in accordance with the invention are composed of a viral particle derived from a naturally-occurring virus which has been genetically altered to render the virus replication-defective and to express a recombinant gene of interest in accordance with the
35 invention. Once the virus delivers its genetic material to a cell, it does not generate additional

infectious virus but does introduce exogenous recombinant genes into the cell, preferably into the genome of the cell.

Numerous viral vectors are well known in the art, including, for example, retrovirus, adenovirus, adeno-associated virus, herpes simplex virus (HSV), cytomegalovirus (CMV), vaccinia and poliovirus vectors. Retroviral vectors are less preferred since retroviruses require replicating cells and secretory glands are composed of mostly slowly replicating and/or terminally differentiated cells. Adenovirus is a preferred viral vector since this virus efficiently infects slowly replicating and/or terminally differentiated cells. The viral vector may be selected according to its preferential infection of the targeted secretory gland (*e.g.*, where the secretory gland is a salivary gland, the viral vector may be derived from an attenuated (*i.e.*, does not cause significant pathology or morbidity in the infected host, *e.g.* the virus is nonpathogenic or causes only minor disease symptoms) and/or replication-deficient mumps virus or other attenuated and/or replication-deficient virus which is substantially specific for salivary gland cells).

Where a replication-deficient virus is used as the viral vector, the production of infective virus particles containing either DNA or RNA corresponding to the DNA of interest can be produced by introducing the viral construct into a recombinant cell line which provides the missing components essential for viral replication. Preferably, transformation of the recombinant cell line with the recombinant viral vector will not result in production of replication-competent viruses, *e.g.*, by homologous recombination of the viral sequences of the recombinant cell line into the introduced viral vector. Methods for production of replication-deficient viral particles containing a nucleic acid of interest are well known in the art and are described in, for example, Rosenfeld et al., *Science* 252:431-434, 1991 and Rosenfeld et al., *Cell* 68:143-155, 1992 (adenovirus); USPN 5,139,941 (adeno-associated virus); USPN 4,861,719 (retrovirus); and USPN 5,356,806 (vaccinia virus). Methods and materials for manipulation of the mumps virus genome, characterization of mumps virus genes responsible for viral fusion and viral replication, and the structure and sequence of the mumps viral genome are described in Tanabayashi et al., *J. Virol.* 67:2928-2931, 1993; Takeuchi et al., *Archiv. Virol.*, 128:177-183, 1993; Tanabayashi et al., *Virol.* 187:801-804, 1992; Kawano et al., *Virol.*, 179:857-861, 1990; Elango et al., *J. Gen. Virol.* 69:2893-28900, 1988.

30 Conditions or Diseases Amenable to Treatment Using the Method of the Invention

Various disease conditions are amenable to treatment using the methods of the invention. One skilled in the art can recognize the appropriate protein which should be produced by the invention for treating specific disease conditions. Exemplary diseases which are amenable to treatment using the subject invention, and exemplary, appropriate proteins which can be used in treating these diseases, are shown in Table 2.

TABLE 2: Exemplary Disease Conditions Amenable to Treatment Using the Invention

5	<u>Enzyme Deficiency</u> Adenosine deaminase ¹ Purine nucleotide phosphorylase Galactosidase β -glucuronidase	<u>Endotoxic Shock/Sepsis</u> Lipid-binding protein (LBP)
	<u>Antioxidants for Cancer Therapy</u> Superoxide dismutase Catalase	<u>Anemia</u> Erythropoietin
10 15 20	<u>Cancer</u> α -Interferon γ -Interferon α -IL 1 Phenylalanine ammonia lyase Arginase L-asparaginase Uricase Granulocyte colony stimulating factor (G-CSF) Monoclonal antibodies Tissue necrosis factor	<u>Growth Factors (for use in wound healing, induction of red blood cell formation, etc.)</u> Epidermal growth factor G-CSF γ -Interferon Transforming growth factor Erythropoietin Thrombopoietin Insulin-like growth factor-I Insulin Human growth hormone
25	<u>Cardiovascular Disease</u> Tissue plasminogen activator Urokinase (native or chimeric) α_1 -antitrypsin Antithrombin-III Other proteases or protease inhibitors Apolipoproteins (particularly B-48) Circulating Scavenger Receptor APO A1 ²	<u>Diabetes</u> Insulin Glucagon Insulinotrophic hormone <u>Clotting disorders</u> Clotting factor VIII
30	<u>Obesity and Feeding</u> Ob gene product Cholecystokinin (CCK)	<u>Gastrointestinal and Pancreatic Deficiencies</u> Pepsin (for esophageal reflux) Trypsin Chymotrypsin Elastase Carboxypeptidase Lactase (for lactose deficiency) Sucrase: Intrinsic Factor (pernicious anemia)
35	<u>Bone diseases</u> Calcitonin PTH-like hormone	

TABLE 2 (cont.)

Organ-Specific Autoimmune diseases (target of antibody in parentheses)	
5	Myasthenia gravis (acetylcholine receptors)
	Graves' disease (thyroid-stimulating hormone receptor)
	Thyroiditis (thyroid, peroxidase)
	Insulin-resistant diabetes with acanthosis nigricans or with ataxia telangiectasia (Insulin receptor)
	Allergic rhinitis, asthma (Beta ₂ -adrenergic receptors)
10	Juvenile insulin-dependent diabetes (insulin, GAD65)
	Pernicious anemia (gastric parietal cells, vitamin B ₁₂ binding site of intrinsic factor)
	Addison's disease (adrenal cells)
	Idiopathic hypoparathyroidism (parathyroid cells)
	Spontaneous infertility (sperm)
	Premature ovarian failure (interstitial cells, corpus luteum cells)
15	Pemphigus (intercellular substance of skin and mucosa)
	Bullous pemphigoid (basement membrane zone of skin and mucosa)
	Primary biliary cirrhosis (mitochondria)
	Autoimmune hemolytic anemia (erythrocytes)
20	Idiopathic thrombocytopenic purpura (platelet)
	Idiopathic neutropenia (neutrophils)
	Vitiligo (melanocytes)
	Osteosclerosis and Meniere's disease (type II collagen)
	Chronic active hepatitis (nuclei of hepatocytes)
Systemic Autoimmune Diseases (defect/organ affected in parentheses)	
25	Goodpasture's syndrome (basement membranes)
	Rheumatoid arthritis (γ-globulin, EBV-related antigens, collagen types II and III)
	Sjögren's syndrome (γ-globulin, SS-A (Ro), SS-B (La))
	Systemic lupus erythematosus (nuclei, double-stranded DNA, single-stranded DNA, Sm
30	ribonucleoprotein, lymphocytes, erythrocytes, neurons, γ-globulin)
	Scleroderm (nuclei, Scl-70, SS-A(Ro), SS-B (La), centromere)
	Polymyositis (nuclei, Jo-1, PL-7, histadyl-tRNA or threonyl-tRNA synthetases, PM-1, Mi-2)
	Rheumatic fever (myocardium heart valves, choroid plexus)
	¹ For treatment of severe combined immunodeficiency
35	² Converts low-density lipoproteins to high-density lipoproteins

Transformation of Secretory Gland Cells

The DNA of interest-containing vector (*i.e.*, either a viral or non-viral vector (including naked DNA)) is introduced into the secretory gland *in vivo* via the duct system (*i.e.*, by retrograde ductal injection, which may be accomplished by perfusion (*i.e.*, continuous injection), or by a single, discontinuous injection). Retrograde ductal injection may be accomplished in the pancreas and liver by endoscopic retrograde cholangio-pancreatography (ERCP). Ductal administration provides several advantages. Because the vector is presented to the cells from "outside" the body (from the lumen), the immunological and inflammatory reactions that are commonly observed as a result of the administration of transforming formulations and their adjuvants into blood and interstitial fluid may be avoided.

Moreover, the cells of secretory glands form a monolayer that encloses the duct system. As a consequence, virtually all of the cells of the glands can be accessed by a single administration into

the duct. In this way it is possible to transfect large masses of cells in a relatively simple manner with a single procedure. The DNA of interest can thus also be administered without substantial dilution (it is only diluted by the fluid in the duct system) and without the need to develop organ specific targeting signals. In contrast, intravenous administration necessarily greatly dilutes the material and requires that it be targeted to the organ of interest in some fashion.

The amount of DNA to transform a sufficient number of secretory gland cells and provide for expression of therapeutic levels of the protein can be readily determined using an animal model (e.g., a rodent (mouse or rat) or other mammalian animal model) to assess factors such as the efficiency of transformation, the levels of protein expression achieved, the susceptibility of the targeted secretory gland cells to transformation, and the amounts of DNA required to transform secretory gland cells.

The precise amount of DNA administered will vary greatly according to a number of factors including the susceptibility of the target cells to transformation, the size and weight of the subject, the levels of protein expression desired, and the condition to be treated. For example, the amount of DNA introduced into a secretory gland of a human is generally from about 1 μ g to 200 mg, preferably from about 100 μ g to 100 mg, more preferably from about 500 μ g to 50 mg, most preferably about 10 mg. Specifically, the amount of DNA introduced into the pancreas of a human is, for example, generally from about 1 μ g to 100 mg, preferably about 100 μ g to 10 mg, more preferably from about 250 μ g to 5 mg, still more preferably from about 500 μ g to 1.5 mg, most preferably about 1 mg. The amount of DNA introduced into the salivary gland of a human is, for example, generally from about 2.5 μ g to 30 mg, more preferably from about 25 μ g to 3 mg, still more preferably from about 100 μ g to 1 mg, most preferably about 250 μ g. The amount of DNA introduced into the liver of a human is, for examples, generally from about 10 μ g to 500 mg, more preferably from about 100 μ g to 300 mg, still more preferably from about 150 μ g to 100 mg, most preferably about 1 mg.

Generally, the amounts of DNA for human therapy according to the invention can be extrapolated from the amounts of DNA effective for therapy in an animal model. For example, the amount of DNA for therapy in a human is roughly 100 times the amount of DNA effective in therapy in a rat. The amount of DNA necessary to accomplish secretory gland cell transformation will decrease with an increase in the efficiency of the transformation method used.

Concurrent Transformation of Multiple Secretory Glands

In a preferred embodiment of the invention, at least two secretory glands are transformed according to the methods of the invention. Any two secretory glands can be transformed concurrently. For example, the DNA of interest can be administered to both the pancreas and the

liver, or to both the salivary gland and the pancreas, or to both the salivary gland and the liver, or to all three. Preferably, cells of the pancreas and the liver are concurrently transformed.

Concurrent transformation of the secretory glands can be carried out several hours to several days apart or, preferably, simultaneously (*i.e.*, DNA is introduced into the two secretory glands during the same procedure. For example, where the liver and pancreas are to be concurrently transformed, the DNA formulation can be introduced simultaneously via a common duct, or separately (*e.g.*, first via the pancreatic duct with occlusion of the hepatic duct, then vice versa).

Concurrent transformation of at least two or more secretory glands can advantageously provide higher levels of expression of the polypeptide of interest in a secretory gland tissue and/or in the bloodstream and can, unexpectedly, provide for synergy between the organs (*e.g.*, to provide for higher levels of tissue expression in a secretory gland than when the secretory gland is transformed alone). For example, concurrent transformation of the pancreas and the liver results in increased levels of tissue expression in the pancreas relative to tissue levels in pancreas when it is transformed alone.

Moreover, the liver releases the polypeptide of interest in a continuous fashion that is not regulated by hormonal stimulation. The pancreas provides a relatively lower level of constitutive secretion and stores most of the polypeptide of interest and only releases large amounts after stimulation (*e.g.*, after the individual eats). Therefore, transformation of both liver and pancreas has the advantage of providing both constitutive secretion primarily from the liver, and hormonally-regulated secretion from the pancreas.

Intravenous protein therapy by transformation of salivary gland, pancreatic, and liver cells

Secretory glands transformed according to the invention facilitate high levels expression of a DNA of interest, particularly where the DNA of interest is operably linked to a strong eukaryotic promoter (*e.g.*, CMV, MMTV). The expressed protein is then secreted at high levels into the bloodstream. The protein so expressed and secreted is thus useful in treating a mammalian subject having a variety of conditions.

In a preferred embodiment, the proteins are secreted into the bloodstream at levels sufficient for intravenous protein therapy. For example, the amount of a specific protein normally released into the blood from the pancreas can be substantial, *e.g.*, a specific protein that is released into the bloodstream can be as much as 25% of the amount of duct-directed secretion of that specific protein. This amounts to as much as 1-2 mg of protein/gram of tissue being directed into the blood per hour.

Bloodstream levels of the therapeutic protein may be enhanced by several different methods. For example, bloodstream levels can be enhanced by increasing the overall level of expression of the desired protein, *e.g.*, by integration of multiple copies of the DNA of interest into the genome of the

target cells, by operably linking a strong promoter (*e.g.*, a promoter from CMV) and/or enhancer elements to the DNA of interest in the construct, or by transformation of a greater number of target cells in the subject (*e.g.*, by administration of multiple doses of the transforming material).

5 Secretion of the therapeutic protein into the bloodstream can also be enhanced by incorporating leader sequences, amino acid sequence motifs, or other elements that mediate intravenous-directed secretion into the sequence of the therapeutic protein. For example, the DNA of interest can be engineered to contain a secretion signal that directs secretion of the protein primarily into the bloodstream, thereby increasing the amount of the protein produced in the secretory gland that reaches in the bloodstream. Intravenous-directed secretion signals can be
10 identified by, for example, site-directed mutagenesis of DNA encoding a bloodstream-targeted protein (*e.g.*, insulin). The mutants can be screened by expression of the mutated DNA in secretory gland cells and subsequently determining the ratio of, for example, salivary to intravenous expression.

Alternatively, intravenous-directed secretion signals can be identified by constructing
15 recombinant, chimeric proteins composed of, for example, a putative intravenous secretion signal inserted into a saliva-directed protein. Intravenous secretion signals would then be identified by their ability to re-direct expression of the saliva-directed protein into the bloodstream. Putative intravenous secretion signals and duct system secretion signals can also be identified by comparison of DNA and amino acid sequences of proteins which are preferentially secreted into the bloodstream.
20 Areas of homology or common motifs among the proteins could then be tested as described above.

Overall secretion from secretory glands can be augmented by hormonal stimulation. For example, where the protein is primarily secreted into the duct system and is secreted at lower levels into the bloodstream, hormonal stimulation enhances intravenous secretion as well as secretion into the duct. Thus, therapeutically effective levels of the protein the bloodstream may be achieved or
25 enhanced by administration of an appropriate, secretory gland specific hormone. For example, secretory gland secretion can be enhanced by administration of a cholinergic agonist such as acetyl- β -methyl choline, or can be augmented or further augmented by control of diet (*i.e.*, eating stimulates pancreatic and salivary gland secretion). Thus, because eating a meal can elicit a secretory response, adjustment of meals (*e.g.*, frequency of meals and/or amounts eaten) can be used
30 as a dosing mechanism for delivery of the desired protein, and can be accomplished without administration of additional protein-encoding DNA.

Bloodstream-directed secretion can also be regulated at either the level of transcription, translation, or secretion. Transcriptional regulation involves the timing and level of transcription directed from the DNA of interest, while translational regulation involves the production of
35 polypeptides from transcribed RNA. Secretory regulation involves the release of polypeptides from

the cell (*e.g.*, from secretory cells in which the polypeptides to be secreted are stored within intracellular vacuoles). Methods for providing transcriptional and/or translational regulation of a DNA of interest are well known in the art (*e.g.*, transcriptional regulation through the use of inducible promoters).

5 Secretory regulation can be achieved by, for example, administration of a hormone that elicits a secretory response in the desired secretory gland, or by activity that stimulates production of such hormone(s) (*e.g.*, eating to stimulate pancreatic secretion). Unlike regulation at the level of transcription or translation, which can take many hours to become effective, regulation of secretion occurs within minutes after stimulation. Moreover, endocrine secretion from the pancreas and
10 salivary glands is stimulated by hormones and neurotransmitters that are natural components of the feeding response; thus feeding itself can act as a dosing mechanism.

 The actual number of transformed secretory gland cells required to achieve therapeutic levels of the protein of interest will vary according to several factors including the protein to be expressed, the level of expression of the protein by the transformed cells, the rate of protein secretion, the
15 partitioning of the therapeutic protein between the gastrointestinal tract and the bloodstream, and the condition to be treated. For example, the desired intravenous level of therapeutic protein can be readily calculated by determining the level of the protein present in a normal subject (for treatment of a protein deficiency), or by determining the level of protein required to effect the desired therapeutic result.

20

Application of the Method of the Invention to Achieve Euglycemia in a Diabetic Syndrome

 In another preferred embodiment of the invention, pancreatic cells are transformed using insulin-encoding DNA to provide for expression and secretion of insulin into the bloodstream of a mammalian subject. Transformation of pancreatic cells with insulin encoding DNA not only
25 provides for regulated expression of insulin in a mammalian subject, but also provides for maintenance of a euglycemic state (*i.e.*, normal blood glucose levels) in diabetic subjects for extended periods of time (*e.g.*, up to 6 to 7 days post transformation). Thus, not only is the exocrine pancreas secreting insulin to reduce blood sugar, but regulating its secretion so that blood levels are maintained at normal levels, *e.g.*, are regulated. Thus, pancreatic transformation with insulin-
30 encoding DNA can be used in the therapy of individuals having a disease or condition associated with elevated blood glucose levels (*e.g.*, diabetes (*e.g.*, type I or type II diabetes), and hyperglycemia). This aspect of the invention may be applied to regulate levels of other proteins in the bloodstream.

Assessment of Protein Therapy

The effects of expression of the protein encoded by the DNA of interest following *in vivo* transfer of the DNA of interest can be monitored in a variety of ways. Generally, a sample of blood from the subject can be assayed for the presence of the therapeutic protein. Appropriate assays for detecting a protein of interest in blood samples are well known in the art. For example, a sample of blood can be tested for the presence of the polypeptide using an antibody which specifically binds the polypeptide in an ELISA assay. This assay can be performed either qualitatively or quantitatively. The ELISA assay, as well as other immunological assays for detecting a polypeptide in a sample, are described in Antibodies: A Laboratory Manual (1988, Harlow and Lane, eds. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).

Alternatively, or in addition, the efficacy of the polypeptide therapy can be assessed by testing a sample of blood for an activity associated with the polypeptide (*e.g.*, an enzymatic activity). Furthermore, the efficacy of the therapy using the methods of the invention can be assessed by monitoring the condition of the mammalian subject for improvement. For example, where the polypeptide is erythropoietin, the subject's blood is examined for iron content or other parameters associated with anemia.

EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to carry out the invention and is not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (*e.g.*, amounts, temperatures, etc.), but some experimental error and deviation should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1: *In vivo* gene transfer of DNA encoding human growth hormone by retrograde injection of DNA into the pancreas

Four constructs for expression of human growth hormone (hGH) were prepared using techniques well known in the art (see, for example, Sambrook et al. *ibid*). The first construct, pFGH, contains the genomic hGH DNA sequence inserted in the commercially available vector pBLUESCRIPT SK+™ (Stratagene, LaJolla California) (Fig. 1). Because the hGH coding sequence is not linked to a promoter, this vector provides for no or only low-level hGH expression. Thus, the pFGH construct serves as a negative control for hGH expression in the pancreas. The second construct, pFGH.CMV, was constructed by operably inserting the promoter from the immediate

early gene of human CMV upstream of the genomic hGH sequence of the pFGH vector (Fig. 2). The third construct, pFGH.chymo, was constructed by operably inserting the rat chymotrypsin B gene promoter upstream of the genomic hGH sequence of the pFGH vector (Fig. 3). The fourth construct, pFGH.RSV, was constructed by operably inserting the promoter from the long terminal repeat (LTR) of RSV upstream of the genomic hGH sequence of the pFGH vector.

Each of the four vectors was used to transfect the pancreas of approximately 300 g adult male, Sprague-Dawley rats (pFGH + lipofectin, 4 rats; pFGH.chymo + lipofectin, 4 rats; pFGH.RSV + lipofectin, 4 rats; pFGH.CMV + lipofectin, 10 rats; pFGH.CMV without lipofectin, 7 rats; negative control (no DNA, no lipofectin), 3 rats). Pancreatic transfection was accomplished by first anesthetizing the rats and performing a laparotomy to expose the duodenum. The pancreas and the associated common bile duct were identified, and the common bile duct was cannulated either extraduodenally or through the papilla of Vater. The hepatic duct was occluded, and 100 µl of phosphate-buffered saline (PBS) containing one of the four vectors, or 100 µl of PBS alone as a negative control, were slowly injected or infused into the pancreatic duct in a retrograde direction. The vector-containing solutions were composed of 8 µg DNA per 100 µl in PBS, either with or without 6% lipofectin, a cationic lipid used to increase transfection efficiency. The solution was left in place for 5 min before secretory flow was allowed to resume and hepatic duct blockage removed. The catheter was left in place and inserted into the duodenum through a small hole to ensure adequate biliary and pancreatic flow post-operatively. The abdomen was then closed with sutures. The animals recovered fully and rapidly from the surgery without obvious side effects. This transfection method provides direct access of the vector to over 90% of the pancreatic gland cells.

At 48 hr after surgery, a blood sample was obtained to measure serum hGH levels, and the rats were sacrificed. At autopsy, the pancreas of both control and test rats appeared normal, and exhibited no gross or microscopic pathology.

The pancreas was dissected free from the mesenteric surface and was homogenized in cold 0.2 M (pH 8.0) sodium phosphate buffer (1:10 w/v) containing protease inhibitors aprotinin, leupeptin, pepstatin, and PEFABLOC SCTM. Homogenization was completed by shearing after 10 passes with a motorized pestle at approximately 4000 rpm in a glass homogenizer. The homogenate was then centrifuged at 1000 g for 15 min. The supernatant was collected and stored at -80°C until analysis. The levels of hGH in the serum and pancreatic protein samples were measured using the hGH radioimmune assay (Nichols Institute). Each assay was performed in duplicate and compared to a set of control samples.

Rats injected with the pFGH.CMV vector expressed higher levels of hGH in the pancreatic tissue (Fig. 4), compared to background levels of pancreatic hGH expression in rats injected with either no DNA (PBS alone) or the pFGH vector (hGH DNA with no promoter). The addition of

lipofectin modestly increased hGH expression in rats injected with the pFGH.CMV construct. In addition, rats transfected with the pFGH.CMV vector secreted hGH in the serum at levels increased relative to background levels and to hGH secretion levels in rats injected with either control samples (no DNA or pFGH) or with samples containing hGH DNA linked to either the chymotrypsin B or RSV promoters (Fig. 5). In Fig. 6, all data from the above experiments (including all controls and vectors) are analyzed by plotting the hGH serum levels against the hGH tissue levels. This graph shows that higher tissue levels result in higher levels of secretion into the blood. Thus, retrograde pancreatic injection of the pFGH.CMV vector successfully transfected pancreatic cells to provide both hGH pancreatic tissue expression and hGH secretion into the bloodstream.

Example 2: *In Vivo* Transformation of Pancreatic Cells by Retrograde Ductal Injection of hGH-Encoding DNA and Regulation of hGH Secretion

Eight rats were anesthetized and control blood samples (no DNA) were collected from the femoral vein of each animal. Pancreatic transfection was accomplished by exposing the duodenum by laparotomy and identifying the pancreas and the associated common bile duct. The common bile duct was cannulated either extraduodenally or through the papilla of Vater, and the hepatic duct was occluded. A 1:50 dilution of replication-defective human adenovirus (Ad5-*di* 342) supernatant in 100 μ l of phosphate-buffered saline (PBS) containing 8 μ g of the hGH-encoding plasmid pFGH.CMV (Fig. 2) was slowly infused into the pancreatic duct in a retrograde direction. The solution was left in place for approximately 5 min before secretory flow was allowed to resume and the hepatic duct blockage removed. The catheter was left in place and inserted into the duodenum through a small hole to ensure adequate biliary and pancreatic flow post-operatively. The abdomen was then closed with sutures. The animals recovered fully and rapidly from the surgery without obvious side effects.

At 48 hr after surgery, a blood sample was obtained to measure serum hGH levels (unstimulated serum levels). The cholinergic agonist McH was injected subcutaneously into each rat at 0.8 mg/kg body weight. Blood samples were collected from the inferior vena cava of each animal at 15 min intervals following McH injection. Serum was separated from the blood of all samples after clotting, and kept at -20°C prior to assay.

As shown in Fig. 9 (one representative animal) plasma levels of hGH increased markedly following McH injection, demonstrating that secretion of hGH expressed by transformed pancreatic cells is regulated by agonist stimulation. Moreover, bloodstream-directed secretion of hGH from the transformed pancreatic cells occurred at relevant, physiological levels useful in therapeutic administration (*i.e.*, at the ng/ml level).

Example 3: *In Vivo* Transformation of Salivary Glands by Retrograde Ductal Injection of DNA Encoding Human Growth Hormone

Twelve adult rats weighing approximately 300 g each were anesthetized with an intraperitoneal injection of sodium pentobarbital. A total volume of 50 μ l containing 4 μ g of the pFGH.CMV plasmid, which contains cDNA encoding human growth hormone (hGH) (Fig. 2), was introduced into each submandibular gland of 8 rats by retrograde ductal injection via the ducts leading from the oral mucosa to the salivary gland. Briefly, both the left and right Wharton's duct were cannulated intraorally with polyethylene (PE) 10 tubing, and the DNA injected into the duct system of each gland in a retrograde fashion (4 μ g/50 μ l of PBS). The material was kept in place for two minutes before normal flow was reestablished.

For three of these animals the DNA was mixed in a 6% solution of the cationic lipid Lipofectin (labeled "liposomes") from Life Technologies (Gaithersburg, MD). For four of these animals, the DNA was mixed with a 1:50 dilution of replication-defective human adenovirus (Ad5-di 342) supernatant. Control rats (4 rats) received 50 μ l 0.9% saline (control) without plasmid. No significant leakage of material or bleeding occurred. After 3 hours, the animals were awake, drinking water, and appearing normal.

Approximately 48 hours after cDNA injection, the animals were sacrificed. The right and left submandibular glands were removed and were homogenized in cold 0.2 M (pH 8.0) sodium phosphate buffer (1:10 w/v) containing the protease inhibitors aprotinin, leupeptin, pepstatin, and PEFABLOC SCTM. Homogenization was completed by shearing after 10 passes with a motorized pestle at approximately 4000 rpm in a glass homogenizer. The homogenates were centrifuged at 1000 g for 15 min, and the supernatant collected and stored at -80°C until analysis. The levels of hGH in the protein samples were measured using the hGH radioimmune assay (Nichols Institute). Each assay was performed in duplicate and compared to a set of control samples.

Each of the submandibular glands of the rats injected with the pFGH.CMV vector expressed hGH in the salivary gland tissue; hGH expression was undetectable in the control rats' salivary glands (Fig. 7).

Example 4: *In Vivo* Transformation of Salivary Glands by Retrograde Ductal Injection of hGH-Encoding DNA and Regulation of hGH Secretion

Three adult rats weighing approximately 300 g each were anesthetized with an intraperitoneal injection of sodium pentobarbital. A control blood sample (prior to DNA) was drawn from the femoral vein of each animal. A total of 4 μ g of the hGH-encoding plasmid pFGH.CMV (Fig. 2) in 50 μ l, was introduced into each submandibular gland of each rat by retrograde ductal

injection via the ducts leading from the oral mucosa to the salivary gland as described above in Example 2. No significant leakage of material or bleeding occurred. After 3 hours, the animals were awake, drinking water, and appearing normal.

Forty-eight hours after cDNA injection, the animals were again anesthetized and a control blood sample was drawn from the femoral vein of each animal (unstimulated serum level). The cholinergic agonist acetyl- β -methyl choline (McH) was injected subcutaneously at 0.8 mg/kg body weight into each animal. Blood samples were collected from the femoral vein of each animal at 10 min, 20 min, 40 min, and 50 min after McH injection. Serum was separated from the blood of all samples after clotting, and kept at -20°C prior to assay.

As shown in Fig. 8 (one representative animal), secretion of hGH into the bloodstream was dramatically increased in response to administration of McH, peaking at 40 min. Thus, these data demonstrate that introduction of hGH-encoding DNA into the salivary gland results in bloodstream-directed secretion of hGH and regulation by cholinergic stimulation. Moreover, regulation is at the level of secretion, not transcription, since transcriptional regulation would not result in increased hGH bloodstream levels in such a short period.

Example 5: Treatment of Diabetes Mellitus Over a Three Day Period by *In Vivo* Transformation of Pancreatic Cells by Retrograde Ductal Injection with Insulin-Encoding DNA

Streptozotocin, which induces diabetes mellitus in rats, was administered to 8 male Sprague-Dawley rats (260-280 g) after overnight fasting by intraperitoneal injection in 1 mM citrate buffer (pH 4.5) (Sigma) at 65 mg/kg of body weight. One hour later, animals were anesthetized with Nembutal and the body cavity opened to expose the gastrointestinal tract. Each animal was given the appropriate DNA construct directly by retrograde injection in the pancreatic duct in a 100 μ l injection volume containing 8 μ g DNA plus adenovirus (Ad5-*di* 342)(3×10^{10} viral particles) as described above. Test animals (4 rats) received the human insulin-encoding construct pBat16.hInsG1.M2. The pBAT16.hInsG1.M2 construct (Fig. 10) encodes an insulin gene containing a site-directed mutation of the second protease site to create a furin recognition site; this construct provides for enhanced expression of processed insulin in non-neuroendocrine cells. In addition, the human β -globin first intron replaces the first insulin gene intron which is inefficiently spliced. Control animals (4 rats) received the control construct CMV-GFP, which contains a green fluorescent protein (GFP)-encoding sequence operably linked to a CMV promoter. The animals recovered fully and rapidly from the surgery without obvious side effects. Body weight and blood glucose were monitored daily for three days post-injection. Blood glucose was measured by the glucose oxidase method (Lifescan, Milpitas, CA).

As shown in Fig. 11, treatment of the streptozotocin-induced diabetic rats with the insulin-encoding construct resulted in maintenance of almost complete euglycemia for 3 days. In contrast, control animals that received the GFP-encoding construct remained hyperglycemic throughout the test period. The data show that introduction of insulin-encoding DNA into the pancreas results in pancreatic cell transformation, as well as secretion of insulin by the transformed pancreatic cells at levels sufficient to overcome diabetes in an animal model. Moreover, these results show that the method of the invention provides regulated and relatively normal blood glucose levels. Surprisingly, the exocrine pancreas regulates the release of insulin such that blood sugar levels are maintained at regulated levels (normally the endocrine pancreas is responsible for regulation of bloodstream-directed secretion).

Example 6: Treatment of Diabetes Mellitus Over a Six Day Period by *In Vivo* Transformation of Pancreatic Cells by Retrograde Ductal Injection with Insulin-Encoding DNA

Streptozotocin was administered to 14 rats at 70 mg/kg body weight by intraperitoneal injection to induce diabetes mellitus. The animals were then anesthetized by intraperitoneal injection of sodium pentobarbital. Two rats did not receive streptozotocin and served as one negative control. Insulin-encoding DNA in the pBAT16.hInsG1.M2 construct (Fig. 10) was administered to 8 of the streptozotocin-injected rats by retrograde ductal injection as described above. Six streptozotocin-treated rats received either 100 µl of saline without DNA (2 animals) or a control DNA without the human insulin gene (4 animals) by pancreatic retrograde ductal injection as additional negative controls. The animals recovered fully and rapidly from the surgery without obvious side effects. Blood samples were collected from the femoral vein of each animal at 24 hr intervals for 6 days. Human insulin was measured using a double antibody radioimmunoassay (Linco Laboratories, Saint Louis, MO).

As shown in (Fig. 12), blood glucose levels were significantly decreased in the diabetic rats that received the insulin-encoding DNA (+Strep, + DNA) relative to diabetic the rats that received no DNA (+Strep, No DNA). Furthermore, these decreased blood glucose levels were observed throughout the entire 6 day course of the experiment. Thus, these data show that introduction of insulin-encoding DNA into the pancreas results in persistent expression of insulin, and that the insulin expressed by the transformed pancreatic cells is secreted into the bloodstream and can function in regulation of blood glucose at levels sufficient to overcome diabetes in an animal model. As shown in Fig. 12, elevated insulin levels for such an extended period additionally demonstrate prolonged expression from the DNA introduced into the pancreatic cells.

Example 7: *In Vivo* Transformation of Pancreatic Cells by Retrograde Ductal Injection of Green Fluorescent Protein-Encoding DNA and Expression in Pancreatic Cells

To identify the pancreatic cells that expressed the recombinant protein, DNA encoding green fluorescent protein (GFP) was used to transform pancreatic cells according to the methods of the invention. EGFP cDNA from plasmid pEGFP.C2 (Clontech) was inserted into pFOX. The EGFP sequence was modified to contain an SV40 nuclear localization signal, in-frame at the 3' end. This addition allowed for partial nuclear localization and facilitated immunohistochemical detection. The CMV immediate early promoter was positioned upstream of the first intron of human β -globin to create the expression vector pFOX.EGFP.N2.CMV.

After fasting overnight, Male Sprague-Dawley rats (260-280 g) were anesthetized and the body cavity opened to expose the gastrointestinal tract. The green fluorescent protein (GFP)-encoding construct pFOX.EGFP.N2.CMV was administered to each animal by retrograde injection in the pancreatic duct in a 100 μ l injection volume containing 8 μ g DNA premixed with adenovirus (3×10^{10} viral particles) as described above. The animals recovered fully and rapidly from the surgery without obvious side effects.

Seventy-two hours post-treatment, the animals were sacrificed, and pancreases were removed and weighed (wet weight). Samples of each pancreas were fixed in 5% buffered formalin for 24-48 hours at room temperature. Fixed tissues were dehydrated and imbedded in paraffin, and 5 μ m sections were processed for immunohistochemistry using standard techniques. Endogenous peroxidase was quenched in 0.7% H_2O_2 /MeOH, and antigen retrieval was performed using Citra solution (Biogenex, San Ramon, CA) according to the manufacturers' instructions. Sections were preincubated for 30 minutes in 5% goat serum/phosphate-buffered saline (PBS), and then incubated overnight at 4°C with primary antisera diluted in 5% goat serum/PBS.

The primary antisera were selected from either anti-GFP antisera (1:1500; Clontech, Palo Alto, CA), anti-insulin antisera (1:500; Dako, Carpinteria, CA), or non-specific rabbit sera (1:1500). The following day all sections were incubated with biotinylated goat anti-rabbit antiserum (5 μ g/ml; Vector, Burlingame, CA) for 30 minutes at room temperature, and then incubated with streptavidin-aminohexanol-biotin horseradish peroxidase (HRP) complex (Vectastain-Elite, Vector). Protein was visualized by reaction with the peroxidase substrate 3,3'-diamino-benzidine tetrahydrochloride (DAB; Sigma). The color reaction was followed by a brief counter stain in 1% methyl green (Sigma) prior to mounting. Negative controls included staining of sections from pancreas not injected with CMV-GFP, and omission of primary antiserum.

Staining for GFP was observed in the pancreas of animals treated with GFP DNA, but not in control animals. GFP expression was restricted to exocrine cells; there was no staining in either ductal or islet cells. Moreover, expression was observed in 0.1-1.0% of exocrine cells. Endogenous

insulin was detected in adjacent sections; but GFP expression did not co-localize with insulin expression, suggesting that the pancreatic cells primarily transformed are exocrine, not endocrine cells. Under the conditions studied there was no histological indication of inflammatory infiltration as a consequence of ductal injection of the vector.

5 These data show that introduction of the DNA construct results in successful transformation of pancreatic cells, despite the introduction of the construct against the flow of pancreatic juices and the high concentrations of DNase in the pancreatic juice. Moreover, these data, combined with the data above showing that transformation of the pancreas results in bloodstream-directed secretion of the encoded protein, and suggest that transformation of exocrine pancreatic cells results in
10 bloodstream-directed secretion of the protein encoded by the introduced construct. Furthermore, because insulin staining and GFP staining did not co-localize, introduction of the GFP-encoding construct resulted in transformation of exocrine tissue, which is normally associated with protein secretion into the gastrointestinal tract, rather than endocrine tissue, which is normally associated with bloodstream-directed secretion. Despite this, bloodstream-directed secretion was still obtained
15 at physiologically relevant levels sufficient to treat diabetes mellitus in an animal model as evidenced in the examples above.

Example 8: *In Vivo* Transformation of Liver Cells by Retrograde Ductal Injection of hGH-Encoding DNA and Bloodstream-Directed hGH Secretion

20 Four rats were anesthetized and control blood samples (no DNA) were collected from the femoral vein of each animal. Transfection of liver cells was accomplished by exposing the duodenum by laparotomy and identifying the liver and the associated common bile duct. The common bile duct was cannulated either extraduodenally or through the papilla of Vater. The tubing was advanced to the bifurcation of the hepatic duct in order to prevent injected material from
25 entering the distally located pancreatic drainage. A 1:50 dilution of replication-defective human adenovirus supernatant in 100 μ l of phosphate-buffered saline (PBS) containing 8 μ g of the hGH-encoding plasmid pFGH.CMV (Fig. 2) or 100 μ l of PBS alone (no DNA) were slowly infused into the hepatic duct in a retrograde direction. The solution was left in place for approximately 2 min to 5 min before secretory flow was allowed to resume and the pancreatic duct blockage removed. The
30 catheter was left in place and inserted into the duodenum through a small hole to ensure adequate biliary and pancreatic flow post-operatively. The abdomen was then closed with sutures. The animals recovered fully and rapidly from the surgery without obvious side effects.

Plasma hGH levels were measured 2 days after treatment; the results are shown in Fig. 13. Each data point in Fig. 13 represents the mean \pm standard error of the mean (SEM) for three animals.
35 These data demonstrate that liver cells were transformed with the hGH-encoding DNA.

Furthermore, hGH was secreted by the transformed liver cells into the bloodstream at physiologically relevant levels.

Example 9: *In Vivo* Transformation of Pancreatic Cells with hGH-Encoding DNA and Expression in Rat Exocrine Pancreas and Plasma

Following overnight fasting and anesthesia with pentobarbital, the abdominal cavity of the rats was opened and the pancreatic duct cannulated external to the duodenum with PE 10 tubing as described above. Eight to twenty-five micrograms of each of pFGH (promoter less construct), pFGH.chymo (construct with the chymotrypsin promoter), pFGH.RSV (construct with the RSV promoter), and pFGH.CMV (construct with the CMV promoter) was injected in a total volume of 100 μ l of PBS into the pancreas via the pancreatic duct as described above. Immediately prior to injection construct samples were optionally premixed with either Lipofectin (6-12% vol:vol) or adenovirus (3×10^{10} viral particles). The material was kept in the duct for 5 min prior to establishing normal flow. The abdomen was closed and the animals allowed to recover.

Forty-eight hours later the pancreas was harvested, plasma obtained, and human growth hormone measured. The animals were anesthetized, blood samples taken (either from the femoral vein or inferior vena cava), and the transfected tissue removed. The tissue was homogenized in PBS containing 5 mM Na_2HPO_4 (pH 7.8) at a tissue to fluid ratio of 1:10 using a motorized mortar and pestle. Large particulate material in the homogenate was removed by sedimentation at $10,000 \times g$ for 30 minutes, and the supernatant assayed for the protein of interest. The results are shown in Figs. 14-17. All data shown are the mean \pm the SEM.

The effects of the various promoters upon tissue expression and secretion of hGH into the bloodstream are shown in Figs. 14 and 15, respectively. In these experiments, the constructs were mixed with lipofection prior to administration. Of the promoters tested, the CMV promoter was by far the most effective, and produced high levels of hGH in tissue (in the range of 150 ng/g tissue wet weight) when compared to either promoter less controls, or plasmids containing RSV and chymotrypsin promoters (Fig. 14). The cationic lipid adjuvant Lipofectin increased expression by about 50%, and pre-mixing the plasmid with adenovirus enhanced tissue expression five fold (Fig. 15). Expression of hGH at 24, 48 or 72 hours after injection was similar under all conditions studied.

As shown in Figs. 16 and 17, hGH was secreted into plasma. Plasmids containing the CMV promoter increased circulating levels of hGH five times above background (Fig. 16). With plasmid alone, plasma hGH concentrations in the range of 60 to 80 pg/ml were routinely observed. Premixing the plasmids with adjuvants also increased circulating hGH levels (Fig. 17). Lipofectin

increased plasma levels by an additional 50%, and adenovirus by 75%, when compared to plasmid alone.

5 Example 10: Comparison of hGH Secretion by Rat Liver, Pancreas, and Combined Liver and Pancreas Transformed with hGH-Encoding DNA

Eight micrograms of the pFGH.CMV construct premixed with adenovirus as described above in Example 9, was injected into the ducts of either the liver, the pancreas, or both organs of the same animal. Where only the liver or the pancreas was transformed (liver alone or pancreas alone), the DNA was introduced according to the methods described above. Where both the liver and pancreas were transformed, the DNA-containing formulation was introduced into the hepatic duct first, and then the tubing partially withdrawn to provide access to the pancreatic duct system. A temporary ligature was then placed around the hepatic duct to prevent the second infusion from entering the parenchyma of the liver. Thus, animals in which both the pancreas and liver were transformed received two doses of the DNA-containing formulation. Plasma hGH levels were measured two days later.

In animals having transformed liver (liver alone) or pancreas (pancreas alone), hGH was expressed in liver or pancreatic tissue, respectively, and hGH detected in plasma under both circumstances. Tissue levels in liver when transformed alone were far lower than in the pancreas when transformed alone (less than 1 ng/g, as compared to about 500 ng/g), but hGH concentration in plasma of animals in which only the liver was transformed was nonetheless comparable to hGH plasma levels in animals having only the pancreas transformed (in the range of 0.15 ng/ml; Fig. 18). These results are consistent with the observation that, in contrast to the exocrine cells of the pancreas and salivary glands, hepatocytes secrete most of what they produce soon after synthesis.

When pancreas and liver were both transfected, plasma levels were higher than seen when the glands were treated individually (nearly 0.3 ng/ml) -- a value approximately equal to the sum of that observed for the two organs separately. Surprisingly, transformation of both liver and pancreas resulted in tissue levels in the pancreas being significantly increased relative to tissue levels in the pancreas when the pancreas was transformed alone (Fig. 19)

30 Example 11: Human growth hormone (hGH) expression in rat salivary gland.

Four micrograms of the pFGH.CMV construct, premixed with either Lipofectin or adenovirus, was injected into each submandibular gland via retrograde ductal injection (via Wharton's duct) as described above. Two days later, each gland was harvested and hGH content was measured

As shown in Fig. 20, tissue levels of hGH averaged about 50 ng/g tissue wet weight. Plasma hGH levels were in the 20-40 pg/ml range. As in the pancreas, the addition of adenovirus increased tissue hGH levels, in this case to 100 ng/g (Fig. 20).

5 Example 12: Stimulation of Human growth hormone (hGH) secretion.

Even when exocrine secretory cells store large amounts of protein, such as after a period of fasting, they secrete these proteins at a low rate under unstimulated conditions (*i.e.* basal or constitutive secretion). Greater rates are achieved when exogenous stimulants (*e.g.*, hormonal stimulants and/or stimulation associated with eating) are applied. To determine whether secretion of
10 the engineered protein would be enhanced during feeding, pancreatic secretion was stimulated with a secretory stimulant. For these experiments we used animals in which both pancreas and liver were transfected. Eight micrograms of the pFGH.CMV construct were injected into ducts of both the pancreas and liver of four rats as described above. A blood sample was taken prior to injection as a control. Two days after transfection, a second control blood sample was taken and the rats were
15 treated with the cholinergic agonist, acetyl- β -methylcholine (McH) (0.8 mg/kg body weight).

As shown in Fig. 21, hGH secretion was increased three fold within 30 minutes of stimulation, with plasma levels approaching 1.0 ng/ml. Similar enhancement of hGH secretion was observed when either the pancreas was studied alone, or when the salivary glands were studied alone. These data show that hGH secretion is enhanced by stimulation with a cholinergic agonist.
20 Thus secretion of hGH is regulated in a manner similar to secretion of endogenous proteins.

Although the concentration of hGH in plasma was correlated to the level of hormone in the pancreas ($r=0.55$, $p<0.01$, $n=41$), at high tissue levels, plasma concentration was not linearly proportional to tissue content. For example, addition of adenovirus to the hGH vector produced a five fold increase in tissue levels relative to the plasmid alone (Fig. 20), but only about a two fold
25 increase in plasma concentration (see, *e.g.*, Figs. 16 and 17, Example 9 above). This lack of proportionality indicates that it is not the concentration of product in the cells alone that determines the rate of secretion into blood, but that at high tissue levels, secretion is limited by other factors. This result is similar to what is observed for endogenous protein secretion and suggests that secretion of the engineered protein is regulated in much the same manner.

30

Example 13: Human Insulin Expression and Secretion in Diabetic Rat Pancreas

In an attempt to treat a disease state, diabetes mellitus, we expressed human insulin in the exocrine pancreas. Fasted experimental and control animals received intra-peritoneal streptozotocin (Sigma; 65 mg/kg body weight, in 1mM citrate buffer, pH4.5) on day zero one hour prior to
35 administration of the insulin-encoding construct. The experimental animals subsequently received 8

µg of the insulin plasmid (pBAT16.hInsG1.M2) premixed with adenovirus and injected into the pancreatic duct, also on day zero. The pBAT16.hInsG1.M2 construct contains the human insulin cDNA linked to a CMV immediate early promoter, which is positioned upstream of the first intron of human β-globin. The human insulin cDNA was mutated to convert the second protease site, between peptides C and A, to a furin recognition site. This allows for correct proteolytic processing of mature insulin in non-endocrine cells.

Plasma insulin and glucose levels were determined for up to six days. Plasma glucose levels in diabetic rats (n=3), and diabetic rats treated with the pBAT16.hInsG1.M2 plasmid (n=3), measured over a three day period, are shown in Fig. 22. Plasma insulin levels in diabetic rats (n=3), and diabetic rats treated with the pBAT16.hInsG1.M2 plasmid (n = 3), measured over a three day period, are shown in Fig. 23. Plasma glucose levels in individual diabetic (n=3) and pBAT16.hInsG1.M2 plasmid- treated diabetic rats (n=3), measured over a six day period, are shown in Fig. 24.

As a consequence of streptozotocin administration, blood glucose levels rose from the normal level of 100 mg/dl to 300-400 mg/dl within 24 hours and remained elevated for the duration of the study (Fig. 22). Treatment with the human insulin plasmid reduced blood glucose levels in diabetic rats to the normal range (Figs. 22 and 24), and concentrations of insulin remained near pre-treatment values (Fig. 23). Blood glucose levels were euglycemic for the duration of the study (6 days; Fig. 24). Animals transfected with a control plasmid remained diabetic (data not shown). These data show that regulation of insulin secretion in response to feeding was effective.

Example 14: *In vivo* gene transfer of DNA encoding human growth hormone by retrograde injection of DNA into the salivary gland

A DNA expression construct encoding human growth hormone (hGH) is prepared by operably linking a CMV promoter to hGH-encoding DNA. The expression cassette is then inserted into a construct such as the bacterial plasmid pBR322. *Escherichia coli* is then transformed with the plasmid using conventional transformation procedures. *E. coli* containing the plasmid are selected by virtue of the tetracycline or ampicillin resistance encoded by pBR322, and the transformed bacterial cells propagated in culture. Plasmid DNA is then isolated from the transformed bacterial cell culture and the DNA purified by cesium gradient.

Approximately 250 µg of the purified plasmid DNA containing hGH DNA is injected into the salivary gland of a human patient by retrograde ductal injection via a salivary gland duct. Expression and intravenous secretion of the protein is assessed using the method described above.

Example 15: *In vivo* gene transfer of DNA encoding human growth hormone by retrograde ductal injection of naked DNA into the pancreas

A construct containing hGH-encoding DNA (Marshall et al., *Biotechnology* 24:293-298, 1992) operably linked to the CMV promoter is resuspended in 0.9% saline and a volume of the DNA solution is administered to a human patient. Approximately 1 mg of DNA is delivered to the pancreas of the patient by cannulation of the pancreatic duct by duodenal intubation using endoscopic retrograde cholangio-pancreatography. Expression and secretion of human growth hormone into the bloodstream is assessed by detection of the protein in the patient's blood.

10 Example 16: *In vivo* gene transfer of DNA encoding human insulin by cannulation of naked DNA into the liver

A construct containing human insulin-encoding DNA operably linked to the CMV promoter is resuspended in 0.9% saline and a volume of the DNA solution is administered to a human patient. Approximately 1 mg of DNA is delivered to the patient's liver by cannulation of the hepatic duct. Expression and secretion of human growth hormone into the bloodstream is assessed by detection of the protein in the patient's blood.

Example 17: *In vivo* gene transfer of human insulin-encoding DNA to both the pancreas and the liver of a patient

20 A construct containing human insulin-encoding DNA operably linked to the CMV promoter is resuspended in 0.9% saline. A volume of the DNA solution is administered to a human patient so as to transform both pancreatic and hepatic cells (e.g., by introducing the DNA solution into the common bile duct before it splits into the hepatic and pancreatic ducts). Approximately 2 mg of DNA is delivered to the patient's liver by cannulation of the hepatic duct; in addition, approximately 25 1 mg of DNA is delivered to the pancreas via retrograde injection via the pancreatic duct. Expression and secretion of human growth hormone into the bloodstream is assessed by detection of the protein in the patient's blood.

Following procedures similar to those described above, other therapeutic proteins can be expressed from DNA inserted in the genome of a secretory gland cell by gene transfer according to the invention.

The invention now being fully described, it will be apparent to one of ordinary skill in the art that many changes and modifications can be made thereto without departing from the spirit or scope of the appended claims.

What is claimed is:

1. A method of delivering a polypeptide to a mammalian subject, the method comprising:
 - 5 introducing *in vivo* into at least two secretory glands of a mammal a construct comprising a DNA of interest that encodes a desired polypeptide and a eukaryotic promoting sequence operably linked to the DNA of interest, said introducing being by retrograde ductal injection;
wherein the introduced construct genetically transforms cells of each secretory gland and
wherein expression of the DNA by the genetically transformed cells of the secretory glands results in
10 bloodstream-directed secretion of the protein encoded by the DNA of interest.
 2. The method of claim 1, wherein the secretory glands are selected from the group consisting of pancreas, liver, and salivary gland.
 - 15 3. The method of claim 1, wherein the secretory glands are liver and pancreas.
 4. The method of claim 1, wherein the DNA of interest encodes a polypeptide selected from the group consisting of insulin, growth hormone, interferon-alpha 2b, Interferon-alpha 2a, interferon-alpha N1, filgrastim, insulinotropin, imiglucerase, clotting factor VIII, interferon-beta1b,
20 erythropoietin, sargramostim, interleukin-2, interferon-gamma, anti-CD3 antibody, GPIIb/IIIa monoclonal antibody, adenosine deaminase, interleukin-8, insulin-like growth factor-I, platelet-derived growth factor, epidermal growth factor, hemoglobin.
 5. The method of claim 1, wherein the DNA of interest encodes human insulin.
 - 25 6. A method of controlling blood glucose levels in a mammalian subject having a disease or condition associated with hyperglycemia, the method comprising:
introducing into a mammalian patient's pancreas a construct comprising insulin-encoding DNA and a eukaryotic promoting sequence operably linked to the insulin-encoding, said introducing
30 being by intraductal injection, thereby accomplishing genetic transformation of a pancreatic cell; and
allowing the genetically transformed cell to express the protein and secrete the protein into the bloodstream of the patient in a therapeutically effective amount thereby treating the patient;
wherein blood glucose levels in the mammalian subject are decreased to achieve a
euglycemic state.

35

7. The method of claim 6, wherein the euglycemic state is maintained for at least three days.
8. The method of claim 6, wherein the euglycemic state is maintained for at least 6
5 days.
9. The method of claim 6, the method further comprising enhancing insulin expression by administration of a cholinergic agonist to the mammalian subject.
10. The method of claim 6, the method further comprising enhancing insulin expression
10 by ingestion of a meal.

1/18




FIG. 1

2/18





FIG. 2

3/18



4/18




FIG. 4

5/18




FIG. 5

6/18







FIG. 6

7/18

Salivary: Tissue Expression of hGH**FIG. 7**



9/18






FIG. 10


10/18



**FIG. 11**

11/18



**FIG. 12**



12/18

**FIG. 13**






14/18



**FIG. 16****FIG. 17**

**FIG. 18****FIG. 19**

16/18

**FIG. 20**

**FIG. 21****FIG. 22**

**FIG. 23****FIG. 24**

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/20199

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01N 43/04, 63/00; C12N 13/00

US CL : 514/44; 424/93.21; 435/455

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44; 424/93.21; 435/455

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
APS, CAS ONLINE, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	O'CONNELL et al. Transfer of a Gene encoding the Anticandidal Protein Histatin 3 to Salivary Glands. Human Gene Therapy. 01 December 1996, Vol. 7, pages 2255-2261, especially pages 2259 and 2260.	1-10
Y	FEHER et al. Exit Routes For Secretory Proteins From The Dog Pancreas. Acta Physiologica Academiae Scientiarum Hungaricae. 1980, Tomus 56, No. 4, pages 401-410, especially pages 407-409.	1-10
Y	KAGAMI et al. Evidence for the systemic Delivery of a Transgene Product from Salivary Glands. Human Gene Therapy. 10 November 1996, Vol. 7, pages 2177-2184, especially pages 2181-2183.	1-10

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

18 DECEMBER 1998

Date of mailing of the international search report

25 JAN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DAVE NGUYEN

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/20199

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ADESANYA et al. Immediate Inflammatory responses to Adenovirus-Mediated Gene Transfer in rat Salivary Glands. Human Gene Therapy. 10 June 1996, Vol. 7, pages 1085-1093, abstract, especially pages 1090 and 1091.	1-10
A,P	US 5,792,751 A (LEDLEY ET AL) 11 August 1998, columns 4-10.	1-10

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*

THIS PAGE BLANK (USPTO)

PCT

LD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A23B 7/10, A23J 1/00, C07K 16/12, C07H 15/11, 21/02, 21/04, C12N 9/16, 15/85, 15/63, 1/20, C12Q 1/68	A1	(11) International Publication Number: WO 99/08539 (43) International Publication Date: 25 February 1999 (25.02.99)
(21) International Application Number: PCT/US98/16728 (22) International Filing Date: 13 August 1998 (13.08.98) (30) Priority Data: 08/910,798 13 August 1997 (13.08.97) US (71) Applicant: DIVERSA CORPORATION [US/US]; 10665 Sorrento Valley Road, San Diego, CA 92121 (US). (72) Inventor: KRETZ, Keith; 973 Hawthorne Drive, San Marcos, CA 92069 (US). (74) Agent: HAILE, Lisa, A.; Fish & Richardson P.C., Suite 1400, 4225 Executive Square, La Jolla, CA 92037 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: NOVEL PHYTASE (57) Abstract <p>The invention provides a purified phytate enzyme derived from <i>Escherichia coli</i> B. The enzyme has a molecular weight of about 47.1 kilodaltons and has phytase activity (SEQ ID NO:2). The enzyme can be produced from native or recombinant host cells and can be used to aid in the digestion of phytate where desired. In particular, the phytase of the present invention can be used in animal feed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-

NOVEL PHYTASE

Field of the Invention

This invention relates to newly identified polynucleotides, polypeptides encoded by such polynucleotides, the use of such polynucleotides and polypeptides, as well as the production and isolation of such polynucleotides and polypeptides. More particularly, the polypeptides of the present invention have been identified as phytases and in particular, enzymes having phytase activity.

Background

Minerals are essential elements for the growth of all organisms. For livestock production of monogastric animals (e.g., pigs, poultry) and fish, feed is commonly supplemented with minerals. Plant seeds are a rich source of minerals since they contain ions that are complexed with the phosphate groups of phytic acid. Ruminants do not require inorganic phosphate and minerals because microorganisms in the rumen produce enzymes that catalyze conversion of phytate (myo-inositol-hexaphosphate) to inositol and inorganic phosphate. In the process, minerals that have been complexed with phytate are released.

Phytate occurs as a source of stored phosphorous in virtually all plant feeds (Phytic Acid, Chemistry and Applications, E. Graf (Ed.), Pilatus Press: Minneapolis, MN, U.S.A., 1986). Phytic acid forms a normal part of the seed in cereals and legumes. It functions to bind dietary minerals that are essential to the new plant as it emerges from the seed. When the phosphate groups of phytic acid are removed by the seed enzyme phytase, the ability to bind metal ions is lost and the minerals become available to the plant. In livestock feed grains, the trace minerals bound by phytic acid are only partially available for absorption by monogastric animals, which lack phytase activity. Although some hydrolysis of phytate occurs in the colon, most phytate passes through the gastrointestinal tract of monogastric animals and is excreted in the manure contributing

-2-

to fecal phosphate pollution problems in areas of intense livestock production. Inorganic phosphorous released in the colon has no nutritional value to livestock because inorganic phosphorous is absorbed only in the small intestine. Thus, a significant amount of the nutritionally important dietary minerals are potentially not available to monogastric animals.

Conversion of phytate to inositol and inorganic phosphorous can be catalyzed by microbial enzymes referred to broadly as phytases. Phytases such as phytase #EC 3.1.3.8 are capable of catalyzing hydrolysis of myo-inositol hexaphosphate to D-myo-inositol 1,2,4,5,6-pentaphosphate and orthophosphate. Certain fungal phytases reportedly hydrolyze inositol pentaphosphate to tetra-, tri-, and lower phosphates; e.g., *A. ficuum* phytases reportedly produce mixtures of myoinositol di- and mono-phosphate (Ullah, 1988). Phytase producing microorganisms comprise bacteria such as *Bacillus subtilis* (V.K. Powar and V.J. Jagannathan, *J. Bacteriol.* 151:1102-1108, 1982) and *Pseudomonas* (D.J. Cosgrove, *Austral. J. Biol. Sci.* 2:1207-1220, 1970); yeasts such as *Saccharomyces cerevisiae* (N.R. Nayini and P. Markakis, *Lebensmittel Wissenschaft und Technologie* 17:24-26, 1984); and fungi such as *Aspergillus terreus* (K. Yamada, et al., *Agric. Biol. Chem.* 32:1275-1282, 1968). The possible use of microbes capable of producing phytase as a feed additive for monogastric animals has been reported previously (Shieh and Ware, U.S. Patent No. 3,297,548; Nelson, T.S. et al., *J. Nutrition* 101:1289-1294, 1971).

Microbial phytases may also reportedly be useful for producing animal feed from certain industrial processes, e.g., wheat and corn waste products. The wet milling process of corn produces glens sold as animal feeds. Addition of phytase may reportedly improve the nutritional value of the feed product. Fungal phytase enzymes and process conditions (t~50(C and pH ~5.5) have been reported previously in European Patent Application 0 321 004. In processing soybean meal the presence of phytate reportedly renders the meal and wastes unsuitable for feeds used in rearing fish, poultry and other non-ruminants as well as calves fed on milk. Phytase is reportedly useful for improving the nutrient and

-3-

commercial value of this high protein soy material (see Finase Enzymes by Alko, Rajamaki, Finland). A combination of phytase and a pH 2.5 optimum acid phosphatase form *A. niger* has been used by Alko, Ltd as an animal feed supplement in their phytic acid degradative product Finas F and Finase S. A cost-effective source of phytase would greatly enhance the value of soybean meals as an animal feed (Shieh et al., 1969).

Phytase and less specific acid phosphatases are produced by the fungus *Aspergillus ficuum* as extracellular enzymes (Shieh et al., 1969). Ullah reportedly purified a phytase from wild-type *A. ficuum* that had an apparent molecular weight of 61.7 kDa (on SDS-PAGE; as corrected for glycosylation); pH optima at pH 2.5 and pH 5.5; a K_m of about 40m; and, a specific activity of about 50U/mg (Ullah, A., Preparative Biochem 18:443-458, 1988); PCT patent application WO 91/05053 also reportedly discloses isolation and molecular cloning of a phytase from *Aspergillus ficuum* with pH optima at pH 2.5 and pH 5.5, a K_m of about 250m, and specific activity of about 100U/mg protein.

Acid phosphatases are enzymes that catalytically hydrolyze a wide variety of phosphate esters and usually exhibit pH optima below 6.0 (Hollander, 1971); e.g., #EC 3.1.3.2 catalyzes hydrolysis of orthophosphoric monoesters to orthophosphate products. An acid phosphatase has reportedly been purified from *A. ficuum*. The deglycosylated form of the acid phosphatase has an apparent molecular weight of 32.6 kDa (Ullah et al., 1987).

The object of the present invention provides a recombinant phytase isolated from *Escherichia coli* B that improves the efficiency of release of phosphorous from phytate and the salts of phytic acid. Another object of the present invention provides a source of a recombinant enzyme that is suitable for commercial use in feeds and industrial processes with minimal processing.

-4-

Summary of the Invention

The present invention provides a polynucleotide and a polypeptide encoded thereby which has been identified as a phytase enzyme having phytase activity. In accordance with one aspect of the present invention, there is provided a novel recombinant enzyme,
5 as well as active fragments, analogs and derivatives thereof.

In accordance with another aspect of the present invention, there are provided isolated nucleic acid molecules encoding the enzyme of the present invention including mRNA, DNA, cDNA, genomic DNA as well as active analogs and fragments of such enzyme.

In accordance with yet a further aspect of the present invention, there is provided a
10 process for producing such polypeptide by recombinant techniques comprising culturing recombinant prokaryotic and/or eukaryotic host cells, containing a nucleic acid sequence encoding an enzyme of the present invention, under conditions promoting expression of said enzyme and subsequent recovery of said enzyme.

In accordance with yet a further aspect of the present invention, there is provided a
15 process for utilizing such enzyme, or polynucleotide encoding such enzymes for use in commercial processes, such as, for example, processes that liberate minerals from phytates in plant materials either in vitro, i.e., in feed treatment processes, or in vivo, i.e., by administering the enzyme to animals.

In accordance with yet a further aspect of the present invention, there is provided a
20 process for utilizing such enzymes, or polynucleotides encoding such enzymes, for *in vitro* purposes related to scientific research, for example, to generate probes for identifying similar sequences which might encode similar enzymes from other organisms.

-5-

These and other aspects of the present invention should be apparent to those skilled in the art from the teachings herein.

BRIEF DESCRIPTION OF THE DRAWINGS

5 The following drawings are illustrative of embodiments of the invention and are not meant to limit the scope of the invention as encompassed by the claims.

Figure 1 shows the nucleotide and deduced amino acid sequences the enzyme of the present invention. Sequencing was performed using a 378 automated DNA sequencer (Applied Biosystems, Inc.).

10 Figure 2 shows the pH and temperature profile and stability data for the phytase enzyme of the present invention. The assay used for these analysis is the following for the detection of phytase activity: Phytase activity is measured by incubating 150(l of the enzyme preparation with 600(l of 2 mM sodium phytate in 100 mM Tris HCl buffer pH 7.5, supplemented with 1mM CaCl₂ for 30 minutes at 37(C. After incubation the reaction is stopped by adding 750(l of 5% trichloroacetic acid. Phosphate released was measured
15 against phosphate standard spectrophotometrically at 700nm after adding 1500(l of the color reagent (4 volumes of 1.5% ammonium molybdate in 5.5% sulfuric acid and 1 volume of 2.7% ferrous sulfate; Shimizu, M., 1992; Biosci. Biotech. Biochem., 56:1266-1269). OD at 700nm is indicated on the Y-axis of the graphs in Figure 2. Temperature or pH is indicated on the X-axis of the graphs.

DETAILED DESCRIPTION OF THE INVENTION

The term "gene" means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons).

5 A coding sequence is "operably linked to" another coding sequence when RNA polymerase will transcribe the two coding sequences into a single mRNA, which is then translated into a single polypeptide having amino acids derived from both coding sequences. The coding sequences need not be contiguous to one another so long as the expressed sequences are ultimately processed to produce the desired protein.

10 "Recombinant" enzymes refer to enzymes produced by recombinant DNA techniques; *i.e.*, produced from cells transformed by an exogenous DNA construct encoding the desired enzyme. "Synthetic" enzymes are those prepared by chemical synthesis.

15 A DNA "coding sequence of" or a "nucleotide sequence encoding" a particular enzyme, is a DNA sequence which is transcribed and translated into an enzyme when placed under the control of appropriate regulatory sequences. A "promotor sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. The promoter is part of the DNA sequence. This sequence region has a start codon at its 3' terminus. The promoter sequence does include the minimum number of bases where elements necessary to initiate transcription at levels detectable above background. However, after the RNA polymerase binds the sequence and transcription is initiated at the start codon (3' terminus with a promoter), transcription proceeds downstream in the 3' direction. Within the promotor sequence will be found a transcription initiation site (conveniently defined by mapping with nuclease S1) as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

20

25

-7-

The present invention provides purified a recombinant enzyme that catalyzes the hydrolysis of phytate to inositol and free phosphate with release of minerals from the phytic acid complex. An exemplary purified enzyme is a phytase derived from *Escherichia coli* B. This exemplary enzyme is shown in Figure 1, SEQ ID NO:2.

5 The polynucleotide encoding SEQ ID NO:2 was originally recovered from genomic DNA isolated from *Escherichia coli* B as described below. It contains an open reading frame encoding a protein of 432 amino acid residues.

10 In one embodiment, the phytase enzyme of SEQ ID NO:2 of the present invention has a molecular weight of about 47,056 kilodaltons as measured by SDS-PAGE gel electrophoresis and an inferred molecular weight from the nucleotide sequence of the gene. The pI is 6.70. The pH and temperature profile and stability data for this enzyme is presented in Figure 2. This purified enzyme may be used to catalyze the hydrolysis of phytate to inositol and free phosphate where desired. The phytase enzyme of the present invention has a high thermostability.

15 In accordance with an aspect of the present invention, there are provided isolated nucleic acid molecules (polynucleotides) which encode for the mature enzyme having the deduced amino acid sequence of Figure 1.

20 This invention can be used to isolate nucleic acid sequences substantially similar to the isolated nucleic acid molecule encoding an phytase enzyme disclosed in Figure 1 (SEQ ID NO:1),. Isolated nucleic acid sequences are substantially similar if: (i) they are capable of hybridizing under stringent conditions, hereinafter described, to SEQ ID NO:1; or (ii) they encode DNA sequences which are degenerate to SEQ ID NO:1. Degenerate DNA sequences encode the amino acid sequence of SEQ ID NO:2, but have variations in the nucleotide coding sequences. As used herein, "substantially similar"

25 refers to the sequences having similar identity to the sequences of the instant invention.

-8-

The nucleotide sequences that are substantially similar can be identified by hybridization or by sequence comparison. Enzyme sequences that are substantially similar can be identified by one or more of the following: proteolytic digestion, gel electrophoresis and/or microsequencing.

5 One means for isolating a nucleic acid molecule encoding a phytase enzyme is to probe a genomic gene library with a natural or artificially designed probe using art recognized procedures (see, for example: Current Protocols in Molecular Biology, Ausubel F.M. *et al.* (EDS.) Green Publishing Company Assoc. and John Wiley Interscience, New York, 1989, 1992). It is appreciated to one skilled in the art that SEQ ID NO:1, or fragments
10 thereof (comprising at least 15 contiguous nucleotides), is a particularly useful probe. Other particular useful probes for this purpose are hybridizable fragments to the sequences of SEQ ID NO:1 (comprising at least 10 contiguous nucleotides and at least 70% complementary to a target sequence), is a particularly useful probe. Other particular useful probes for this purpose are hybridizable fragments to the sequences
15 of SEQ ID NO:1 (*i.e.*, comprising at least 10 contiguous nucleotides and at least 70% complementary to a target sequence).

With respect to nucleic acid sequences which hybridize to specific nucleic acid sequences disclosed herein, hybridization may be carried out under conditions of reduced stringency, medium stringency or even stringent conditions. As an example of
20 oligonucleotide hybridization, a polymer membrane containing immobilized denatured nucleic acid is first prehybridized for 30 minutes at 45°C in a solution consisting of 0.9 M NaCl, 50 mM NaH₂PO₄, pH 7.0, 5.0 mM Na₂EDTA, 0.5% SDS, 10X Denhardt's, and 0.5 mg/mL polyriboadenylic acid. Approximately 2×10^7 cpm (specific activity $4-9 \times 10^8$ cpm/ug) of ³²P end-labeled oligonucleotide probe are then added to the solution.
25 After 12-16 hours of incubation, the membrane is washed for 30 minutes at room temperature in 1X SET (150 mM NaCl, 20 mM Tris hydrochloride, pH 7.8, 1 mM Na₂EDTA) containing 0.5% SDS, followed by a 30 minute wash in fresh 1X SET at

SUBSTITUTE SHEET (RULE 26)

-9-

Tm-10(C for the oligo-nucleotide probe. The membrane is then exposed to auto-radiographic film for detection of hybridization signals.

Stringent conditions means hybridization will occur only if there is at least 90% identity, preferably at least 95% identity and most preferably at least 97% identity between the sequences. See J. Sambrook *et al.*, Molecular Cloning, A Laboratory Manual (2d Ed. 1989) (Cold Spring Harbor Laboratory) which is hereby incorporated by reference in its entirety.

"Identity" as the term is used herein, refers to a polynucleotide sequence which comprises a percentage of the same bases as a reference polynucleotide (SEQ ID NO:1). For example, a polynucleotide which is at least 90% identical to a reference polynucleotide, has polynucleotide bases which are identical in 90% of the bases which make up the reference polynucleotide and may have different bases in 10% of the bases which comprise that polynucleotide sequence.

The present invention also relates to polynucleotides which differ from the reference polynucleotide such that the changes are silent changes, for example the changes do not alter the amino acid sequence encoded by the polynucleotide. The present invention also relates to nucleotide changes which result in amino acid substitutions, additions, deletions, fusions and truncations in the enzyme encoded by the reference polynucleotide (SEQ ID NO:1). In a preferred aspect of the invention these enzymes retain the same biological action as the enzyme encoded by the reference polynucleotide.

It is also appreciated that such probes can be and are preferably labeled with an analytically detectable reagent to facilitate identification of the probe. Useful reagents include but are not limited to radioactivity, fluorescent dyes or enzymes capable of catalyzing the formation of a detectable product. The probes are thus useful to isolate complementary copies of DNA from other animal sources or to screen such sources for

-10-

related sequences.

The present invention provides a substantially pure phytase enzyme. The term "substantially pure" is used herein to describe a molecule, such as a polypeptide (*e.g.*, a phytase polypeptide, or a fragment thereof) that is substantially free of other proteins, lipids, carbohydrates, nucleic acids, and other biological materials with which it is naturally associated. For example, a substantially pure molecule, such as a polypeptide, can be at least 60%, by dry weight, the molecule of interest. The purity of the polypeptides can be determined using standard methods including, *e.g.*, polyacrylamide gel electrophoresis (*e.g.*, SDS-PAGE), column chromatography (*e.g.*, high performance liquid chromatography (HPLC)), and amino-terminal amino acid sequence analysis.

The phytase polypeptide included in the invention can have the amino acid sequences of Phytase shown in Figure 1 (SEQ ID NO:1). Phytase polypeptides, such as those isolated from *E.coli* B, can be characterized by catalyzing the hydrolysis of phytate to inositol and free phosphate with the release of minerals from the phytic acid complex.

Also included in the invention are polypeptides having sequences that are "substantially identical" to the sequence of a phytase polypeptide, such as one of SEQ ID 1. A "substantially identical" amino acid sequence is a sequence that differs from a reference sequence only by conservative amino acid substitutions, for example, substitutions of one amino acid for another of the same class (*e.g.*, substitution of one hydrophobic amino acid, such as isoleucine, valine, leucine, or methionine, for another, or substitution of one polar amino acid for another, such as substitution of arginine for lysine, glutamic acid for aspartic acid, or glutamine for asparagine).

Fragments of the phytase polypeptide of the present invention can retain at least one phytase-specific activity or epitope. Phytase activity can be assayed by examining the

-11-

5 catalysis of phytate to inositol and free phosphate. For example, a phytase polypeptide fragment containing, *e.g.*, at least 8-10 amino acids can be used as an immunogen in the production of phytase-specific antibodies. The fragment can contain, for example, an amino acid sequence that is conserved in phytases, and this amino acid sequence can contain amino acids that are conserved in phytases. Such fragments can easily be identified by comparing the sequences of phytases found in Figure 1. In addition to their use as peptide immunogens, the above-described phytase fragments can be used in immunoassays, such as ELISAs, to detect the presence of phytase-specific antibodies in samples.

10 Other phytase polypeptides included in the invention are polypeptides having amino acid sequences that are at least 50% identical to the amino acid sequence of a phytase polypeptide, such as SEQ ID NO:2. The length of comparison in determining amino acid sequence homology can be, for example, at least 15 amino acids, for example, at least 20, 25, or 35 amino acids. Homology can be measured using standard sequence analysis software (e.g., Sequence Analysis Software Package of the Genetics Computer
15 Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705; also see Ausubel, et al., *supra*). Such procedures and algorithms include, for example, a BLAST program (Basic Local Alignment Search Tool at the National Center for Biological Information), ALIGN, AMAS (Analysis of Multiply Aligned Sequences), AMPS (Protein Multiple Sequence Alignment), ASSET (Aligned Segment Statistical Evaluation Tool), BANDS, BESTSCOR, BIOSCAN (Biological Sequence Comparative Analysis Node), BLIMPS (BBlocks IMProved Searcher), FASTA, Intervals & Points, BMB, CLUSTAL V, CLUSTAL W, CONSENSUS, LCONSENSUS, WCONSENSUS, Smith-Waterman algorithm, DARWIN, Las Vegas algorithm, FNAT
20 (Forced Nucleotide Alignment Tool), Framealign, Framesearch, DYNAMIC, FILTER, FSAP (Fristensky Sequence Analysis Package), GAP (Global Alignment Program), GENAL, GIBBS, GenQuest, ISSC (Sensitive Sequence Comparison), LALIGN (Local Sequence Alignment), LCP (Local Content Program), MACAW (Multiple Alignment

-12-

Construction & Analysis Workbench), MAP (Multiple Alignment Program), MBLKP, MBLKN, PIMA (Pattern-Induced Multi-sequence Alignment), SAGA (Sequence Alignment by Genetic Algorithm) and WHAT-IF.

5 The phytase polypeptide of the invention can be obtained using any of several standard methods. For example, phytase polypeptides can be produced in a standard recombinant expression systems (see below), chemically synthesized (this approach may be limited to small phytase peptide fragments), or purified from organisms in which they are naturally expressed.

10 The invention also provides isolated nucleic acid molecules that encode the phytase polypeptide described above. For example, nucleic acids that encode SEQ ID NO:1 are included in the invention. These nucleic acids can contain naturally occurring nucleotide sequences, or sequences that differ from those of the naturally occurring nucleic acids that encode phytases, but encode the same amino acids, due to the degeneracy of the genetic code. The nucleic acids of the invention can contain DNA or RNA nucleotides,
15 or combinations or modifications thereof. Exemplary nucleic acids of the invention are shown in SEQ ID NO:1.

20 By "isolated nucleic acid" is meant a nucleic acid, *e.g.*, a DNA or RNA molecule, that is not immediately contiguous with the 5' and 3' flanking sequences with which it normally is immediately contiguous when present in the naturally occurring genome of the organism from which it is derived. The term thus describes, for example, a nucleic acid that is incorporated into a vector, such as a plasmid or viral vector; a nucleic acid that is incorporated into the genome of a heterologous cell (or the genome of a homologous cell, but at a site different from that at which it naturally occurs); and a nucleic acid that exists as a separate molecule, *e.g.*, a DNA fragment produced by PCR
25 amplification or restriction enzyme digestion, or an RNA molecule produced by *in vitro* transcription. The term also describes a recombinant nucleic acid that forms part of a

-13-

hybrid gene encoding additional polypeptide sequences that can be used, for example, in the production of a fusion protein.

The nucleic acid molecules of the invention can be used as templates in standard methods for production of phytase gene products (*e.g.*, phytase RNAs and phytase polypeptides). In addition, the nucleic acid molecules that encode phytase polypeptides (and fragments thereof) and related nucleic acids, such as (1) nucleic acids containing sequences that are complementary to, or that hybridize to, nucleic acids encoding phytase polypeptides, or fragments thereof (*e.g.*, fragments containing at least 10, 12, 15, 20, or 25 nucleotides); and (2) nucleic acids containing sequences that hybridize to sequences that are complementary to nucleic acids encoding phytase polypeptides, or fragments thereof (*e.g.*, fragments containing at least 10, 12, 15, 20, or 25 nucleotides); can be used in methods focused on their hybridization properties. For example, as is described in further detail below, such nucleic acid molecules can be used in the following methods: PCR methods for synthesizing phytase nucleic acids, methods for detecting the presence of a phytase nucleic acid in a sample, screening methods for identifying nucleic acids encoding new phytase family members. Oligonucleotide probes useful for screening methods are from 10 to about 150 nucleotides in length. Further, such probes are preferably 10 to about 100 nucleotides in length and more preferably from 10 to about 50 nucleotides in length.

The invention also includes methods for identifying nucleic acid molecules that encode members of the phytase polypeptide family in addition to SEQ ID NO:1. In these methods, a sample, *e.g.*, a nucleic acid library, such as a cDNA library, that contains a nucleic acid encoding a phytase polypeptide is screened with a phytase-specific probe, *e.g.*, a phytase-specific nucleic acid probe. Phytase-specific nucleic acid probes are nucleic acid molecules (*e.g.*, molecules containing DNA or RNA nucleotides, or combinations or modifications thereof) that specifically hybridize to nucleic acids encoding phytase polypeptides, or to complementary sequences thereof. The term

-14-

"phytase-specific probe," in the context of this method of invention, refers to probes that bind to nucleic acids encoding phytase polypeptides, or to complementary sequences thereof, to a detectably greater extent than to nucleic acids encoding other enzymes, or to complementary sequences thereof.

5 The invention facilitates production of phytase-specific nucleic acid probes. Methods for obtaining such probes can be designed based on the amino acid sequences shown in Figure 1. The probes, which can contain at least 10, *e.g.*, at least 15, 25, 35, 50, 100, or 150 nucleotides, can be produced using any of several standard methods (see, *e.g.*, Ausubel, *et al.*, *supra*). For example, preferably, the probes are generated using PCR
10 amplification methods. In these methods, primers are designed that correspond to phytase-conserved sequences (see Figure 1), which can include phytase-specific amino acids, and the resulting PCR product is used as a probe to screen a nucleic acid library, such as a cDNA library.

15 The coding sequences for the phytase enzymes of the present invention were identified by preparing *E.coli* B genomic DNA, for example, and recovering (via, for example, PCR amplification) from the genomic DNA, DNA encoding phytase activity. Such methods for recovery are well-known in the art. One means, for example, comprises designing amplification primers to recover the coding sequence, amplifying the gene from the genomic DNA, subcloning the DNA into a vector, transforming the resulting
20 construct into a host strain, and expressing the phytase enzyme for evaluation. Such procedures are well known in the art and methods are provided, for example, in Maniatis, T., *et al.*, Molecular Cloning, Cold Spring Harbor Press, New York, 1982, which is hereby incorporated by reference in its entirety.

25 In a preferred embodiment, the enzyme of the present invention, was isolated from an *E.coli* B genomic DNA by the following technique:

E.coli B genomic DNA was obtained from Sigma (Catalog # D-2001), St. Louis, New

-15-

Jersey.

The following primers were used to amplify the gene directly from the genomic DNA:

5' primer gtttctgaattcaaggaggaatttaaATGAAAGCGATCTTAATCCCATT

3' primer gtttctggatccTTACAAACTGCACGCCGGTAT

5 Pfu polymerase was used according to manufacturers protocol (Stratagene Cloning Systems, Inc., La Jolla, CA).

PCR product and pQE60 vector (Qiagen) were both digested with EcoRI and BglII restriction endonucleases (New England Biolabs) according to manufacturers protocols. Ligation and transformation into, and expression in M15 pREP4 host cells (Qiagen) yields c-term 6X-His tagged protein.

15 Fragments of the full length gene of the present invention may be used as a hybridization probe for a cDNA or a genomic library to isolate the full length DNA and to isolate other DNAs which have a high sequence similarity to the gene or similar biological activity. Probes of this type have at least 10, preferably at least 15, and even more preferably at least 30 bases and may contain, for example, at least 50 or more bases. The probe may also be used to identify a DNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete gene including regulatory and promoter regions, exons, and introns.

20 The isolated nucleic acid sequences and other enzymes may then be measured for retention of biological activity characteristic to the enzyme of the present invention, for example, in an assay for detecting enzymatic phytase activity. Such enzymes include truncated forms of phytase, and variants such as deletion and insertion variants.

Examples of such assays include the following assay for the detection of phytase activity: Phytase activity can be measured by incubating 150(l of the enzyme preparation with 600(l of 2 mM sodium phytate in 100 mM Tris HCl buffer pH 7.5, supplemented with

-16-

1mM CaCl₂ for 30 minutes at 37°C. After incubation the reaction is stopped by adding 750(1 of 5% trichloroacetic acid. Phosphate released was measured against phosphate standard spectrophotometrically at 700nm after adding 1500(1 of the color reagent (4 volumes of 1.5% ammonium molybdate in 5.5% sulfuric acid and 1 volume of 2.7% ferrous sulfate; Shimizu, M., 1992; Biosci. Biotech. Biochem., 56:1266-1269). One unit of enzyme activity is defined as the amount of enzyme required to liberate one (mol Pi per min under assay conditions. Specific activity can be expressed in units of enzyme activity per mg of protein.

The enzyme of the present invention has enzymatic activity with respect to the hydrolysis of phytate to inositol and free phosphate.

The polynucleotide of the present invention may be in the form of DNA which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. The coding sequence which encodes the mature enzyme may be identical to the coding sequences shown in Figure 1 and/or that of the deposited clone (SEQ ID NO:1), or may be a different coding sequence which coding sequence, as a result of the redundancy or degeneracy of the genetic code, encodes the same mature enzyme as the DNA of Figure 1 (e.g., SEQ ID NO:1).

The polynucleotide which encodes for the mature enzyme of Figure 1 (e.g., SEQ ID NO:2) may include, but is not limited to: only the coding sequence for the mature enzyme; the coding sequence for the mature enzyme and additional coding sequence such as a leader sequence or a proprotein sequence; the coding sequence for the mature enzyme (and optionally additional coding sequence) and non-coding sequence, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the mature enzyme.

-17-

Thus, the term "polynucleotide encoding an enzyme (protein)" encompasses a polynucleotide which includes only coding sequence for the enzyme as well as a polynucleotide which includes additional coding and/or non-coding sequence.

5 The present invention further relates to variants of the hereinabove described polynucleotides which encode for analogs and derivatives of the enzyme having the deduced amino acid sequence of Figure 1 (*e.g.*, SEQ ID NO:2). The variant of the polynucleotide may be a naturally occurring allelic variant of the polynucleotide or a non-naturally occurring variant of the polynucleotide.

10 Thus, the present invention includes polynucleotides encoding the same mature enzyme as shown in Figure 1 as well as variants of such polynucleotides which variants encode for a derivative or analog of the enzyme of Figure 1. Such nucleotide variants include deletion variants, substitution variants and addition or insertion variants.

15 As hereinabove indicated, the polynucleotide may have a coding sequence which is a naturally occurring allelic variant of the coding sequence shown in Figure 1. As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which may have a substitution, deletion or addition of one or more nucleotides, which does not substantially alter the function of the encoded enzyme.

20 The present invention also includes polynucleotides, wherein the coding sequence for the mature enzyme may be fused in the same reading frame to a polynucleotide sequence which aids in expression and secretion of an enzyme from a host cell, for example, a leader sequence which functions to control transport of an enzyme from the cell. The enzyme having a leader sequence is a preprotein and may have the leader sequence cleaved by the host cell to form the mature form of the enzyme. The polynucleotides may also encode for a proprotein which is the mature protein plus additional 5' amino
25 acid residues. A mature protein having a prosequence is a proprotein and is an inactive

-18-

form of the protein. Once the prosequence is cleaved an active mature protein remains.

Thus, for example, the polynucleotide of the present invention may encode for a mature enzyme, or for an enzyme having a prosequence or for an enzyme having both a prosequence and a presequence (leader sequence).

5 The present invention further relates to a enzyme which has the deduced amino acid sequence of Figure 1, as well as analogs and derivatives of such enzyme.

The terms "derivative" and "analog" when referring to the enzyme of Figure 1 means a enzyme which retains essentially the same biological function or activity as such enzyme. Thus, an analog includes a proprotein which can be activated by cleavage of
10 the proprotein portion to produce an active mature enzyme.

The enzyme of the present invention may be a recombinant enzyme, a natural enzyme or a synthetic enzyme, preferably a recombinant enzyme.

The derivative or analog of the enzyme of Figure 1 may be (i) one in which one or more of the amino acid residues are substituted with an amino acid residue which is not
15 encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature enzyme is fused with another compound, such as a compound to increase the half-life of the enzyme (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature enzyme, such as a leader or secretory sequence or a sequence which is
20 employed for purification of the mature enzyme or a proprotein sequence. Such derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

-19-

The enzymes and polynucleotides of the present invention are preferably provided in an isolated form, and preferably are purified to homogeneity.

The term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or enzyme present in a living animal is not isolated, but the same polynucleotide or enzyme, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or enzymes could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

The enzymes of the present invention includes an enzyme of Figure 1 (in particular the mature enzyme) as well as enzymes which have at least 70% similarity (preferably at least 70% identity) to an enzyme of Figure 1 and more preferably at least 90% similarity (more preferably at least 90% identity) to an enzyme of Figure 1 and still more preferably at least 95% similarity (still more preferably at least 95% identity) to an enzyme of Figure 1 and also include portions of such enzymes with such portion of the enzyme generally containing at least 30 amino acids and more preferably at least 50 amino acids.

As known in the art "similarity" between two enzymes is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one enzyme to the sequence of a second enzyme. Similarity in nucleic acid and amino acid sequences may be determined by procedures and algorithms which are well-known in the art. Such procedures and algorithms include, for example, a BLAST program (Basic Local Alignment Search Tool at the National Center for Biological Information), ALIGN, AMAS (Analysis of Multiply Aligned Sequences), AMPS (Protein Multiple Sequence Alignment), ASSET (Aligned Segment Statistical Evaluation Tool), BANDS, BESTSCOR, BIOSCAN (Biological Sequence Comparative Analysis Node), BLIMPS (BBlocks IMProved Searcher), FASTA, Intervals & Points, BMB, CLUSTAL V,

-20-

CLUSTAL W, CONSENSUS, LCONSENSUS, WCONSENSUS, Smith-Waterman
algorithm, DARWIN, Las Vegas algorithm, FNAT (Forced Nucleotide Alignment Tool),
Framealign, Framesearch, DYNAMIC, FILTER, FSAP (Fristensky Sequence Analysis
Package), GAP (Global Alignment Program), GENAL, GIBBS, GenQuest, ISSC
5 (Sensitive Sequence Comparison), LALIGN (Local Sequence Alignment), LCP (Local
Content Program), MACAW (Multiple Alignment Construction & Analysis Workbench),
MAP (Multiple Alignment Program), MBLKP, MBLKN, PIMA (Pattern-Induced
Multi-sequence Alignment), SAGA (Sequence Alignment by Genetic Algorithm) and
WHAT-IF.

10 A variant, i.e. a "analog" or "derivative" enzyme, and reference enzyme may differ in
amino acid sequence by one or more substitutions, additions, deletions, fusions and
truncations, which may be present in any combination.

Among preferred variants are those that vary from a reference by conservative amino
acid substitutions. Such substitutions are those that substitute a given amino acid in a
15 polypeptide by another amino acid of like characteristics. Typically seen as conservative
substitutions are the replacements, one for another, among the aliphatic amino acids Ala,
Val, Leu and Ile; interchange of the hydroxyl residues Ser and Thr, exchange of the
acidic residues Asp and Glu, substitution between the amide residues Asn and Gln,
exchange of the basic residues Lys and Arg and replacements among the aromatic
20 residues Phe, Tyr.

Most highly preferred are variants which retain the same biological function and activity
as the reference polypeptide from which it varies.

Fragments or portions of the enzymes of the present invention may be employed for
producing the corresponding full-length enzyme by peptide synthesis; therefore, the
25 fragments may be employed as intermediates for producing the full-length enzymes.
Fragments or portions of the polynucleotides of the present invention may be used to

-21-

synthesize full-length polynucleotides of the present invention.

The present invention also relates to vectors which include polynucleotides of the present invention, host cells which are genetically engineered with vectors of the invention and the production of enzymes of the invention by recombinant techniques.

5 Host cells are genetically engineered (transduced or transformed or transfected) with the vectors containing the polynucleotides of this invention. Such vectors may be, for example, a cloning vector or an expression vector. The vector may be, for example, in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters,
10 selecting transformants or amplifying the genes of the present invention. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

The polynucleotides of the present invention may be employed for producing enzymes by recombinant techniques. Thus, for example, the polynucleotide may be included in
15 any one of a variety of expression vectors for expressing an enzyme. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other vector may be used as long as it
20 is replicable and viable in the host.

The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and others are deemed to be within the scope of those skilled in the art.

-22-

The DNA sequence in the expression vector is operatively linked to an appropriate expression control sequence(s) (promoter) to direct mRNA synthesis. As representative examples of such promoters, there may be mentioned: LTR or SV40 promoter, the *E. coli. lac* or *trp*, the phage lambda P_L promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. The vector may also include appropriate sequences for amplifying expression.

In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in *E. coli*.

The vector containing the appropriate DNA sequence as hereinabove described, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein.

As representative examples of appropriate hosts, there may be mentioned: bacterial cells, such as *E. coli*, *Streptomyces*, *Bacillus subtilis*; fungal cells, such as yeast; insect cells such as *Drosophila S2* and *Spodoptera Sf9*; animal cells such as CHO, COS or Bowes melanoma; adenoviruses; plant cells, *etc.* The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein.

More particularly, the present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter,

-23-

operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. The following vectors are provided by way of example; Bacterial: pQE70, pQE60, pQE-9 (Qiagen), pBluescript II (Stratagene); pTRC99a, pKK223-3, pDR540, pRIT2T (Pharmacia); Eukaryotic: pXT1, pSG5 (Stratagene) pSVK3, pBPV, pMSG, pSVLSV40 (Pharmacia). However, any other plasmid or vector may be used as long as they are replicable and viable in the host.

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda P_R, P_L, and trp. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art.

In a further embodiment, the present invention relates to host cells containing the above-described constructs. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (Davis, L., Dibner, M., Battey, L., Basic Methods in Molecular Biology, (1986)).

The constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the enzymes of the invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells

-24-

under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), the disclosure of which is hereby incorporated by reference.

Transcription of the DNA encoding the enzymes of the present invention by higher eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp that act on a promoter to increase its transcription. Examples include the SV40 enhancer on the late side of the replication origin bp 100 to 270, a cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), λ -factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated enzyme. Optionally, the heterologous sequence can encode a fusion enzyme including an N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product.

Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation

-25-

and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

As a representative but nonlimiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced by appropriate means (e.g., temperature shift or chemical induction) and cells are cultured for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, such methods are well known to those skilled in the art.

-26-

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, Cell, 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

The enzyme can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

The enzymes of the present invention may be a naturally purified product, or a product of chemical synthetic procedures, or produced by recombinant techniques from a prokaryotic or eukaryotic host (for example, by bacterial, yeast, higher plant, insect and mammalian cells in culture). Depending upon the host employed in a recombinant production procedure, the enzymes of the present invention may be glycosylated or may be non-glycosylated. Enzymes of the invention may or may not also include an initial methionine amino acid residue.

The enzyme of this invention may be employed for any purpose in which such enzyme activity is necessary or desired. In a preferred embodiment the enzyme is employed for

-27-

catalyzing the hydrolysis of phytate. The degradation of phytate may be used in animal feed.

In a preferred embodiment, the enzyme of the present invention is a phytase enzyme which is stable to heat and is heat resistant and catalyzes the enzymatic hydrolysis of phytate, *i.e.*, the enzyme is able to renature and regain activity after a brief (*i.e.*, 5 to 30 seconds), or longer period, for example, minutes or hours, exposure to temperatures of 50(C optimum above 50(C.

The enzymes, their fragments or other derivatives, or analogs thereof, or cells expressing them can be used as an immunogen to produce antibodies thereto. These antibodies can be, for example, polyclonal or monoclonal antibodies. The present invention also includes chimeric, single chain, and humanized antibodies, as well as Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

Antibodies generated against the enzymes corresponding to a sequence of the present invention can be obtained by direct injection of the enzymes into an animal or by administering the enzymes to an animal, preferably a nonhuman. The antibody so obtained will then bind the enzymes itself. In this manner, even a sequence encoding only a fragment of the enzymes can be used to generate antibodies binding the whole native enzymes. Such antibodies can then be used to isolate the enzyme from cells expressing that enzyme.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, 1975, Nature, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, Immunology Today 4:72), and

-28-

the EBV-hybridoma technique to produce human monoclonal antibodies (Cole, et al., 1985, in Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96).

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic enzyme products of this invention. Also, transgenic mice may be used to express humanized antibodies to immunogenic enzyme products of this invention.

Antibodies generated against the enzyme of the present invention may be used in screening for similar enzymes from other organisms and samples. Such screening techniques are known in the art. Antibodies may also be employed as a probe to screen gene libraries generated from this or other organisms to identify this or cross reactive activities.

Isolation and purification of polypeptides produced in the systems described above can be carried out using conventional methods, appropriate for the particular system. For example, preparative chromatography and immunological separations employing antibodies, such as monoclonal or polyclonal antibodies, can be used.

The term "antibody," as used herein, refers to intact immunoglobulin molecules, as well as fragments of immunoglobulin molecules, such as Fab, Fab', (Fab')₂, Fv, and SCA fragments, that are capable of binding to an epitope of an endoglucanase polypeptide. These antibody fragments, which retain some ability to selectively bind to the antigen (*e.g.*, an endoglucanase antigen) of the antibody from which they are derived, can be made using well known methods in the art (see, *e.g.*, Harlow and Lane, *supra*), and are described further, as follows.

-29-

(1) A Fab fragment consists of a monovalent antigen-binding fragment of an antibody molecule, and can be produced by digestion of a whole antibody molecule with the enzyme papain, to yield a fragment consisting of an intact light chain and a portion of a heavy chain.

5 (2) A Fab' fragment of an antibody molecule can be obtained by treating a whole antibody molecule with pepsin, followed by reduction, to yield a molecule consisting of an intact light chain and a portion of a heavy chain. Two Fab' fragments are obtained per antibody molecule treated in this manner.

10 (3) A (Fab')₂ fragment of an antibody can be obtained by treating a whole antibody molecule with the enzyme pepsin, without subsequent reduction. A (Fab')₂ fragment is a dimer of two Fab' fragments, held together by two disulfide bonds.

(4) An Fv fragment is defined as a genetically engineered fragment containing the variable region of a light chain and the variable region of a heavy chain expressed as two chains.

15 (5) A single chain antibody ("SCA") is a genetically engineered single chain molecule containing the variable region of a light chain and the variable region of a heavy chain, linked by a suitable, flexible polypeptide linker.

20 As used in this invention, the term "epitope" refers to an antigenic determinant on an antigen, such as a phytase polypeptide, to which the paratope of an antibody, such as an phytase-specific antibody, binds. Antigenic determinants usually consist of chemically active surface groupings of molecules, such as amino acids or sugar side chains, and can have specific three-dimensional structural characteristics, as well as specific charge characteristics.

-30-

As is mentioned above, antigens that can be used in producing phytase-specific antibodies include phytase polypeptides, *e.g.*, any of the phytase shown in Figures 1 polypeptide fragments. The polypeptide or peptide used to immunize an animal can be obtained by standard recombinant, chemical synthetic, or purification methods. As is well known in the art, in order to increase immunogenicity, an antigen can be conjugated to a carrier protein. Commonly used carriers include keyhole limpet hemocyanin (KLH), thyroglobulin, bovine serum albumin (BSA), and tetanus toxoid. The coupled peptide is then used to immunize the animal (*e.g.*, a mouse, a rat, or a rabbit). In addition to such carriers, well known adjuvants can be administered with the antigen to facilitate induction of a strong immune response.

Phytase-specific polyclonal and monoclonal antibodies can be purified, for example, by binding to, and elution from, a matrix containing a phytase polypeptide, *e.g.*, the phytase polypeptide (or fragment thereof) to which the antibodies were raised. Additional methods for antibody purification and concentration are well known in the art and can be practiced with the phytase-specific antibodies of the invention (see, for example, Coligan, *et al.*, Unit 9, *Current Protocols in Immunology*, Wiley Interscience, 1994).

Anti-idiotypic antibodies corresponding to phytase-specific antigens are also included in the invention, and can be produced using standard methods. These antibodies are raised to phytase-specific antibodies, and thus mimic phytase-specific epitopes.

The members of a pair of molecules (*e.g.*, an antibody-antigen pair or a nucleic acid pair) are said to "specifically bind" to each other if they bind to each other with greater affinity than to other, non-specific molecules. For example, an antibody raised against an antigen to which it binds more efficiently than to a non-specific protein can be described as specifically binding to the antigen. (Similarly, a nucleic acid probe can be described as specifically binding to a nucleic acid target if it forms a specific duplex with the target by base pairing interactions (see above).)

-31-

The present invention is further described with reference to the following examples; however, it is to be understood that the present invention is not limited to such examples. All parts or amounts, unless otherwise specified, are by weight.

5 In one aspect of the invention, a method for producing an phytase enzyme, such as those shown in Figures 1, is provided. The method includes growing a host cell which contains a polynucleotide encoding the enzyme (*e.g.*, SEQ ID NO: 1), under conditions which allow the expression of the nucleic acid, and isolating the enzyme encoded by the nucleic acid. Methods of culturing the host cell are described in the Examples and are known by those of skill in the art.

10 In another embodiment, the invention provides a method for catalyzing the hydrolysis of phytate to inositol and free phosphate with release of minerals from the phytic acid complex. The method includes contacting phytate with a degrading effective amount of an enzyme of the invention, such as the enzyme shown in SEQ ID NO:1. The term "degrading effective" amount refers to the amount of enzyme which is required to
15 degrade at least 50% of the phytate, as compared to phytate not contacted with the enzyme. Preferably, at least 80% of the phytate is degraded.

In another embodiment, the invention provides a method for hydrolyzing phospho-mono-ester bonds in phosphate, the method including administering an effective amount of an enzyme of the invention (*e.g.*, SEQ ID NO:1), to yield inositol and free
20 phosphate. An "effective" amount refers to the amount of enzyme which is required to hydrolyze at least 50% of the phospho-mono-ester bonds, as compared to phytate not contacted with the enzyme. Preferably, at least 80% of the bonds are hydrolyzed.

In order to facilitate understanding of the following examples certain frequently occurring methods and/or terms will be described.

-32-

"Plasmids" are designated by a lower case p preceded and/or followed by capital letters and/or numbers. The starting plasmids herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available plasmids in accord with published procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled artisan.

"Digestion" of DNA refers to catalytic cleavage of the DNA with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes used herein are commercially available and their reaction conditions, cofactors and other requirements were used as would be known to the ordinarily skilled artisan. For analytical purposes, typically 1 ög of plasmid or DNA fragment is used with about 2 units of enzyme in about 20 öl of buffer solution. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 ög of DNA are digested with 20 to 250 units of enzyme in a larger volume. Appropriate buffers and substrate amounts for particular restriction enzymes are specified by the manufacturer. Incubation times of about 1 hour at 37°C are ordinarily used, but may vary in accordance with the supplier's instructions. After digestion the reaction is electrophoresed directly on a polyacrylamide gel to isolate the desired fragment.

Size separation of the cleaved fragments is generally performed using 8 percent polyacrylamide gel described by Goeddel, D. *et al.*, Nucleic Acids Res., 8:4057 (1980), for example.

"Oligonucleotides" refers to either a single stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands which may be chemically synthesized. Such synthetic oligonucleotides may or may not have a 5' phosphate. Those that do not will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide will ligate to a fragment that has not been dephosphorylated.

"Ligation" refers to the process of forming phosphodiester bonds between two double

SUBSTITUTE SHEET (RULE 26)

-33-

stranded nucleic acid fragments (Maniatis, T., et al., Id., p. 146). Unless otherwise provided, ligation may be accomplished using known buffers and conditions with 10 units of T4 DNA ligase ("ligase") per 0.5 (g of approximately equimolar amounts of the DNA fragments to be ligated.

5 Unless otherwise stated, transformation was performed as described in the method of Sambrook, Fritsch and Maniatus, 1989. The following examples are intended to illustrate, but not to limit, the invention. While the procedures described in the examples are typical of those that can be used to carry out certain aspects of the invention, other procedures known to those skilled in the art can also be used. The following materials
10 and methods were used in carrying out the experiments described in the examples.

Example 1

Isolation, Bacterial Expression and Purification of Phytase

E.coli B genomic DNA was obtained from Sigma (Catalog # D-2001), St. Louis, New Jersey.

15 The following primers were used to PCR amplify the gene directly from the genomic DNA:

5' primer	gtttctgaattcaaggaggaatttaaATGAAAGCGATCTTAATCCCATT
3' primer	gtttctggatccTTACAAACTGCACGCCGGTAT

20 Pfu polymerase in the PCR reaction, and amplification was performed according to manufacturers protocol (Stratagene Cloning Systems, Inc., La Jolla, CA).

PCR product was purified and purified product and pQE60 vector (Qiagen) were both digested with EcoRI and BglII restriction endonucleases (New England Biolabs) according to manufacturers protocols. Overnight ligations were performed using

standard protocols to yield pQE60.

The amplified sequences were inserted in frame with the sequence encoding for the RBS. The ligation mixture was then used to transform the *E. coli* strain M15/pREP4 (Qiagen, Inc.) by electroporation. M15/pREP4 contains multiple copies of the plasmid pREP4, which expresses the *lacI* repressor and also confers kanamycin resistance (Kan^r). Plasmid DNA was isolated and confirmed by restriction analysis. Clones containing the desired constructs were grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture was used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells were grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG ("Isopropyl-B-D-thiogalacto pyranoside") was then added to a final concentration of 1 mM. IPTG induces by inactivating the *lacI* repressor, clearing the P/O leading to increased gene expression. Cells were grown an extra 3 to 4 hours. Cells were then harvested by centrifugation.

The primer sequences set out above may also be employed to isolate the target gene from the deposited material by hybridization techniques described above.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, within the scope of the appended claims, the invention may be practiced otherwise than as particularly described. It is to be understood that, while the invention has been described with reference to the above detailed description, the foregoing description is intended to illustrate, but not to limit, the scope of the invention. Other aspects, advantages, and modifications of the invention are within the scope of the following claims. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety.

What Is Claimed Is:

1. Substantially pure phytase having an amino acid sequence selected from the group consisting of SEQ ID NO:2.
2. An isolated polynucleotide sequence encoding a phytase of claim 1.
3. An isolated polynucleotide selected from the group consisting of:
 - a) SEQ ID NO:1;
 - b) SEQ ID NO:1 wherein T can also be U;
4. The polynucleotide of claim 2, wherein the polynucleotide is isolated from a prokaryote.
5. An expression vector including the polynucleotide of claim 2.
6. The vector of claim 5, wherein the vector is a plasmid.
7. The vector of claim 5, wherein the vector is a virus-derived.
8. A host cell transformed with the vector of claim 5.
9. The host cell of claim 8, wherein the cell is prokaryotic.
10. Antibodies that bind to the polypeptide of claim 1.
11. The antibodies of claim 10, wherein the antibodies are polyclonal.

-36-

12. The antibodies of claim 10, wherein the antibodies are monoclonal.
13. A method for producing an enzyme comprising growing a host cell of claim 8 under conditions which allow the expression of the nucleic acid and isolating the enzyme encoded by the nucleic acid.
14. A method for degrading phytase comprising contacting phytate with a degrading effective amount of the enzyme of claim 1.
15. A method for hydrolyzing the phospho-mono-ester bond in phytate comprising contacting an effective amount of the enzyme of claim 1 with phytate to hydrolyze the bond.
16. An animal feed composition comprising a phytase.
17. The composition of claim 16, wherein the phytase has an amino acid sequence as set forth in SEQ ID NO:2.
18. An isolated polynucleotide having at least 70% identity to a member selected from the group consisting of:
 - a) a polynucleotide encoding an enzyme comprising an amino acid sequence as set forth in SEQ ID NO:2;
 - b) a polynucleotide which is complementary to the polynucleotide of a); and
 - c) a polynucleotide comprising at least 15 bases of the polynucleotide of a) or b).
19. An oligonucleotide probe that hybridizes to a nucleic acid target region corresponding to a region selected from the nucleic acid sequence set forth in SEQ ID NO:1
20. The probe of claim 19, wherein the probe is from 10 to about 150 nucleotides in length.

-37-

21. The probe of claim 19, wherein the probe is from 10 to about 100 nucleotides in length.
22. The probe of claim 19, wherein the probe is from 10 to about 50 nucleotides in length.
23. The probe of claim 19, wherein the probe is from 10 to about 30 nucleotides in length.
24. The probe of claim 19, wherein the probe is from 10 to about 15 nucleotides in length.
25. The oligonucleotide probe of claim 19, wherein the probe comprises a segment of 10 contiguous bases which is at least 70% complementary to a target sequence of 10 contiguous nucleotides present in the target region.
26. The probe of claim 19, wherein the probe is detectably labeled.
27. The probe of claim 26, wherein the detectable label is selected from the group consisting of a radioisotope, a fluorescent compound, a bioluminescent compound, a chemiluminescent compound, a metal chelator or an enzyme.
28. The antibody of claim 10, wherein the antibody is detectably labeled.
29. An enzyme selected from the group consisting of:
 - a) an enzyme comprising an amino acid sequence which is at least 70% identical to the amino acid sequence set forth in SEQ ID NO:2; and
 - b) an enzyme which comprises at least 30 consecutive amino acid residues homologous with an enzyme of a).

(SEQ ID NO:1-nucleotide sequence and SEQ ID NO:2-amino acid sequence)



Escherichia coli B Phytase Sequence

1
 ATG AAA GCG ATC TTA ATC CCA TTT TTA TCT CTT CTG ATT CCG TTA ACC CCG
 Met Lys Ala Ile Leu Ile Pro Phe Leu Ser Leu Leu Ile Pro Leu Thr Pro
 CAA TCT GCA TTC GCT CAG AGT GAG CCG GAG CTG AAG CTG GAA AGT GTG GTG
 Gln Ser Ala Phe Ala Gln Ser Glu Pro Glu Leu Lys Leu Glu Ser Val Val
 ATT GTC AGT CGT CAT GGT GTG CGT GCT CCA ACC AAG GCC ACG CAA CTG ATG
 Ile Val Ser Arg His Gly Val Arg Ala Pro Thr Lys Ala Thr Gln Leu Met
 CAG GAT GTC ACC CCA GAC GCA TGG CCA ACC TGG CCG GTA AAA CTG GGT TGG
 Gln Asp Val Thr Pro Asp Ala Trp Pro Thr Trp Pro Val Lys Leu Gly Trp
 CTG ACA CCG CGN GGT GGT GAG CTA ATC GCC TAT CTC GGA CAT TAC CAA CGC
 Leu Thr Pro Arg Gly Gly Glu Leu Ile Ala Tyr Leu Gly His Tyr Gln Arg
 CAG CGT CTG GTA GCC GAC GGA TTG CTG GCG AAA AAG GGC TGC CCG CAG TCT
 Gln Arg Leu Val Ala Asp Gly Leu Leu Ala Lys Lys Gly Cys Pro Gln Ser
 GGT CAG GTC GCG ATT ATT GCT GAT GTC GAC GAG CGT ACC CGT AAA ACA GGC
 Gly Gln Val Ala Ile Ile Ala Asp Val Asp Glu Arg Thr Arg Lys Thr Gly
 GAA GCC TTC GCC GCC GGG CTG GCA CCT GAC TGT GCA ATA ACC GTA CAT ACC
 Glu Ala Phe Ala Ala Gly Leu Ala Pro Asp Cys Ala Ile Thr Val His Thr
 CAG GCA GAT ACG TCC AGT CCC GAT CCG TTA TTT AAT CCT CTA AAA ACT GGC
 Gln Ala Asp Thr Ser Ser Pro Asp Pro Leu Phe Asn Pro Leu Lys Thr Gly
 GTT TGC CAA CTG GAT AAC GCG AAC GTG ACT GAC GCG ATC CTC AGC AGG GCA
 Val Cys Gln Leu Asp Asn Ala Asn Val Thr Asp Ala Ile Leu Ser Arg Ala
 GGA GGG TCA ATT GCT GAC TTT ACC GGG CAT CGG CAA ACG GCG TTT CGC GAA
 Gly Gly Ser Ile Ala Asp Phe Thr Gly His Arg Gln Thr Ala Phe Arg Glu
 CTG GAA CGG GTG CTT AAT TTT CCG CAA TCA AAC TTG TGC CTT AAA CGT GAG
 Leu Glu Arg Val Leu Asn Phe Pro Gln Ser Asn Leu Cys Leu Lys Arg Glu
 AAA CAG GAC GAA AGC TGT TCA TTA ACG CAG GCA TTA CCA TCG GAA CTC AAG
 Lys Gln Asp Glu Ser Cys Ser Leu Thr Gln Ala Leu Pro Ser Glu Leu Lys
 GTG AGC GCC GAC AAT GTC TCA TTA ACC GGT GCG GTA AGC CTC GCA TCA ATG
 Val Ser Ala Asp Asn Val Ser Leu Thr Gly Ala Val Ser Leu Ala Ser Met
 CTG ACG GAG ATA TTT CTC CTG CAA CAA GCA CAG GGA ATG CCG GAG CCG GGG
 Leu Thr Glu Ile Phe Leu Leu Gln Gln Ala Gln Gly Met Pro Glu Pro Gly
 TGG GGA AGG ATC ACC GAT TCA CAC CAG TGG AAC ACC TTG CTA AGT TTG CAT
 Trp Gly Arg Ile Thr Asp Ser His Gln Trp Asn Thr Leu Leu Ser Leu His
 AAC GCG CAA TTT TAT TTG CTA CAA CGC ACG CCA GAG GTT GCC CGC AGC CGC
 Asn Ala Gln Phe Tyr Leu Leu Gln Arg Thr Pro Glu Val Ala Arg Ser Arg
 GCC ACC CCG TTA TTG GAT TTG ATC ATG GCA GCG TTG ACG CCC CAT CCA CCG

FIGURE 1 CONT'D

Ala Thr Pro Leu Leu Asp Leu Ile Met Ala Ala Leu Thr Pro His Pro Pro
 CAA AAA CAG GCG TAT GGT GTG ACA TTA CCC ACT TCA GTA CTG TTT ATT GCC
 Gln Lys Gln Ala Tyr Gly Val Thr Leu Pro Thr Ser Val Leu Phe Ile Ala
 GGA CAC GAT ACT AAT CTG GCA AAT CTC GGC GGC GCA CTG GAG CTC AAC TGG
 Gly His Asp Thr Asn Leu Ala Asn Leu Gly Gly Ala Leu Glu Leu Asn Trp
 ACG CTT CCC GGT CAG CCG GAT AAC ACG CCG CCA GGT GGT GAA CTG GTG TTT
 Thr Leu Pro Gly Gln Pro Asp Asn Thr Pro Pro Gly Gly Glu Leu Val Phe
 GAA CGC TGG CGT CGG CTA AGC GAT AAC AGC CAG TGG ATT CAG GTT TCG CTG
 Glu Arg Trp Arg Arg Leu Ser Asp Asn Ser Gln Trp Ile Gln Val Ser Leu
 GTC TTC CAG ACT TTA CAG CAG ATG CGT GAT AAA ACG CCG CTG TCA TTA AAT
 Val Phe Gln Thr Leu Gln Gln Met Arg Asp Lys Thr Pro Leu Ser Leu Asn
 ACG CCG CCC GGA GAG GTG AAA CTG ACC CTG GCA GGA TGT GAA GAG CGA AAT
 Thr Pro Pro Gly Glu Val Lys Leu Thr Leu Ala Gly Cys Glu Glu Arg Asn
 GCG CAG GGC ATG TGT TCG TTG GCA GGT TTT ACG CAA ATC GTG AAT GAA GCA
 Ala Gln Gly Met Cys Ser Leu Ala Gly Phe Thr Gln Ile Val Asn Glu Ala
 CGC ATA CCG GCG TGC AGT TTG AGA TCT CAT CAC CAT CAC CAT CAC TAA 1323
 Arg Ile Pro Ala Cys Ser Leu Arg Ser His His His His His His End

FIGURE 2
pH/Temperature Profile and Stability



-1-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: KRETZ
- (ii) TITLE OF INVENTION: NOVEL PHYTASE
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Fish & Richardson P.C.
 - (B) STREET: 4225 Executive Square, Suite 1400
 - (C) CITY: La Jolla
 - (D) STATE: California
 - (E) COUNTRY: US
 - (F) ZIP: 92037
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/910,798
 - (B) FILING DATE: August 13, 1997
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: HAILE, PH.D., LISA A.
 - (B) REGISTRATION NUMBER: 38,347
 - (C) REFERENCE/DOCKET NUMBER: 09010/029001
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 619/678-5070
 - (B) TELEFAX: 619/678-5099

-2-

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1272 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vii) IMMEDIATE SOURCE:

- (B) CLONE: PHYTASE

(ix) FEATURE:

- (A) NAME/KEY:
- (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG AAA GCG ATC TTA ATC CCA TTT TTA TCT CTT CTG ATT CCG TTA ACC	48
Met Lys Ala Ile Leu Ile Pro Phe Leu Ser Leu Leu Ile Pro Leu Thr	
1 5 10 15	
CCG CAA TCT GCA TTC GCT CAG AGT GAG CCG GAG CTG AAG CTG GAA AGT	96
Pro Gln Ser Ala Phe Ala Gln Ser Glu Pro Glu Leu Lys Leu Glu Ser	
20 25 30	
GTG GTG ATT GTC AGT CGT CAT GGT GTG CGT GCT CCA ACC AAG GCC ACG	144
Val Val Ile Val Ser Arg His Gly Val Arg Ala Pro Thr Lys Ala Thr	
35 40 45	
CAA CTG ATG CAG GAT GTC ACC CCA GAC GCA TGG CCA ACC TGG CCG GTA	192
Gln Leu Met Gln Asp Val Thr Pro Asp Ala Trp Pro Thr Trp Pro Val	
50 55 60	
AAA CTG GGT TGG CTG ACA CCG CGN GGT GGT GAG CTA ATC GCC TAT CTC	240
Lys Leu Gly Trp Leu Thr Pro Arg Gly Gly Glu Leu Ile Ala Tyr Leu	
65 70 75 80	
GGA CAT TAC CAA CGC CAG CGT CTG GTA GCC GAC GGA TTG CTG GCG AAA	288
Gly His Tyr Gln Arg Gln Arg Leu Val Ala Asp Gly Leu Leu Ala Lys	
85 90 95	
AAG GGC TGC CCG CAG TCT GGT CAG GTC GCG ATT ATT GCT GAT GTC GAC	336
Lys Gly Cys Pro Gln Ser Gly Gln Val Ala Ile Ile Ala Asp Val Asp	
100 105 110	
GAG CGT ACC CGT AAA ACA GGC CAG GCA GAT ACG TCC AGT CCC GAT CCG	384
Glu Arg Thr Arg Lys Thr Gly Gln Ala Asp Thr Ser Ser Pro Asp Pro	
115 120 125	
TTA TTT AAT CCT CTA AAA ACT GGC GTT TGC CAA CTG GAT AAC GCG AAC	432
Leu Phe Asn Pro Leu Lys Thr Gly Val Cys Gln Leu Asp Asn Ala Asn	
130 135 140	
GTG ACT GAC GCG ATC CTC AGC AGG GCA GGA GGG TCA ATT GCT GAC TTT	480
Val Thr Asp Ala Ile Leu Ser Arg Ala Gly Gly Ser Ile Ala Asp Phe	
145 150 155 160	
ACC GGG CAT CGG CAA ACG GCG TTT CGC GAA CTG GAA CGG GTG CTT AAT	528
Thr Gly His Arg Gln Thr Ala Phe Arg Glu Leu Glu Arg Val Leu Asn	
165 170 175	

SUBSTITUTE SHEET (RULE 26)

-3-

TTT CCG CAA TCA AAC TTG TGC CTT AAA CGT GAG AAA CAG GAC GAA AGC	576
Phe Pro Gln Ser Asn Leu Cys Leu Lys Arg Glu Lys Gln Asp Glu Ser	
180 185 190	
TGT TCA TTA ACG CAG GCA TTA CCA TCG GAA CTC AAG GTG AGC GCC GAC	624
Cys Ser Leu Thr Gln Ala Leu Pro Ser Glu Leu Lys Val Ser Ala Asp	
195 200 205	
AAT GTC TCA TTA ACC GGT GCG GTA AGC CTC GCA TCA ATG CTG ACG GAG	672
Asn Val Ser Leu Thr Gly Ala Val Ser Leu Ala Ser Met Leu Thr Glu	
210 215 220	
ATA TTT CTC CTG CAA CAA GCA CAG GGA ATG CCG GAG CCG GGG TGG GGA	720
Ile Phe Leu Leu Gln Gln Ala Gln Gly Met Pro Glu Pro Gly Trp Gly	
225 230 235 240	
AGG ATC ACC GAT TCA CAC CAG TGG AAC ACC TTG CTA AGT TTG CAT AAC	768
Arg Ile Thr Asp Ser His Gln Trp Asn Thr Leu Leu Ser Leu His Asn	
245 250 255	
GCG CAA TTT TAT TTG CTA CAA CGC ACG CCA GAG GTT GCC CGC AGC CGC	816
Ala Gln Phe Tyr Leu Leu Gln Arg Thr Pro Glu Val Ala Arg Ser Arg	
260 265 270	
GCC ACC CCG TTA TTG GAT TTG ATC ATG GCA GCG TTG ACG CCC CAT CCA	864
Ala Thr Pro Leu Leu Asp Leu Ile Met Ala Ala Leu Thr Pro His Pro	
275 280 285	
CCG CAA AAA CAG GCG TAT GGT GTG ACA TTA CCC ACT TCA GTA CTG TTT	912
Pro Gln Lys Gln Ala Tyr Gly Val Thr Leu Pro Thr Ser Val Leu Phe	
290 295 300	
ATT GCC GGA CAC GAT ACT AAT CTG GCA AAT CTC GGC GGC GCA CTG GAG	960
Ile Ala Gly His Asp Thr Asn Leu Ala Asn Leu Gly Gly Ala Leu Glu	
305 310 315 320	
CTC AAC TGG ACG CTT CCC GGT CAG CCG GAT AAC ACG CCG CCA GGT GGT	1008
Leu Asn Trp Thr Leu Pro Gly Gln Pro Asp Asn Thr Pro Pro Gly Gly	
325 330 335	
GAA CTG GTG TTT GAA CGC TGG CGT CGG CTA AGC GAT AAC AGC CAG TGG	1056
Glu Leu Val Phe Glu Arg Trp Arg Arg Leu Ser Asp Asn Ser Gln Trp	
340 345 350	
ATT CAG GTT TCG CTG GTC TTC CAG ACT TTA CAG CAG ATG CGT GAT AAA	1104
Ile Gln Val Ser Leu Val Phe Gln Thr Leu Gln Gln Met Arg Asp Lys	
355 360 365	
ACG CCG CTG TCA TTA AAT ACG CCG CCC GGA GAG GTG AAA CTG ACC CTG	1152
Thr Pro Leu Ser Leu Asn Thr Pro Pro Gly Glu Val Lys Leu Thr Leu	
370 375 380	
GCA GGA TGT GAA GAG CGA AAT GCG CAG GGC ATG TGT TCG TTG GCA GGT	1200
Ala Gly Cys Glu Glu Arg Asn Ala Gln Gly Met Cys Ser Leu Ala Gly	
385 390 395 400	
TTT ACG CAA ATC GTG AAT GAA GCA CGC ATA CCG GCG TGC AGT TTG AGA	1248
Phe Thr Gln Ile Val Asn Glu Ala Arg Ile Pro Ala Cys Ser Leu Arg	
405 410 415	
TCT CAT CAC CAT CAC CAT CAC TAA	1272
Ser His His His His His His	
420	

SUBSTITUTE SHEET (RULE 26)

(A) LENGTH: 423 amino acids

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEOUENCE DESCRIPTION: SEO ID NO:2:

SUBSTITUTE SHEET (RULE 26)

-5-

420

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 49 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (ix) FEATURE:
 - (A) NAME/KEY:
 - (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GTTTCTGAAT TCAAGGAGGA ATTTAAATGA AAGCGATCTT AATCCCATT

49

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (ix) FEATURE:
 - (A) NAME/KEY:
 - (B) LOCATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GTTTCTGGAT CCTTACAAAC TGCACGCCGG TAT

33

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16728

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 426/53, 656; 435/6, 196, 325, 320.1, 252.3; 530/387.9, 388.26; 536/23.1, 23.2, 24.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	DASSA, J. et al. The Complete Nucleotide Sequence of the Escherichia coli Gene appA Reveals Significant Homology between pH 2.5 Acid Phosphatase and Glucose-1-Phosphatase. J. Bacteriol. September 1990, Vol. 172, No. 9, pages 5497-5500, especially page 5498.	18 ----- 10-12, 31
X	US 5,436,156 A (VAN GORCOM et al.) 25 July 1995, col. 5, lines 10-34.	16
X	Database GenBank on STN. DOEBELI, H. et al. 'Oligonucleotide sequence (adaptor 9) from patent EP0282042'. GenBank. January 1994. Accession No. A02249.	19-23, 25 ----- 26,27



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

13 OCTOBER 1998

Date of mailing of the international search report

23 OCT 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231
Facsimile No. (703) 305-3230Authorized officer
PETER TUNG

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16728

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6): A23B 7/10; A23J 1/00; C07K 16/12; C07H 15/11, 21/02, 21/04; C12N 9/16, 15/85, 15/63, 1/20; C12Q 1/68

A. CLASSIFICATION OF SUBJECT MATTER:

US CL : 426/53, 656; 435/6, 196, 325, 320.1, 252.3; 530/387.9, 388.26; 536/23.1, 23.2, 24.3

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

MasPar Search (DNA and Protein sequences), APS, MEDLINE, CAPLUS, CANCERLIT, BIOSIS, EMBASE, WPIDS, DISSABS

search terms: Keith Kretz, phytase, antibody, feed, E. coli, bacteria

THIS PAGE BLANK (USPTO)
